Rozprawa doktorska

Efektywność technik modyfikujących procesy biodegradacji zanieczyszczeń ropopochodnych w gruncie z terenu dołu urobkowego

Autor:
mgr Katarzyna Wojtowicz

Promotor:
dr hab. inż. Teresa Steliga, prof. INiG-PIB



Politechnika Śląska

Dyscyplina naukowa: Inżynieria Środowiska, Górnictwo i Energetyka

Gliwice, 2026

Autor:
mgr Katarzyna Wojtowicz
Zakład Technologii Eksploatacji Płynów Złożowych
Instytut Nafty i Gazu – Państwowy Instytut Badawczy
ul. Lubicz 25A, 31-503 Kraków
e-mail: katarzyna.wojtowicz@inig.pl

Promotor:
dr hab. inż. Teresa Steliga, prof. INiG-PIB
Kierownik Zakładu Technologii Eksploatacji Płynów Złożowych
Instytut Nafty i Gazu – Państwowy Instytut Badawczy
ul. Lubicz 25A, 31-503 Kraków
e-mail: teresa.steliga@inig.pl

Tytuł w języku angielskim:

Effectiveness of Techniques Enhancing the Biodegradation of Petroleum Hydrocarbon Contaminants in Soil from a Waste Pit Area

©Copyright 2026 by Katarzyna Wojtowicz

Podziękowania

Składam serdeczne podziękowania wszystkim osobom, które przyczyniły się do powstania niniejszej pracy doktorskiej oraz wsparły mnie w jej realizacji.

W szczególności pragnę wyrazić głęboką wdzięczność Pani Promotor, Pani prof. dr hab. Teresie Stelidze, za opiekę naukową, nieocenione wsparcie merytoryczne, cierpliwość oraz czas poświęcony na konsultacje i dyskusje, które miały kluczowe znaczenie dla kształtu tej pracy.

Szczególne podziękowania kieruję również do Kierownictwa Instytutu Nafty i Gazu – Państwowego Instytutu Badawczego, a zwłaszcza do Pana Dyrektora prof. dr hab. Piotra Kaszy, za życzliwość, wsparcie oraz stworzenie warunków sprzyjających rozwojowi naukowemu i realizacji celów badawczych i zawodowych.

Dziękuję wszystkim Pracownikom, Koleżankom i Kolegom z INiG-PIB za pomoc, współpracę oraz życzliwą atmosferę, która towarzyszyła mi na różnych etapach realizacji pracy.

Wyrazy wdzięczności kieruję także do pracowników Politechniki Śląskiej, a w szczególności do Pana prof. dr hab. Krzysztofa Labusa oraz Pani mgr inż. Izabeli Herman za pomoc w sprawach organizacyjnych i formalnych.

Na koniec pragnę podziękować Rodzicom za nieustanne wsparcie, zrozumienie i pomoc, bez których ukończenie tej pracy nie byłoby możliwe.

Spis treści

1. Wstęp 11

1.1. Zanieczyszczenia środowiska glebowego węglowodorami ropopochodnymi 11

1.2. Charakterystyka zanieczyszczeń ropopochodnych 12

1.3. Strategie prowadzenia remediacji gruntów zanieczyszczonych węglowodorami ropopochodnymi 14

1.4. Biodegradacja węglowodorów ropopochodnych 16

1.4.1. Czynniki wpływające na efektywność biodegradacji 16

1.4.2. Metody wspomagania procesu biodegradacji 23

1.5. Charakterystyka gleb z dołów urobkowych 33

1.6. Motywacja, hipoteza i cele pracy 38

1.7. Zakres pracy 40

2. Materiały i metody 43

2.1. Materiały badawcze 43

2.2. Metody badawcze 46

2.2.1. Analizy respirometryczne 48

2.2.2. Analizy chromatograficzne 49

2.2.3. Ocena ekotoksykologiczna 51

2.2.4. Analizy mikrobiologiczne 56

2.2.5. Analiza statystyczna 60

3. Wpływ γ-PGA na efektywność biodegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego wspomaganej bioaugmentacją- Artykuł I 61

3.1. Omówienie wyników zaprezentowanych w artykule I 62

3.2 Podsumowanie artykułu I 68

4. Rola biosurfaktantów w poprawie efektywności biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego – Artykuł II 70

4.1. Omówienie wyników zaprezentowanych w artykule II 70

4.2 Podsumowanie artykułu II 75

5. Wpływ wzbogacenia biopreparatu bakteryjnego w wybrane autochtoniczne szczepy grzybów na efektywność biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego oraz zastosowania fitoremediacji w celu jej głębszego doczyszczenia – Artykuły III i IV 78

5.1. Omówienie wyników badań biodegradacji zaprezentowanych w artykule III 79

5.2. Omówienie wyników fitoremediacji wspomaganej bioaugmentacją zaprezentowanych w artykułach III i IV 82

5.3. Podsumowanie artykułów III i IV 89

6. Zastosowanie roślin dziko rosnących w celu zwiększenia efektywności fitoremediacji gleb zanieczyszczonych węglowodorami ropopochodnymi – Artykuł V 92

6.1. Omówienie wyników zaprezentowanych w artykule V 93

6.2. Podsumowanie artykułu V 100

7. Wpływ nanomateriałów na efektywność bioremediacji gleby z dołu urobkowego – Artykuł VI . 104

7.1. Omówienie wyników badań biodegradacji zaprezentowanych w artykule VI 105

7.2. Omówienie wyników badań fitoremediacji wspomaganej bioaugmentacją i nanomateriałami zaprezentowanych w artykule VI 108

7.3 Podsumowanie artykułu VI 112

8. Efektywności biodegradacji węglowodorów ropopochodnych przy współwystępowaniu skażenia polichlorowanymi bifenylami (PCB) – Artykuł VII 115

8.1. Omówienie wyników zaprezentowanych w artykule VII 115

8.2. Podsumowanie artykułu VII 121

9. Podsumowanie i wnioski 123

Bibliografia 129

Streszczenie 151

Załączniki 156

Artykuł I 157

Artykuł II 187

Artykuły III 215

Artykuły IV 246

Artykuł V 269

Artykuł VI 295

Artykuł VII 230

Lista publikacji:

Niniejsza rozprawa doktorska została przygotowana w oparciu o siedem powiązanych tematycznie publikacji naukowych, zamieszczonych w załącznikach do pracy. Poniżej przedstawiono szczegółowe dane bibliograficzne dotyczące poszczególnych artykułów. Zakres wkładu współautorów w przygotowanie każdej z publikacji określono w oświadczeniach autorskich dołączonych do rozprawy.

I. **Katarzyna Wojtowicz**, T. Steliga, P., Kapusta, J. Brzeszcz, T. Skalski, *Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil*, Materials, Volume 15(2), 400, 2022.
DOI: 10.3390/ma15020400
(**IF**$_{2022}$ **= 3,4**); 28 cytowań

II. **Katarzyna Wojtowicz**, T. Steliga, T. Skalski, P., Kapusta, *Influence of Biosurfactants on the Efficiency of Petroleum Hydrocarbons Biodegradation in Soil*, Sustainability, Volume 17(14), 6520, 2025.
DOI: 10.3390/su17146520
(**IF**$_{2025}$ **= 3,3**);5 cytowań

III. **Katarzyna Wojtowicz**, T. Steliga, P., Kapusta, J. Brzeszcz, *Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (Zea mays)*, Molecules, Volume 28(16), 6104, 2023.
DOI: 10.3390/molecules28166104
(**IF**$_{2023}$ **= 4,2**); 29 cytowań

IV. **Katarzyna Wojtowicz**, T. Steliga, P., Kapusta, *Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using Echinacea purpurea*, Applied Sciences (Basel), Volume 13(24), 13077, 2023.
DOI: 10.3390/app132413077
(**IF**$_{2023}$ **= 2,5**); 14 cytowań

V. **Katarzyna Wojtowicz**, T. Steliga, J. Brzeszcz, J. Fyda, T. Skalski, P., Kapusta, *Phytoremediation of soil contaminated with petroleum hydrocarbons using the wild plants Scirpus sylvaticus and Cirsium oleraceum, supported by bioaugmentation*, International Biodeterioration & Biodegradation, Volume 206, 106217, 2026.
DOI: 10.1016/j.ibiod.2025.106217
(**IF**$_{2025}$ **= 4,1**); 6 cytowań

VI. **Katarzyna Wojtowicz**, T. Steliga, P., Kapusta, J. Brzeszcz, *The Role of Graphene Oxide and Zinc Oxide Nanoparticles in Enhancing the Effectiveness of Phytoremediation of Petroleum Hydrocarbon-Contaminated Soils Using Lolium perenne*, Molecules, Volume 31(5), 890, 2026. DOI: 10.3390/molecules31050890
(**IF**$_{2026}$ **= 4,6**); 0 cytowań

VII. T. Steliga, **Katarzyna Wojtowicz**, P., Kapusta, J. Brzeszcz, *Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum*

Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture, Molecules, Volume 25(3), 709, 2020.
DOI: 10.3390/molecules25030709
(**IF_{2020} = 4,927**); 69 cytowań

Publikacje wspomagające:

Katarzyna Wojtowicz, T. Steliga, *Wpływ mieszaniny biosurfaktantów na efektywność bioremediacji gruntu zanieczyszczonego węglowodorami ropopochodnymi*, 80(5), 269-278, 2024. DOI: 10.18668/ng.2024.05.02

Katarzyna Wojtowicz, *Opracowanie metodyki oznaczania BTEX w próbkach gleb z wykorzystaniem chromatografii gazowej z przystawką headspace*, Nafta-Gaz, 74(3), 201-207, 2018. DOI: 10.18668/NG.2018.03.03

Katarzyna Wojtowicz, *Opracowanie metodyki oznaczania polichlorowanych bifenyli w próbkach gleb*, Nafta-Gaz, 75(7), 420-429, 2019. DOI: 10.18668/NG.2019.07.06

Katarzyna Wojtowicz, Piotr Jakubowicz, *Opracowanie metodyki oznaczania WWA w próbkach gleb z wykorzystaniem chromatografii cieczowej HPLC,* Nafta-Gaz, 78(2), 141-153, 2022, DOI: 10.18668/ng.2022.02.06

T. Steliga, P. Jakubowicz, P. Kapusta, **Katarzyna Wojtowicz**, *Efektywność biodegradacji zanieczyszczeń ropopochodnych w zestarzałym odpadzie z dołu urobkowego*, Nafta-Gaz, 74(10), 752-758, 2018. DOI: 10.18668/NG.2018.10.07

T. Steliga, P. Jakubowicz, **Katarzyna Wojtowicz**, D. Kluk, *Zastosowanie testów toksykologicznych w przemyśle naftowym*, Nafta-Gaz, 74(9), 684-689, 2018. DOI: 10.18668/NG.2018.09.07

Wkład autora rozprawy w przygotowanie każdego z artykułów załączonych do rozprawy doktorskiej był następujący:

I. Wkład **Katarzyny Wojtowicz**, zgodnie z oświadczeniem autorskim wynosił **40**% i obejmował: opracowanie koncepcji artykułu, pozyskanie finansowania, analizę formalną, przeprowadzenie i opracowanie wyników badań, przygotowanie metodologii, opracowanie manuskryptu oraz wykonanie przeglądu literaturowego, a także przegląd i redakcję tekstu.

II. Wkład **Katarzyny Wojtowicz**, zgodnie z oświadczeniem autorskim wynosił **70**% i obejmował: opracowanie koncepcji artykułu, przygotowanie metodologii, walidację wyników, analizę formalną, przeprowadzenie i opracowanie wyników badań, opracowanie manuskryptu oraz wykonanie przeglądu literaturowego, a także przegląd i redakcję tekstu.

III. Wkład **Katarzyny Wojtowicz**, zgodnie z oświadczeniem autorskim wynosił **48**% i obejmował: opracowanie koncepcji artykułu, przygotowanie oprogramowania, walidację wyników, analizę formalną, przeprowadzenie i opracowanie wyników badań, a także opracowanie manuskryptu oraz wykonanie przeglądu literaturowego.

IV. Wkład **Katarzyny Wojtowicz**, zgodnie z oświadczeniem autorskim wynosił **50**% i obejmował: opracowanie koncepcji artykułu, przygotowanie oprogramowania, walidację wyników, analizę formalną, przeprowadzenie i opracowanie wyników badań, a także opracowanie manuskryptu oraz wykonanie przeglądu literaturowego.

V. Wkład **Katarzyny Wojtowicz**, zgodnie z oświadczeniem autorskim wynosił **50**% i obejmował: opracowanie koncepcji artykułu, analizę formalną, przeprowadzenie i opracowanie wyników badań, przygotowanie metodologii, opracowanie manuskryptu oraz wykonanie przeglądu literaturowego, a także przegląd i redakcję tekstu.

VI. Wkład **Katarzyny Wojtowicz**, zgodnie z oświadczeniem autorskim wynosił **40**% i obejmował: opracowanie koncepcji artykułu, przygotowanie metodologii, walidację wyników, analizę formalną, przeprowadzenie i opracowanie wyników badań, opracowanie manuskryptu oraz wykonanie przeglądu literaturowego, a także przegląd i redakcję tekstu.

VII. Wkład **Katarzyny Wojtowicz**, zgodnie z oświadczeniem autorskim wynosił **25**% i obejmował: opracowanie koncepcji artykułu, analizę formalną, przeprowadzenie i opracowanie wyników badań, przygotowanie metodologii, a także opracowanie manuskryptu oraz wykonanie przeglądu literaturowego.

Terminologia

A	Antracen (*z ang. Anthracene*)
Ac	Acenaften (*z ang. Acenaphthene*)
Ames	Test mutagenności Amesa (*z ang. Ames Test*)
BaA	Benzo(a)antracen (*z ang. Benzo(a)anthracene*)
BaP	Benzo(a)piren (*z ang. Benzo(a)pyrene*)
BbF	Benzo(b)fluoranten (*z ang. Benzo(b)fluoranthene*)
BghiP	Benzo(g,h,i)perylen (*z ang. Benzo(g,h,i)perylene*)
BkF	Benzo(k)fluoranten (*z ang. Benzo(k)fluoranthene*)
BTEX	Benzen, toluen, etylobenzen, ksyleny *(z ang. Benzene, Toluene, Ethylbenzene, Xylenes)*
CFU	Jednostki tworzące kolonie (*z ang. Colony Forming Units*)
CH	Chryzen (*z ang. Chrysene*)
DahA	Dibenzo(a,h)antracen (*z ang. Dibenzo(a,h)anthracene*)
DNA	Kwas deoksyrybonukleinowy (*z ang. Deoxyribonucleic Acid*)
EC_{50}	Stężenie efektywne powodujące 50% efektu (*z ang. Effective Concentration 50%*)
ECD	Detektor wychwytu elektronów (*z ang. Electron Capture Detector*)
Metoda *ex situ*	Metoda prowadzona poza miejscem skażenia *(z ang. ex situ)*
Metoda *in situ*	Metoda prowadzona w miejscu skażenia *(z ang. in situ)*
F	Fluoranten (*z ang. Fluoranthene*)
FID	Detektor płomieniowo-jonizacyjny (*z ang. Flame Ionization Detector*)
FLD	Detektor fluorescencyjny (*z ang. Fluorescence Detector*)
Fluo	Fluoren (*z ang. Fluorene*)
G-X	Oznaczenia próbek gleby z dołów urobkowego
GC	Chromatografia gazowa (*z ang. Gas Chromatography*)
GC-HS	Chromatografia gazowa z przystawką headspace (*z ang. Gas Chromatography–Headspace)*
GO-NP	Nanocząstki tlenku grafenu *(z ang. Graphene Oxide Nanoparticles)*
HPLC	Wysokosprawna chromatografia cieczowa *(z ang. High Performance Liquid Chromatography)*
Ind123P	Indeno(1,2,3-c,d)piren *(z ang. Indeno(1,2,3-c,d)pyrene)*
INiG-PIB	Instytut Nafty i Gazu – Państwowy Instytut Badawczy *(z ang. Oil and Gas Institute – National Research Institute)*
MALDI-TOF MS	Spektrometria mas z desorpcją/jonizacją laserową wspomaganą matrycą z analizatorem czasu przelotu *(z ang. Matrix-Assisted Laser Desorption/Ionization Time-of-Flight Mass Spectrometry)*
MARA	Test oceny ryzyka mikrobiologicznego *(z ang. Microbial Assay for Risk Assessment)*
n-C_x	n-alkany o różnej długości łańcucha węglowego, gdzie x oznacza liczbę atomów węgla w cząsteczce *(z ang. n-alkanes, where x is the carbon number)*
N	Naftalen *(z ang. Naphthalene)*
NGS	Sekwencjonowanie nowej generacji *(z ang. Next-Generation Sequencing)*
P	Pristan *(z ang. Pristane)*

PAH	Wielopierścieniowe węglowodory aromatyczne *(z ang. Polycyclic Aromatic Hydrocarbons)*
PCB	Polichlorowane bifenyle *(z ang. Polychlorinated Biphenyls)*
Phen	Fenantren *(z ang. Phenanthrene)*
Ph	Fitan *(z ang. Phytane)*
Pr	Piren *(z ang. Pyrene)*
QuEChERS	Metoda ekstrakcji *(z ang. Quick, Easy, Cheap, Effective, Rugged and Safe)*
Analiza RA	Analiza rarefakcji *(z ang. Rarefaction Analysis)*
Analiza RDA	Analiza redundancji *(z ang. Redundancy Analysis)*
Soil DW	Gleba z dołu urobkowego *(z ang. Drill Waste Soil)*
Soil OS	Gleba z miejsca wycieku ropy naftowej *(z ang. Oil Spill Soil)*
TPH	Całkowita zawartość węglowodorów ropopochodnych *(z ang. Total Petroleum Hydrocarbons)*
TU	Wpółczynnik toksyczności *(z ang. Toxicity Unit)*
UV-VIS	Spektrofotometria w zakresie ultrafioletu i światła widzialnego *(z ang. Ultraviolet–Visible Spectroscopy)*
WWA	Wielopierścieniowe węglowodory aromatyczne *(z ang. PAHs Polycyclic Aromatic Hydrocarbons)*
ZnO-NP	Nanocząstki tlenku cynku *(z ang. Zinc Oxide Nanoparticles)*
γ-PGA	Kwas γ-poliglutaminowy *(z ang. gamma-Polyglutamic Acid)*

1. Wstęp

1.1. Zanieczyszczenia środowiska glebowego węglowodorami ropopochodnymi


W ostatnich dziesięcioleciach obserwuje się znaczące zwiększenie presji antropogenicznej na środowisko przyrodnicze, wynikające z postępującego rozwoju przemysłu, wzrostu liczby ludności oraz rosnącego zapotrzebowania na energię i produkty petrochemiczne, co może prowadzić do trwałych zmian w funkcjonowaniu ekosystemów (Mekonnen et al., 2024; Ossai et al., 2020; Singha and Pandey, 2021). Wśród elementów środowiska szczególnie podatnych na zanieczyszczenie różnego typu substancjami chemicznymi, zarówno pochodzenia naturalnego jak i antropogenicznego na pierwszy plan wysuwa się gleba, która jak wskazują dostępne źródła literaturowe może pełnić podwójną funkcje na terenach o podwyższonym stopniu narażenia na zanieczyszczenia środowiskowe. Z jednej strony może pełnić rolę naturalnej bariery ograniczającej migrację zanieczyszczeń do innych części ekosystemu (wody gruntowe, powietrze czy organizmy żywe), z drugiej natomiast źródła wtórnej emisji substancji toksycznych, co ma miejsce w przypadku zestarzałych gleb z terenów dołów urobkowych.

Zanieczyszczenia gleb dzieli się na dwie główne grupy, tj. zanieczyszczenia nieorganiczne oraz organiczne. Wśród zanieczyszczeń nieorganicznych najczęściej wymienia się metale ciężkie oraz związki mineralne, takie jak azotany, fosforany oraz nadmiar soli mineralnych, których obecność wiąże się najczęściej z działalnością rolniczą i przemysłową. Z kolei do najpowszechniejszych zanieczyszczeń organicznych należą substancje ropopochodne, w skład których wchodzą węglowodory alifatyczne i aromatyczne, w tym związki z grupy BTEX oraz wielopierścieniowe węglowodory aromatyczne (WWA) odznaczające się dużą trwałością w środowisku glebowym, polichlorowane bifenyle (PCB) oraz pozostałości środków ochrony roślin. W licznych badaniach naukowych wskazuje się, że nagromadzenie szkodliwych substancji w matrycy glebowej, będące wynikiem wieloletniej działalności przemysłowej, górniczej i rolniczej, może prowadzić do stopniowego zaburzania równowagi biologicznej ekosystemów glebowych oraz ograniczenia ich naturalnej aktywności mikrobiologicznej (Steliga, 2009). Konsekwencją tych procesów może być stopniowe obniżanie żyzności gleb oraz utrata ich właściwości użytkowych. Ponadto omawiając problem zanieczyszczenia gleb różnego rodzaju ksenobiotykami, w szczególności węglowodorami ropopochodnymi nie sposób przejść obojętnie wobec ryzyka negatywnego oddziaływania tych substancji na zdrowie ludzi i zwierząt, co w warunkach wieloletniego oddziaływania może prowadzić do rozwoju chorób i zaburzeń fizjologicznych, nierzadko prowadzących do śmierci (Daâssi & Qabil Almaghribi, 2022).

W związku z powyższym, w obecnych czasach coraz większy nacisk kładzie się na rozwój i wdrażanie skutecznych metod oczyszczania zanieczyszczonych gruntów, umożliwiających przywrócenie ich właściwości użytkowych oraz na działania zmierzające do ograniczenia dalszej degradacji środowiska naturalnego w wyniku działalności górniczej i przemysłowe.

1.2. Charakterystyka zanieczyszczeń ropopochodnych

Węglowodory ropopochodne stanowią złożoną i niejednorodną mieszaninę związków chemicznych, którą klasyfikuje się cztery główne grupy: węglowodory alifatyczne, węglowodory aromatyczne, asfalteny i żywice (Pondolfo et al., 2023) (Rys. 1). Węglowodory alifatyczne, określane również jako węglowodory łańcuchowe, to związki organiczne zbudowane wyłącznie z atomów węgla i wodoru, których struktura opiera się na prostych lub rozgałęzionych łańcuchach węglowych, rzadziej przyjmujących formę układów pierścieniowych, które pozbawione są aromatycznego układu wiązań. W obrębie tej grupy wyróżnia się związki nasycone, czyli alkany, oraz związki nienasycone, do których zalicza się alkeny i alkiny (Mekonnen et al., 2024). Węglowodory aromatyczne, podobnie jak węglowodory alifatyczne, zbudowane są wyłącznie z atomów węgla i wodoru, jednak w ich cząsteczkach występuje jeden lub więcej pierścieni aromatycznych. Węglowodory jednopierścieniowe, takie jak benzen, toluen, etylobenzen oraz izomery ksylenu określane są zbiorczo w literaturze jako węglowodory BTEX, natomiast węglowodory posiadające co najmniej dwa pierścienie aromatyczne w cząsteczce jako wielopierścieniowe węglowodory aromatyczne (WWA). Do najcięższych frakcji węglowodorów ropopochodnych o złożonej, wielkocząsteczkowej budowie zalicza się żywice oraz asfalteny. Żywice charakteryzują się większą zawartością związków polarnych i stanowią frakcję pośrednią między olejami a asfaltenami, natomiast asfalteny są frakcją najbardziej skondensowaną, nierozpuszczalną w n-alkanach i odpowiedzialną za tworzenie się osadów w instalacjach kopalnianych.



Rysunek 1 Klasyfikacja węglowodorów ropopochodnych (opracowanie własne na podstawie Mekonnen et. al. 2024 i poddane obróbce przez OpenAI)

Po przedostaniu się do gleby węglowodory ropopochodne mogą wchodzić w liczne interakcje z innymi jej komponentami, co determinuje ich dalsze zachowanie w środowisku glebowym. Związki te mogą ulegać adsorpcji na powierzchni cząstek mineralnych i materii organicznej oraz przenikać do porów glebowych, gdzie wiążą się z frakcją substancji organicznej. Jak wskazuje Siuta, już pierwszy kontakt substancji ropopochodnych z gruntem może ograniczać ich podatność na desorpcję, parowanie, biodegradację oraz ekstrakcję, co może znacząco obniżać skuteczność przyszłych procesów remediacyjnych (Siuta et al., 2001).

W zależności od swoich właściwości fizykochemicznych substancje ropopochodne mogą występować w gruncie w postaci wolnej fazy ciekłej, związków rozpuszczonych w wodzie glebowej lub składników fazy gazowej (Malina & Szczepański, 1994; Surygała et al., 2000). Dostępna literatura przedmiotu wskazuje, że udział poszczególnych form zanieczyszczeń ropopochodnych w glebie może zależeć zarówno od charakterystyki zanieczyszczenia, jak i od warunków hydrogeologicznych ośrodka gruntowego.

Na transport oraz przemiany substancji ropopochodnych mogą wpływać m.in. sorpcja, rozpuszczalność, prężność par oraz podatność na biodegradację. Malina w swojej publikacji wskazuje, że lekkie frakcje alifatyczne i aromatyczne mogą łatwo odparowywać z matrycy glebowej ze względu na wysoką prężność par, natomiast związki o niewielkim powinowactwie do ośrodka gruntowego mogą być wymywane do wód podziemnych (Malina, 2007). Cięższe frakcje węglowodorowe wykazują natomiast silniejszą sorpcję do cząstek gleby, co może ograniczać ich mobilność, ale jednocześnie zwiększać trwałość w środowisku. Procesy te mogą wpływać również na biodostępność zanieczyszczeń dla mikroorganizmów uczestniczących w ich biodegradacji, która ma istotny wpływ na efektywność procesów bioremediacyjnych (Feng et al., 2021; Sakshi et al., 2019). Warto zauważyć, że długotrwała obecność substancji ropopochodnych w glebie może prowadzić do zmian jej właściwości fizycznych, chemicznych i biologicznych. Siuta podaje, że zanieczyszczenia tego typu mogą powodować zmiany w ilości i składzie substancji organicznych w glebie, a także ograniczenie pojemności wodnej gleby oraz utrudnienie wymiany gazowej przy jednocześnie zwiększonym zapotrzebowaniu na tlen (Siuta, 2001). Ponadto wskazuje się, że zbyt wysokie stężenia węglowodorów ropopochodnych w glebie prowadzą do zaburzenia w niej właściwego stosunku węgla, azotu i fosforu, który jest niezbędny do prawidłowego funkcjonowania mikroorganizmów glebowych (Czarny et al., 2020; Dos Santos & Maranho, 2018). W konsekwencji na skutek obecności zanieczyszczeń ropopochodnych w gruncie może dochodzić do destabilizacji mikrobioty glebowej, osłabienia procesów biologicznych i biochemicznych w glebie oraz pogorszenie właściwości sorpcyjnych i jonowymiennych gruntu, często połączone z jej wtórnym zakwaszeniem (Siuta, 2001, Steliga, 2009). Ponadto ze względu na potwierdzoną toksyczność, mutagenność i rakotwórczość substancji ropopochodnych ich obecność w środowisku glebowym uznawana jest za istotne zagrożenie ekologiczne (Cui et al., 2020; Morillo et al., 2020).

1.3. Strategie prowadzenia remediacji gruntów zanieczyszczonych węglowodorami ropopochodnymi

Remediacja gruntów zanieczyszczonych węglowodorami ropopochodnymi może być prowadzona przy użyciu różnorodnych metod, które klasyfikuje się według takich kryteriów jak miejsce prowadzenia procesu oczyszczania, charakter zastosowanych technik oraz typ procesów odpowiedzialnych za eliminację lub transformację zanieczyszczeń (Rakowska et al., 2012).

Uwzględniając miejsce prowadzenia działań rekultywacyjnych, wyróżnia się dwie podstawowe strategie remediacji, do których zalicza się metody *in-situ* oraz *ex-situ*. Metody *in-situ* polegają na oczyszczaniu gruntu bezpośrednio w miejscu występowania skażenia, bez konieczności jego wydobywania i transportu poza teren zanieczyszczony (Steliga, 2009). Do najczęściej stosowanych metod *in-situ* zalicza się odżywianie gleby, biowentylację, bioekstrakcję, przemywanie oraz mobilizację zanieczyszczeń (Chaudhary & Kim, 2019; Steliga, 2009). Metody *in-situ* znajdują zastosowanie przede wszystkim do oczyszczania terenów zurbanizowanych, przemysłowych, a także pod istniejącą już infrastrukturą budowlaną. Do ich głównych zalet należą ograniczona ingerencja w strukturę oczyszczanego terenu oraz stosunkowo niskie koszty procesu, co sprawia że są one często wykorzystywane do oczyszczania terenów o dużej powierzchni.

Alternatywą są metody *ex-situ*, które opierają się na wydobyciu zanieczyszczonego gruntu i jego oczyszczaniu poza miejscem pierwotnego występowania. Takie podejście pozwala na prowadzenie procesów remediacyjnych w kontrolowanych warunkach technologicznych, co zwiększa intensywność reakcji i skuteczność eliminacji odcieków zawierających substancje szkodliwe. W ramach tej grupy metod wyróżnia się strategie *ex-situ on site*, czyli oczyszczanie wydobytej gleby na terenie skażonego obszaru, oraz *ex-situ off site*, obejmujące transport gruntu do specjalistycznych instalacji remediacyjnych. Do najczęściej stosowanych metod *ex-situ* zalicza się odżywianie gleby, kompostowanie, odmywanie oraz oczyszczanie w biopryzmach i bioreaktorach (Chaudhary & Kim,2019, Radwan et al., 2012; Steliga 2009).

Biorąc pod uwagę charakter zastosowanych technik, metody remediacji można podzielić na inżynieryjne i procesowe. Metody inżynieryjne obejmują rozwiązania techniczne, które pozwalają mechanicznie usuwać zanieczyszczoną matrycę glebową lub ograniczać migrację substancji szkodliwych poprzez zastosowanie specjalnych barier ochronnych, takich jak skimmery, materiały sorpcyjne czy bariery izolacyjne, natomiast metody procesowe wykorzystują procesy fizyczne, chemiczne lub biologiczne, prowadzące do stabilizacji, transformacji lub eliminacji zanieczyszczeń. Do najważniejszych metod procesowych należą między innymi stabilizacja i solidyfikacja zanieczyszczeń oraz różne formy obróbki termicznej.

Istotnym kryterium klasyfikacji jest również rodzaj procesów odpowiedzialnych za eliminację zanieczyszczeń. W tym ujęciu wyróżnia się metody fizyczne, chemiczne oraz biologiczne (Steliga 2009). Metody fizyczne opierają się na przemieszczaniu lub oddzielaniu zanieczyszczeń od matrycy glebowej, przy wykorzystaniu materiałów sorpcyjnych, barier izolacyjnych, systemów odprowadzania wód skażonych substancjami

ropopochodnymi, przemywania gleby lub mechanicznego usuwania skażonej roślinności. Szczególną grupą metod fizycznych są metody termiczne, obejmujące spalanie (Gan et. al. 2009; Kuppusamy et al., 2017; Zhang et al., 2024) lub ogrzewanie zanieczyszczeń, na przykład w procesach spalania termicznego czy dopalania plazmowego (Kowalska & Więch, 2004). Skuteczność metod fizycznych często bywa ograniczona jednak ze względu na szybkość wykonania, wciąż są powszechnie stosowane, szczególnie w połączeniu z innymi metodami remediacyjnymi (Arghiropol et al., 2025).

Metody chemiczne wykorzystują reakcje chemiczne do modyfikacji struktury zanieczyszczeń, co pozwala przekształcić je w formy mniej toksyczne i bardziej stabilne środowiskowo. W ramach tych metod stosuje się m.in. procesy utleniania i redukcji, zmianę odczynu środowiska glebowego czy sorpcję chemiczną. Typowe techniki remediacyjne oparte o metody chemiczne obejmują solidyfikację (Heidarzadeh et al., 2021), desorpcję próżniową (Fudryn & Kowala, 1996), metody elektrokinetyczne (Guo et al., 2020; Kuppusamy et al., 2017; Patel et al., 2020), fotolizę oraz utlenianie chemiczne (Abdel-Shafy & Mansour, 2016; Patel et al., 2020). W tym miejscu warto zwrócić uwagę na metody fizyczno-chemiczne, które łączą elementy obydwu z tych podejść, wśród których do najpopularniejszyc zaliczane jest płukanie gleby (z ang. *soil washing) (*Gan et al.,. 2009; García-Díaz et al., 2015; Kuppusamy et al., 2017).

Szczególną rolę w remediacji gruntów zanieczyszczonych węglowodorami ropopochodnymi odgrywają metody biologiczne, które wykorzystują naturalne procesy zachodzące w środowisku glebowym do przekształcania szkodliwych związków chemicznych w formy mniej toksyczne lub obojętne dla środowiska. Wśród metod biologicznych wyróżnia się przede wszystkim biodegradację mikrobiologiczną oraz fitoremediację. Biodegradacja mikrobiologiczna opiera się na zdolności mikroorganizmów glebowych, w tym bakterii, grzybów i drożdży, do produkcji enzymów odpowiedzialnych za degradację i transformację węglowodorów ropopochodnych obecnych w glebie, co istotnie wpływa na tempo naturalnego rozkładu zanieczyszczeń oraz procesy oczyszczania gleby (Guirado et al., 2023; Radhakrishnan et al., 2023; Salar et al., 2022). Fitoremediacja natomiast opiera się na wykorzystaniu roślin, które poprzez swoje naturalne procesy fizjologiczne wpływają na poprawę jakości gleby, w wyniku pobierania i akumulacji zanieczyszczeń, unieruchamiania ksenobiotyków w strefie ryzosfery lub częściowej ich transformacji przy współudziale mikroorganizmów glebowych. Dzięki temu fitoremediacja może stanowić istotne uzupełnienie procesów mikrobiologicznych i przyczynić się do eliminacji zanieczyszczeń ze środowiska glebowego (Elijah, 2022; Gupta et al., 2024; Yemele et al., 2024).

Metody biologiczne są obecnie jednymi z najczęściej wykorzystywanych metod oczyszczania gruntów zanieczyszczonych węglowodorami ropopochodnymi, gdyż charakteryzują się wysoką efektywnością, szerokim spektrum działania, a także relatywnie niskimi kosztami oczyszczania. Ograniczeniem metod biologicznych mogą być z kolei zbyt wysokie stężenia wyjściowe zanieczyszczeń w oczyszczanym gruncie, mogące wykazywać działanie hamujące względem mikroorganizmów odpowiedzialnych za rozkład zanieczyszczeń. W takich przypadkach przed rozpoczęciem procesu remediacji gruntów z wykorzystaniem metod biologicznych wskazane jest obniżenie stężenia zanieczyszczeń do poziomu tolerowanego przez mikroorganizmy glebowe np.


poprzez wstępne oczyszczenie gruntu metodami fizycznymi lub chemicznymi (Leahy & Colwell, 1990).

1.4. Biodegradacja węglowodorów ropopochodnych

Biodegradacja jest procesem biologicznym, w którym organizmy autochtoniczne przekształcają substancje toksyczne w formy mniej szkodliwe lub całkowicie neutralne dla środowiska. W przypadku zanieczyszczeń organicznych, takich jak węglowodory ropopochodne, proces ten stanowi jeden z kluczowych mechanizmów ich naturalnego usuwania z gleby i środowiska wodnego (Ghorbannezhad et al., 2022). Węglowodory ropopochodne mogą być wykorzystywane przez mikroorganizmy jako źródło węgla i energii, a ich rozkład zachodzi stopniowo w wyniku zachodzących po sobie przemian metabolicznych. Końcowymi produktami biodegradacji węglowodorów ropopochodnych w warunkach tlenowych są dwutlenek węgla i woda, natomiast w warunkach beztlenowych metan. Powstałe związki są nietoksyczne i mogą ulegać dalszym przemianom w naturalnych cyklach biogeochemicznych lub być wykorzystywane przez mikroorganizmy i rośliny, bez negatywnego wpływu na środowisko (Brzeszcz & Kaszycki, 2018; Horel & Schiewer, 2020; Laczi et al., 2020).

1.4.1. Czynniki wpływające na efektywność biodegradacji

Efektywność biodegradacji zanieczyszczeń ropopochodnych, zależy od wielu czynników, które można podzielić na trzy zasadnicze grupy do których zalicza sie: czynniki związane z zanieczyszczeniem, czynniki środowiskowe oraz czynniki biologiczne (Rys. 2).



Rysunek 2 Zestawienie czynników wpływających na efektywność biodegradacji gruntów zanieczyszczonych węglowodorami ropopochodnymi (obraz wygenerowany przez Open AI i poddany obróbce przez autora).

- ***Czynniki związane z zanieczyszczeniem***

Podatność węglowodorów na rozkład mikrobiologiczny determinowana jest przede wszystkim przez czynniki związane z zanieczyszczeniem, obejmujące rodzaj, strukturę cząsteczki, właściwości fizyczno-chemiczne zanieczyszczeń, stężenie ksenobiotyku w glebie, a także wiek zanieczyszczenia. Cooney i in. w swoim opracowaniu z 1985 roku wykazał, że poszczególne klasy węglowodorów różnią się podatnością na degradację mikrobiologiczną, którą ogólnie można uszeregować w kolejności malejącej: alkany liniowe > alkany rozgałęzione > małe węglowodory aromatyczne (BTEX) > alkany cykliczne > wielopierścieniowe węglowodory aromatyczne (WWA) (Cooney et al., 1985; Ulrici, 2001). Zależność ta jest bezpośrednio związana ze strukturą cząsteczki zanieczyszczenia. Generalnie przyjmuje się, że wraz ze wzrostem długości łańcucha węglowego węglowodorów alifatycznych lub liczby pierścieni aromatycznych w cząsteczce węglowodorów aromatycznych szybkość rozkładu maleje (Leahy & Colwell, 1990). W przypadku n-alkanów obserwuje się lekkie odstępstwo od tej reguły, gdyż najłatwiej biodegradacji ulegają węglowodory o długości łańcucha węglowego C_{10}–C_{22}, natomiast węglowodory krótkołańcuchowe ulegają biodegradacji nieznacznie gorzej. Jak wskazuje Steliga (Steliga, 2009) może to wynikać z toksycznego wpływu węglowodorów krótkołańcuchowych na mikroorganizmy degradujące zanieczyszczenia, a także mniejszej liczby gatunków mikroorganizmów zdolnych do ich efektywnego rozkładu i wykorzystania jako źródła węgla oraz energii (Steliga et al., 2012; Wu et al., 2017). W przypadku alifatycznych węglowodorów długołańcuchowych, rozgałęzionych i cyklicznych obserwuje się natomiast znacznie ograniczony stopień ich rozkładu biologicznego, co wskazuje na ich wysoką odporność na procesy biodegradacyjne. Liczne artykuły naukowe dotyczące biodegradacji węglowodorów ropopochodnych wskazują również, że zasadniczo węglowodory aromatyczne są trudniej rozkładalne przez mikroorganizmy niż krótkie alkany i alkeny, głównie ze względu na ich większą masę cząsteczkową oraz wyższą toksyczność (Abdel-Shafy & Monsour, 2016, Kalia et al., 2022), natomiast silnie skondensowane związki aromatyczne takie jak: smoły, struktury cykloparafinowe oraz asfalteny charakteryzują się najmniejszą podatnością na biodegradację (Hong et al., 2015; Kalia et al., 2022). Struktura cząsteczki węglowodoru skorelowana jest ich właściwościami fizyczno-chemicznymi. Węglowodory ropopochodne są związkami hydrofobowymi, co ogranicza ich rozpuszczalność w wodzie i utrudnia transport przez błony komórkowe mikroorganizmów. W przypadku n-alkanów hydrofobowość wzrasta wraz z długością łańcucha węglowego, natomiast w przypadku wielopierścieniowych węglowodorów aromatycznych wraz z liczbą pierścieni aromatycznych w cząsteczce. Wraz ze wzrostem hydrofobowości zanieczyszczeń spada zatem ich biodostępność dla mikroorganizmów, co z kolei prowadzi do obniżenia efektywności biodegradacji. Istnieje kilka sposobów umożliwiających poprawę biodostępności związków hydrofobowych dla mikroorganizmów, z których najważniejsze to zastosowanie surfaktantów lub biosurfaktantów, które poprzez zdolność do tworzenia emulsji pozwalają na zwiększenie powierzchni kontaktu między zanieczyszczeniem, a mikroorganizmami, wpływając tym samym na tempo biodegradacji (Hossein et al., 2022; Patowary et al., 2017; Souza et al., 2014). Kolejnym czynnikiem związanym z

zanieczyszczeniem i mającym istotny wpływ na efektywność biodegradacji jest stężenie ksenobiotyku w glebie. Wykazano, że zarówno zbyt duże, jak i zbyt małe stężenia węglowodorów w matrycy glebowej mogą znacząco spowolnić proces rozkładu biologicznego. Przyjmuje się, że wysokie stężenia węglowodorów ropopochodnych w glebie, rzędu 8–10% wag., mogą wywoływać działanie toksyczne wobec mikroorganizmów przeprowadzających biodegradację, prowadząc w konsekwencji do zahamowania ich wzrostu oraz aktywności metabolicznej (Podsiadło & Krzyśko-Łupicka, 2013; Steliga, 2009). Istnieją również doniesienia wskazujące, że poziom ten może być jeszcze niższy. Przykładowo badania Vasilyeva i in. wykazały, że zawartość ropy naftowej w glebie już powyżej 5% wag. może wywołać efekt toksyczny względem mikroorganizmów uczestniczących w procesach biodegradacji (Vasilyeva et al., 2020). Warto tutaj dodać, że toksyczne stężenie zanieczyszczeń (LC50) zależy od rodzaju ksenobiotyku, np. wartość LC50 benzenu względem bakterii wynosi około 100 mg/l, natomiast chloroformu około 670 mg/l. (Podsiadło & Krzyśko-Łupicka, 2013). Dlatego określenie toksycznego poziomu zanieczyszczeń ropopochodnych względem mikroorganizmów w przypadku gleb zanieczyszczonych mieszaniną różnych substancji ropopochodnych może być niezwykle trudne. Z drugiej strony warto zaznaczyć, że zbyt małe stężenie ksenobiotyku w glebie może hamować tempo poprzez dostarczenie mikroorganizmom niewystarczającej ilości źródła węgla, co z kolei ogranicza indukcję i syntezę enzymów uczestniczących w jego rozkładzie oraz prowadzi do opóźnienia rozpoczęcia aktywnej biodegradacji. (Podsiadło & Krzyśko-Łupicka, 2013; Towell et al., 2011). Nie mniej istotnym czynnikiem mogącym istotnie ograniczać tempo biodegradacji jest wiek zanieczyszczenia. W wyniku przedostania się węglowodorów ropopochodnych do środowiska glebowego mogą one ulegać sorpcji na powierzchni materii organicznej oraz w mikroporach gleby, gdzie z biegiem czasu ulegają przemianom fizykochemicznym i częściowej mineralizacji (Ławniczak et al., 2020). Długi kontakt hydrofobowych zanieczyszczeń ropopochodnych z matrycą glebową sprawia, że sorpcja staje się coraz trudniej odwracalna, co w konsekwencji prowadzi do obniżenia ich biodostępność. Zjawisko to, określane jest w literaturze przedmiotu jako „starzenie się” (z ang. *weathering*) zanieczyszczenia, i prowadzi do selektywnej utraty łatwo biodegradowalnych frakcji węglowodorów (np. n-alkanów i lekkich aromatów) oraz względnego wzbogacenia pozostałości w związki o wysokiej masie cząsteczkowej i strukturach opornych na biodegradację, takich jak asfalteny i żywice (Khan et al., 2018; Ławniczak et al., 2020). W konsekwencji gleby historycznie zanieczyszczone węglowodorami ropopochodnymi stają się trudniej biodegradowalne, a tempo procesów bioremediacji może ulegać istotnemu ograniczeniu (Dandie et al., 2010; Hossein et al., 2022).

- ***Czynniki środowiskowe***

Poza czynnikami związanymi z zanieczyszczeniem istotny wpływ na efektywność biodegradacji węglowodorów ropopochodnych w glebie mają czynniki środowiskowe. Jednym z najważniejszych jest temperatura, która istotnie wpływa na przebieg procesów życiowych społeczności mikroorganizmów zaangażowanych w biodegradację

zanieczyszczeń (Chandra et al., 2013; Varjani & Upasani, 2019). Dostępne dane literaturowe wskazują, że temperatura może działać na mikroorganizmy przeprowadzające biodegradację w sposób bezpośredni, poprzez przyspieszenie lub hamowanie ich wzrostu, aktywności enzymatycznej oraz wymagań pokarmowych, jak również w sposób pośredni poprzez regulację rozpuszczalności substancji w komórkach, transport jonów i właściwości osmotyczne błon komórkowych (Steliga, 2009). Wskazuje się, że zarówno zbyt niska, jak i zbyt wysoka temperatura może ograniczać metabolizm mikroorganizmów, a w skrajnych przypadkach prowadzić do ich śmierci. Das i Chandran wskazali, że dla większości bakterii biodegradacja węglowodorów najbardziej efektywnie zachodzi w przedziale temperaturowym 30–40°C w środowiskach glebowych, 20–30°C w środowiskach słodkowodnych oraz 15–20°C w środowiskach morskich (Das & Chandran, 2010). Istnieją jednak mikroorganizmy zdolne do rozkładu zanieczyszczeń ropopochodnych w bardzo niskich (mikroorganizmy psychrofilne) jak i bardzo wysokich temperaturach (mikroorganizmy termofilne) (Brakstad & Bonaunet, 2006; Feitkenhauer et al., 2003), dlatego na etapie projektowania procesu bioremediacji z wykorzystaniem techniki bioaugmentacji wskazane jest określenie warunków panujących w trakcie oczyszczania. Poza tym temperatura wpływa na właściwości zanieczyszczeń organicznych. Foeght i in. wskazali, że w przypadku węglowodorów ropopochodnych wraz ze wzrostem temperatury zwiększa się ich rozpuszczalność w wodzie, co wpływa bezpośrednio na wzrost ich biodostępności, a tym samym efektywność biodegradacji (Foght et al., 1996). Z drugiej strony, w zbyt wysokiej temperaturze obserwuje się wyraźny spadek stężenia rozpuszczonego tlenu, co może ograniczać aktywność mezofilnych mikroorganizmów tlenowych oraz stanowić czynnik ograniczający tempo biodegradacji w warunkach tlenowych (Chandra et al., 2013; Zhou & Crawford, 1995). Kolejnym czynnikiem środowiskowym odgrywającym istotną rolę w przebiegu biodegradacji węglowodorów ropopochodnych jest odczyn gleby (pH). Parametr ten oddziałuje na aktywność metaboliczną mikroorganizmów oraz aktywność enzymów uczestniczących w procesach rozkładu zanieczyszczeń. Dostępne dane literaturowe wskazują, że biodegradacja węglowodorów ropopochodnych w glebie efektywnie zachodzi przy pH gleby mieszczącym się w granicach 6–9 (Das & Chandran, 2010), a jej największą skuteczność obserwuje się przy wartościach 7,5-7,8 (Atlas, 1981). Wartości pH wykraczające poza optymalny zakres mogą hamować metabolizm drobnoustrojów, zakłócać równowagę między wytwarzaniem a zużyciem ATP oraz zmniejszać wydajność biomasy, co w konsekwencji prowadzi do obniżenia aktywności mikroorganizmów i spadku efektywności procesów biodegradacji zanieczyszczeń ropopochodnych (Kluk & Steliga, 2017; Steliga, 2009). Nie mniej istotnym czynnikiem środowiskowym jest wilgotność gleby. Obecność wody w matrycy glebowej umożliwia ruch mikroorganizmów odpowiedzialnych za degradację zanieczyszczeń, dyfuzję substratów i związków odżywczych do komórki oraz usuwanie produktów metabolizmu z komórki, a także utrzymanie prawidłowego ciśnienia osmotycznego i pH. Analiza dostępnych danych literaturowych wykazała, że optymalna wilgotność gleby w trakcie biodegradacji tlenowej powinna mieścić się w zakresie 7–24% (Hossein et al., 2021; Kluk & Steliga, 2017; Steliga, 2009). Na efektywność biodegradacji zanieczyszczeń ropopochodnych w glebie wpływa również dostępność składników biogennych.. Zapewnienie optymalnej

zawartości azotu, fosforu, siarki, wapnia, magnezu czy potasu, w glebie podczas procesu jej oczyszczania sprzyja wzrostowi oraz aktywności mikroorganizmów glebowych (Steliga, 2009). W glebach silnie zanieczyszczonych węglowodorami ropopochodnymi (gleby z obszarów górniczych i kopalnianych) obserwuje się zaburzenie optymalnego stosunku C:N:P, dlatego w celu prawidłowego przebiegu procesu biodegradacji zanieczyszczeń wskazane jest wzbogacenie gleby w odpowiednie pierwiastki biogenne. Kluk i Steliga wskazują, że substancje biogenne mogą być dostarczane do gleby w formie rozpuszczalnych w wodzie soli mineralnych (KNO_3, $NaNO_3$, NH_3PO_4, NH_4NO_3, K_2HPO_4, $MgNH_4PO_4$), nawozów oleofilowych (występujących zazwyczaj w postaci granulek pokrytych substancją hydrofobową, umożliwiających lokalne dostarczenie składników na granicy faz woda–węglowodory) oraz nawozów nieorganicznych o powolnym uwalnianiu azotu i fosforu, których tempo uwalniania zależy od powierzchni tabletki i mechanizmu osmotycznego (Kluk & Steliga, 2017; Rosa & Triguis, 2007; Xu et al., 2004). Kolejnym niezwykle ważnym czynnikiem środowiskowym jest obecność tlenu, ponieważ większość mikroorganizmów zdolnych do degradacji węglowodorów to organizmy tlenowe. Tlen warunkuje przebieg procesu biodegradacji, uczestnicząc w reakcjach enzymatycznych katalizowanych przez hydroksylazy alkanowe oraz mono- i dioksygenazy w trakcie rozkładu n-alkanów, a także w procesach utleniania pierścieni aromatycznych w cząsteczkach WWA (Das & Chandran, 2011; Rojo, 2009; Van Hamme et al., 2003). Niedobór tlenu może zatem istotnie ograniczać skuteczność procesu biodegradacji, dlatego w celu zapewnienia wysokiej skuteczności oczyszczania skażonego gruntu zaleca się jego napowietrzanie, które może odbywać się poprzez bronowanie (mechaniczne spulchnianie gruntu, z ang. *landfarming*), natlenianie lub metody chemiczne. Spośród wymienionych metod do napowietrzania powierzchniowych warstw gruntu najczęściej wykorzystuje się bronowanie, natomiast iniekcja powietrza lub napowietrzonej wody poprzez układ przewodów lub drenów umożliwia dotlenienie głębszych warstw gleby (Kluk & Steliga, 2017). Alternatywą dla tych metod są metody chemiczne, w tym chemiczne utlenianie, które polega na dostarczeniu do zanieczyszczonego gruntu alternatywnych źródeł tlenu, czyli substancji wolno uwalniających tlen (np. CaO_2) lub silnych utleniaczy (np. $KMnO_4$). W dostępnej literaturze przedmiotu wskazuje się również możliwość suplementacji gruntu nadtlenkiem wodoru lub ozonem, jednak praktyczne wykorzystanie tych związków jest ograniczone ze względu na możliwość toksycznego oddziaływania na mikrobiotę glebową (Hossein et al., 2021; Souza et al., 2014; Steliga, 2009). Ostatnim z omawianych czynników środowiskowych determinujących skuteczność biodegradacji zanieczyszczeń ropopochodnych jest rodzaj gleby, który wpływa zarówno na aktywność mikroorganizmów, jak i zachowanie zanieczyszczeń w środowisku glebowym. Jak powszechnie wiadomo, gleby różnią się pod względem składu granulometrycznego oraz zawartości materii organicznej. W glebach o większej zawartości materii organicznej oraz frakcji drobnoziarnistej obserwuje się silniejszą sorpcję zanieczyszczeń ropopochodnych, takich jak TPH, BTEX i WWA. Prowadzi to do ograniczenia ich mobilności i biodostępności dla mikroorganizmów, co może znacząco spowalniać tempo biodegradacji zanieczyszczeń (Ghosal et al., 2016; Hossein et al., 2021). W tym kontekście można wysunąć ogólny wniosek, że biodegradacja węglowodorów ropopochodnych w glebach o większej średnicy ziaren i


małej zawartości materii organicznej (np. gleby piaszczyste) będzie zachodziła bardziej efektywnie aniżeli w glebach o mniejszej średnicy ziaren i dużej zawartości materii organicznej (np. gleby gliniaste) (Ortega-Calvo et al., 1997).

- ***Czynniki biologiczne***

Czynniki biologiczne odgrywają często najważniejszą rolę w procesie oczyszczania gruntów zanieczyszczonych węglowodorami ropopochodnymi, ponieważ to aktywność mikroorganizmów decyduje o rzeczywistej skuteczności biodegradacji. Należą do nich przede wszystkim liczebność i skład jakościowy mikrobioty, biodostępność zanieczyszczeń dla mikroorganizmów degradujących zanieczyszczenia, zdolność adaptacyjna organizmów do warunków środowiskowych oraz efektywność enzymatyczna w rozkładzie związków toksycznych (Steliga, 2009).

Pierwszym z wymienionych czynników biologicznych jest liczebność mikroorganizmów. Maleci i in. w swojej publikacji wskazują, że typowej glebie wynosi ona zazwyczaj od 10^5 do 10^7 CFU/g s.m., jednak aby biodegradacja zanieczyszczeń przebiegała pomyślnie ich ilość nie powinna spaść poniżej 10^3 CFU/g s.m. (Maletic et al., 2013). Liczebność mikroorganizmów poniżej 10^3 CFU/g s.m. gleby może wskazywać na przekroczenie poziomu stężeń zanieczyszczeń tolerowanego przez mikroorganizmy glebowe, prowadząc w konsekwencji do zahamowania ich aktywności metabolicznej oraz ograniczenia liczebności (Petrović et al., 2008). Równie istotnym czynnikiem biologicznym jest skład jakościowy mikrobioty, gdyż jak wiadomo różne gatunki bakterii, grzybów i drożdży obecnych w glebie wykazują odmienne zdolności do rozkładu konkretnych grup zanieczyszczeń, co warunkuje efektywność procesu oczyszczania. W typowych glebach jedynie 0,01%-1% ogólnej liczby bakterii glebowych wykazuje zdolność do rozkładu węglowodorów (Gałązka, 2009), natomiast w przypadku grzybów wartości te mieszczą się w przedziale 0,1-10% (Atlas & Bartha, 1998). Liczba ta może się jednak zmieniać w zależności od poziomu skażenia. Liczne dane literaturowe wskazują, że w glebach silnie zanieczyszczonych węglowodorami ropopochodnymi obserwuje się wyraźny wzrost udziału mikroorganizmów degradujących te związki, sięgający nawet od kilku do kilkudziesięciu procent całkowitej mikrobioty. (Atlas, 1981; Leahy & Colwell, 1990). Chandra i in. wskazali bardziej szczegółowe dane, zaznaczając że w glebach zanieczyszczonych substancjami ropopochodnymi 0,13%-50% populacji bakterii glebowych może wykazywać zdolność do wykorzystania węglowodorów ropopochodnych jako jedynego źródła węgla, natomiast w przypadku grzybów wartości te mogą wynosić nawet 6%-82% (Chandra et al., 2013). Kolejnym kluczowym czynnikiem biologicznym determinującym przebieg biodegradacji jest biodostępność zanieczyszczeń, która zależy zarówno od właściwości matrycy glebowej, jak i procesów sorpcji i desorpcji zachodzących w glebie. Koshlaf i Ball wskazują, że węglowodory silnie zaadsorbowane w porach gleby wykazują zazwyczaj mniejszą biodostępność dla mikroorganizmów niż węglowodory zaadsorbowane na jej powierzchni. Wynika to przede wszystkim z ograniczonego dostępu mikroorganizmów do wnętrza porów glebowych, gdzie zanieczyszczenia są fizycznie odizolowane od strefy ich aktywności metabolicznej. (Koshlaf & Ball, 2017). Ponadto jak wskazano wcześniej, gleby

zawierające duże ilości frakcji ilastych oraz materii organicznej silniej sorbują zanieczyszczenia, aniżeli gleby piaszczyste i ubogie w próchnicę, co sprzyja ich trwałemu unieruchomieniu wskutek silnego wiązania w strukturze tych gleb, a tym samym ogranicza ich dostępność biologiczną. Na rysunku 3 przedstawiono mechanizm inicjowania biodegradacji zanieczyszczeń ropopochodnych przez mikroorganizmy w warunkach ograniczonej dostępności substratu.



Rysunek 3 Interakcje zachodzące w układzie gleba-węglowodory ropopochodne-mikroorganizmy (opracowanie własne na podstawie Koshlaf & Ball, 2017).

Omawiając czynniki biologiczne mające wpływ na efektywność biodegradacji zanieczyszczeń ropopochodnych w glebach, nie sposób nie wspomnieć o zdolności mikroorganizmów do adaptacji w zmiennych warunkach środowiskowych. Czynnik ten jest szczególnie istotny podczas zabiegu bioaugmentacji, ponieważ skuteczność wprowadzanych do środowiska szczepów drobnoustrojów w dużym stopniu zależy od ich zdolności do przystosowania się do warunków panujących w skażonej glebie, takich jak wysokie stężenie zanieczyszczeń, ograniczona dostępność składników pokarmowych czy konkurencja ze strony autochtonicznej mikrobioty glebowej. Dlatego wielu badaczy wskazuje, że najbardziej korzystne jest wykorzystanie biopreparatów opracowanych na bazie mikroorganizmów autochtonicznych, które naturalnie są przystosowane do lokalnych warunków środowiskowych i wykazują większą przeżywalność oraz aktywność metaboliczną w środowisku skażonym, co może przekładać się na wyższą efektywność biodegradacji związków ropopochodnych (Brzeszcz, 2017; Steliga et al., 2012;). Ostatnim z omówionych czynników biologicznych jest efektywność enzymatyczna, która bezpośrednio determinuje tempo rozkładu zanieczyszczeń. Aktywność enzymów odpowiedzialnych za degradację węglowodorów wpływa na szybkość przekształcania związków toksycznych w prostsze mniej toksyczne produkty

pośrednie lub produkty o właściwościach obojętnych, a jej poziom zależy zarówno od właściwości samych mikroorganizmów, jak i od warunków środowiskowych (Koshlaf & Ball, 2017).

1.4.2. Metody wspomagania procesu biodegradacji

W warunkach naturalnych proces rozkładu węglowodorów przebiega stosunkowo wolno, co wynika z fizykochemicznych właściwości związków ropopochodnych, ograniczonej aktywności enzymatycznej mikroorganizmów oraz spontanicznego przebiegu reakcji degradacyjnych. Samooczyszczanie (bioattuneacja) to proces usuwania zanieczyszczeń oparty na naturalnej biodegradacji mikroorganizmów autochtonicznych, zachodzący bez dodatkowej ingerencji człowieka (Yu et al., 2005). Proces ten jest ściśle powiązany z naturalnym obiegiem pierwiastków w glebie i stanowi podstawę do projektowania skutecznych strategii wspomagania procesu biodegradacji. Jednakże tempo biodegradacji w warunkach samooczyszczania jest zazwyczaj niskie i może trwać wiele miesięcy lub nawet lat, zarówno w przypadku węglowodorów alifatycznych, jak i aromatycznych. W przypadku gleb historycznie skażonych, takich jak doły urobkowe, samooczyszczanie jest zazwyczaj niewystarczające ze względu na wysokie stężenia zanieczyszczeń oraz liczne związki silnie zaadsorbowane na matrycy mineralnej i organicznej, dlatego w praktyce do oczyszczania terenów zanieczyszczonych węglowodorami ropopochodnymi wykorzystuję się biodegradację wspomaganą (Steliga, 2009). Bioremediacja wspomagana polega na zwiększeniu efektywności naturalnego procesu samooczyszczania poprzez zastosowanie odpowiednich działań inżynieryjnych i biotechnologicznych. Wśród najczęściej stosowanych metod wyróżnia się biostymulację oraz bioaugmentację:

- **Biostymulacja** – polega na wprowadzeniu do środowiska gruntowo-wodnego czynników wspierających rozwój i aktywność mikroorganizmów autochtonicznych. Obejmuje ona dostarczenie tlenu, wody, substancji biogennych (np. azotu, fosforu) a także w niektórych przypadkach również surfaktantów zwiększających biodostępność zanieczyszczeń. Zapewnienie optymalnych warunków dla rodzimej mikrobioty w ramach biostymulacji prowadzi do zwiększenia efektywności naturalnych procesów samooczyszczania środowiska glebowego (Greco Lucchina et al., 2026; Mekonnen et al., 2024).
- **Bioaugmentacja** – polega na inokulacji zanieczyszczonego środowiska odpowiednio wyselekcjonowanymi mikroorganizmami charakteryzującymi się wysoką skutecznością biodegradacji różnych grup węglowodorów, takich jak n-alkany, BTEX czy WWA. W praktyce w zabiegu inokulacji można stosować zarówno szczepy autochtoniczne (pozyskane ze skażonej gleby), jak i allochtoniczne (szczepy obce), a niektórych przypadkach szczepy modyfikowane genetycznie zapewniające większą efektywność degradacji zanieczyszczeń. Warto zauważyć, że szczepy pochodzące z miejsca skażenia wykazują zwykle lepszą adaptację i mniejsze ryzyko antagonistycznych interakcji z mikrobiotą glebową aniżeli szczepy obce, co może być szczególnie istotne na etapie opracowania

biopreparatów wykorzystywanych w zabiegu bioaugmentacji (Greco Lucchina et al., 2026; Mekonnen et al., 2024).

Obecnie znanych jest wiele gatunków mikroorganizmów zdolnych do degradacji różnych frakcji węglowodorów ropopochodnych, które mogą być z powodzeniem wykorzystywane do opracowywania biopreparatów stosowanych w bioaugmentacji. Obejmują one bakterie, grzyby i drożdże różniące się mechanizmami metabolicznymi oraz preferencjami względem określonych grup związków, takich jak n-alkany, BTEX czy wielopierścieniowe węglowodory aromatyczne. Pojedynczy gatunek mikroorganizmu zwykle metabolizuje ograniczony zakres substratów, dlatego w bioremediacji gruntów zanieczyszczonych różnego typu substancjami chemicznymi wskazane jest stosowanie biopreparatów opartych na kilku lub kilkunastu szczepach. Takie konsorcja mikroorganizmów pozwalają na komplementarne wykorzystanie różnych szlaków metabolicznych wynikających z różnorodności metabolicznej drobnoustrojów, co zapewnia szersze spektrum zanieczyszczeń ulegających degradacji oraz zwiększa stabilność procesu w zmiennych warunkach środowiskowych (Steliga et al., 2018). Dodatkowo współdziałanie kilku szczepów mikroorganizmów może prowadzić do sekwencyjnego rozkładu złożonych związków, w którym produkty pośrednie jednych organizmów stanowią substrat dla innych.

Wskazuje się, że do inokulacji rzeczywistych gleb zanieczyszczonych węglowodorami ropopochodnymi najbardziej efektywne są biopreparaty opracowane z wykorzystaniem mikroorganizmów pochodzących ze środowiska naturalnego, najlepiej z miejsca skażenia. Takie szczepy są już przystosowane do obecności zanieczyszczeń oraz specyficznych warunków fizykochemicznych, co przekłada się na wyższą skuteczność degradacji i większą przeżywalność po inokulacji. Ponadto wykorzystanie w biopreparatach szczepów autochtonicznych zmniejsza ryzyko występowania interakcji antagonistycznych z mikrobiotą glebową oraz wypierania rodzimych populacji (Brzeszcz et al., 2013; Polyaka et al., 2018; Sarkar et al., 2016). Warto nadmienić, że w trakcie opracowywania biopreparatu oraz przeprowadzania zabiegu inokulacji należy również zwrócić uwagę na bezpieczeństwo środowiskowe stosowanych mikroorganizmów. Wprowadzane szczepy nie powinny bowiem wykazywać właściwości patogennych względem roślin, zwierząt ani ludzi, nie powinny produkować toksycznych metabolitów pośrednich, a także nie powinny być nośnikami genów oporności na antybiotyki, które mogłyby być przenoszone w środowisku (Kluk & Steliga, 2017). Istotne jest również monitorowanie aktywności oraz przeżywalności mikroorganizmów wprowadzanych do oczyszczanej gleby po aplikacji, aby móc ocenić rzeczywistą skuteczność zastosowanego biopreparatu (Gałązka, 2009; Mrozik et al., 2005; Vogel 1996). W tabeli 1 zestawiono wybrane bakterie i grzyby zdolne do degradacji różnych grup węglowodorów ropopochodnych, które mogą być wykorzystane w biopreparatach stosowanych w zabiegu bioagmentacji gleb zanieczyszczonych substancjami ropopochodnymi.

Tabela 1 Zestawienie bakterii i grzybów rozkładających wybrane grupy węglowodorów ropopochodnych

Rodzaj skażenia	**Bakterie i grzyby**	**Literatura**
Alkany C_2–C_8	*Desulfosarcina* sp., *Mycobacterium* sp., *Pseudomonas butanovora*, *Pseudonocardia* sp. *Rhodococcus* sp., *Rhodococcus ruber*	Jaekel et al., 2013; Kurth et al., 2008; Kotani et al., 2006; Ivshina et al., 2025
TPH (alkany C_{10}–C_{36}), oleje, asfalteny	*Achromobacter* sp., *Acinetobacter baumannii*, *Alcanivorax* sp., *Alcanivorax borkumensis*, *Bacillus* spp., *Bacillus licheniformis, Bacillus subtilis, Dietzia* sp., *Gordonia paraffinivorans, Gordonia sihwensis*, *Mycolicibacterium* sp., *Mycobacterium frederiksbergense* IN53 (*Actinobacteria*, K-strategia) i *Acinetobacter* sp. IN47 (*Gammaproteobacteria*, R-strategia), *Pseudomonas* spp., *Pseudomonas aeruginosa*, *Pseudomonas stutzeri, Pseudomonas veronii, Rhodococcus* sp., *Rhodococcus erythropolis, Rhodococcus ruber, Stenotrophomonas maltophilia*.	Li, 2021; Liu et al., 2026; Baltaci et al., 2026; Kadri et al. 2018; Das et al., 2024; Fan et al., 2020; Parthipan et al., 2017; Hu et al., 2020; Silva et al., 2019; Brzeszcz et al., 2023;Brzeszcz et al., 2016; Wang et al., 2025; Mullaeva et al., 2022; Xiang et al., 2022; Zhang et al., 2025; Naloka et al., 2022; Stancu, 2022
Węglowodory monoaromatyczne (BTEX)	*Achromobacter insolitus, Achromobacter xylosoxidans, Aeribacillus pallidus, Archaeoglobus fulgidus*, *Bacillus cereus*, *Bacillus cibi, Bacillus megaterium, Mycobacterium cosmeticum*, *Pseudomonas aeruginosa, Pseudomonas putida, Pseudomonas fluoresce* , *Rhodococcus rhodochrous*, *Sphingobacterium* sp.,	Ma et al., 2015; Mnif et al., 2014; Wilkes et al., 2016; Janbandhu and Fulekar, 2011; Zhang et al., 2013; Mukherjee et al., 2017; Chicca et al., 2016; Huizenga and Semprini, 2023
Wielopierścieniowe węglowodory aromatyczne (WWA)	*Achromobacter xylosoxidans, Aeribacillus pallidus*, *Aspergillus flavus*, *Aspergillus niger*, *Bacillus Licheniformis, Bacillus Mojavensis*, *Bacillus subtilis*, *Brevundimonas* sp., *Burkholderia* sp., *Fusarium solani*, *Gordonia* sp., *Mycobacterium* sp., *Mycobacterium frederiksbergense*, *Nocardioides* sp., *Pleurotus* spp., *Pseudomonas* spp., *Pseudomonas aeruginosa*, *Pseudomonas putida, Pseudomonas stutzeri*, *Rhodococcus* sp., *Rhodococcus erythropolis, Sphingomonas sp., Sphingomonas yanoikuyae*, *Stenotrophomonas* sp., *Stenotrophomonas maltophilia*, *Streptomyces* sp.	Ma et al., 2015; Mnif et al., 2014; Al-Dossary et al., 2020; Manjunatha et al., 2025; Eskandari et al., 2017; Firmin and Laurette, 2026; Smułek et al., 2020; Revathy et al., 2015; Hesham et al., 2017, Qi et al., 2017; Dudhagara & Dave, 2018; Willumsen et al., 2001; Ma et al., 2023; Innes, 2023; Sivasamy et al., 2025, Gao et al., 2023; Tang et al., 2011; Singh and Tiwary, 2017; Ma et al., 2023; Song et al., 2021; Lopez et al. 2022; Tiwari et al., 2016; Xiao et al., 2021; Baoune et al., 2019

Obecnie, w celu poprawy efektywności remediacji terenów zanieczyszczonych węglowodorami ropopochodnymi, coraz częściej stosuje się tzw. strategie łączone, w których bioaugmentacja wykorzystywana jest w połączeniu z dodatkowymi metodami wspomagającymi proces biodegradacji (Rys. 4). Takie podejście pozwala na jednoczesne zwiększenie aktywności mikroorganizmów oraz poprawę dostępności zanieczyszczeń i

składników pokarmowych, co przekłada się na przyspieszenie procesów oczyszczania zanieczyszczonych gruntów. Bioaugmentacja w tego typu strategiach nie ogranicza się wyłącznie do inokulacji pojedynczymi szczepami lub biopreparatami na bazie wyselekcjonowanych szczepów mikroorganizmów, lecz obejmuje również modyfikację biopreparatów oraz warunków prowadzenia procesu oczyszczania. Najważniejsze z wymienionych podejść oraz mechanizmy ich działania zostaną pokrótce omówione w dalszej części pracy oraz w artykułach I-VII



Rysunek 4 Schemat oczyszczania środowiska wodno-gruntowego z wykorzystaniem strategii łączonych (opracowanie własne na podstawie Eras-Muñoz et al. 2022 i poddane obróbce przez OpenAI)

W kontekście wspomagania biodegradacji węglowodorów ropopochodnych w glebach, ciekawą strategią jest modyfikacja procesu bioaugmentacji surfaktantami. Związki powierzchniowo czynne dzięki obecności w swoich cząseczkach zarówno części hydrofilowej jak i hydrofobowej mają zdolność do gromadzenia się się na granicy faz o różnej polarności (olej–woda), co prowadzi do zmniejszenia napięcia powierzchniowego oraz międzyfazowego w układzie, w efekcie czego zwiększa się dostępność związków


hydrofobowych dla mikroorganizmów, co ułatwia ich rozkład i wpływa na poprawę efektywności oczyszczania skażonych gleb (Badmus et al., 2021; Zajic et al., 1983). W niniejszej rozprawie doktorskiej szczególną uwagę zwrócono na możliwość wzbogacenia gleby w biosurfaktanty, które stanowią grupę związków powierzchniowo czynnych wytwarzanych przez mikroorganizmy, głównie bakterie. Powstają one jako metabolity wtórne i mogą być wydzielane do środowiska zewnątrzkomórkowo lub być związane z powierzchnią komórek bakteryjnych. Podobnie jak inne środki powierzchniowo czynne, biosurfaktanty mają budowę amfifilową, gdzie część hydrofilową tworzą mono- lub polisacharydy, peptydy albo aminokwasy, natomiast część lipofilową stanowią jeden lub kilka długołańcuchowych kwasów tłuszczowych, hydroksykwasów tłuszczowych lub α-alkilo-β-hydroksykwasów zawierających od 30 do 40 atomów węgla (obie części połączone są najczęściej grupą acylową lub wiązaniem glikozydowym, co umożliwia lokalizację biosurfaktantów na granicy faz i redukcję napięcia powierzchniowego oraz międzyfazowego) (Eras-Muñoz et al., 2022; Sharma et al., 2021). Różnice w budowie chemicznej części hydrofilowej i lipofilowej stanowią podstawę klasyfikacji biosurfaktantów na: glikolipidy, lipoproteiny i lipopeptydy, fosfolipidy i kwasy tłuszczowe, biopolimery oraz struktury biologiczne (Sharma et al., 2021).

W praktyce bioremediacyjnej do modyfikacji procesu bioaugmentacji gleb zanieczyszczonych węglowodorami ropopochodnymi można wykorzystywać zarówno gotowe biosurfaktanty dodawane bezpośrednio do biopreparatu lub gleby, jak również mikroorganizmy zdolne do ich wytwarzania, co pozwala na stopniowe uwalnianie tych związków i wspiera późniejszą aktywność inokulowanych szczepów. Wykorzystanie w procesie bioremediacji biosurfaktantów pozwala na zwiększenie częstotliwości kontaktu drobnoustrojów z ropą, co wynika bezpośrednio z ich zdolności do lokalizowania się na powierzchniach międzyfazowych oraz zmiany właściwości tych powierzchni. Poza tym Bjerk i in. w swoim artykule przeglądowym podkreślił, że właściwości biosurfaktantów ułatwiają kontakt drobnoustrojów z zanieczyszczeniami silnie zadsorbowanymi na cząstkach mineralnych i organicznych, nie tylko poprzez oddziaływanie na zanieczyszczenie, ale również poprzez oddziaływanie na powierzchnię komórek drobnoustrojów glebowych. co prowadzi do zwiększenia ich hydrofobowości i sprzyja łączeniu się z zanieczyszczeniami o charakterze hydrofobowym (Bjerk et al., 2021). W konsekwencji wprowadzenie biosurfaktantów do gleby może istotnie wpłynąć na poprawę efektywności biodegradacji węglowodorów ropopochodnych, stymulację rozwoju populacji mikroorganizmów autochtonicznych oraz ograniczenie toksycznego działania niektórych frakcji zanieczyszczeń na organizmy żywe (Bjerk et al., 2021; Dini et al., 2024).

Mechanizm działania biosurfaktantów w bioremediacji gruntów zanieczyszczonych węglowodorami ropopochodnymi obejmuje trzy podstawowe procesy fizykochemiczne: mobilizację, solubilizację oraz emulgację (Pacwa-Płociniczak et al., 2011). Mobilizacja to proces, który zachodzi przy stężeniach poniżej krytycznego stężenia miceli (CMC) i polega na zmniejszeniu napięcia powierzchniowego oraz międzyfazowego w układach powietrze–woda i gleba–woda, co osłabia siły kapilarne utrzymujące frakcje olejowe w strukturze gleby. Po przekroczeniu krytycznego stężenia miceli zachodzi solubilizacja. W procesie solubilizacji cząsteczki biosurfaktantów tworzą

micele, które zwiększają pozorną rozpuszczalność związków hydrofobowych w środowisku wodnym. Jak wskazują dostępne źródła literaturowe, w wyniku solublizaji hydrofobowe fragmenty cząsteczek biosurfaktantów są skierowane do wnętrza miceli, natomiast ich części hydrofilowe pozostają zwrócone w stronę fazy wodnej. Taka organizacja struktury micelarnej umożliwia włączanie hydrofobowych związków ropopochodnych do wnętrza miceli, co wpływa na zwiększenie ich biodostępności dla mikroorganizmów glebowych (Bhanu Revathi et al., 2022; Pacwa-Płociniczak et al., 2011; Patowary et al., 2017) (Rys. 5).

Pacwa-Płociniczak i in. podkreślili również znaczenie emulgacji w procesie biodegradacji zanieczyszczeń ropopochodnych, która polega na tworzeniu emulsji składającej się z bardzo małych kropli oleju rozproszonych w fazie wodnej. W swoim artykule wskazali, że w wyniku emulgacji zanieczyszczeń olejowych dochodzi do zwiększenia powierzchni kontaktu pomiędzy zanieczyszczeniem, a środowiskiem wodnym, co sprzyja szybszemu rozpuszczaniu węglowodorów i intensyfikuje ich rozkład biologiczny (Pacwa-Płociniczak et al., 2011). Znaczenie emulgacji w procesie biodegradacji węglowodorów ropopochodnych potwiedzili Brakstad i in., którzy w swoich badaniach wykazali, że przy tej samej ilości oleju stopień biodegradacji wzrasta wraz ze zmniejszaniem średnicy kropli zemulgowanego oleju (Brakstad et al., 2015). Z kolei Karlapudi i in. wskazali, że do emulgacji węglowodorów ropopochodnych najlepiej sprawdzą się biosurfaktanty o dużej masie cząsteczkowej, takie jak lichenyzyny, ramnolipidy oraz surfaktyna (Karlapudi et al., 2018).




Rysunek 5 Schemat oddziaływania biosurfaktantów na zanieczyszczenia ropopochodne w trakcie a) mobilizacji, b) solubilizacji (opracowanie własne na podstawie Selva Filho et al., 2023 oraz Dhanya 2021 i poddane obróbce przez OpenAI)

Kolejną strategią wspomagania procesu biodegradacji węglowodorów ropopochodnych w glebach jest fitoremediacja. Fitoremediacja to technologia oczyszczania gruntów, która polega na zastosowaniu roślin wraz z mikroorganizmami zasiedlającymi strefę korzeniową, do pobierania, gromadzenia lub biodegradacji substancji toksycznych (Barathi & Vasudevan, 2001; Tonelli et al., 2022). Choć od lat wykorzystywana jest głównie do oczyszczania gleb zanieczyszczonych metalami

ciężkimi, w połączeniu z bioaugmentacją może stanowić skuteczne narzędzie również w przypadku gleb skażonych substancjami hydrofobowymi, takimi jak węglowodory ropopochodne (Sarfaraz et al., 2025).

Wykorzystanie roślin w procesie bioremediacji gruntów zanieczyszczonych węglowodorami ropopochodnymi opiera się przede wszystkim na modyfikacji właściwości fizykochemicznych gleby w ryzosferze oraz stymulacji aktywności mikroorganizmów odpowiedzialnych za biodegradację zanieczyszczeń. Liczne badania wykazały, że obecność roślin prowadzi do poprawy struktury gleby, zwiększenia jej porowatości i napowietrzenia, a także wzrostu zawartości materii organicznej, co sprzyja rozwojowi mikrobioty glebowej oraz intensyfikacji procesów biodegradacyjnych (Lü et al., 2024; Xiao et al., 2024) . Jednym z najważniejszych czynników odpowiedzialnych za stymulujący wpływ roślin na procesy biodegradacji jest wydzielanie eksudatów korzeniowych. Yang i in. wykazali, że stanowią one cenne źródło składników odżywczych oraz metabolitów pośrednich dla mikroorganizmów zasiedlających ryzosferę. Związki te zwiększają aktywność enzymatyczną drobnoustrojów oraz ich potencjał metaboliczny, co przekłada się na wyższą efektywność degradacji węglowodorów ropopochodnych (Gałązka & Gałązka, 2015; Yang et al., 2023). Ponadto wydzieliny korzeniowe wpływają na skład i liczebność społeczności mikroorganizmów, selektywnie promując rozwój szczepów zdolnych do rozkładu związków toksycznych, dzięki czemu proces biodegradacji zanieczyszczeń w strefie korzeniowej zachodzi bardziej intensywnie (Lü et al., 2024; Rohrbacher i St-Arnaud, 2016). Dodatkowo jak podkreślają Rohrbacher i St-Arnaud eksudaty korzeniowe mogą zwiększać biodostępność węglowodorów ropopochodnych poprzez modyfikację procesów sorpcji i desorpcji zachodzących w glebie oraz zmianę właściwości chemicznych środowiska ryzosferowego. W rezultacie ułatwiony zostaje kontakt zanieczyszczeń z mikroorganizmami degradującymi, zarówno autochtonicznymi, jak i wprowadzanymi do środowiska w ramach bioaugmentacji, co może prowadzić do zwiększenia efektywności procesów biodegradacyjnych (Rohrbacher & St-Arnaud, 2016). Mechanizmy wspomagające remediację nie ograniczają się jednak wyłącznie do stymulacji aktywności mikroorganizmów. Badania Huang i in. wykazały, że niektóre gatunki roślin posiadają zdolność do pobierania, akumulacji lub adsorpcji niewielkich ilości zanieczyszczeń ropopochodnych, przyczyniając się tym samym do obniżenia ich stężenia w glebie (Huang et al., 2018). Proces ten może pośrednio wspomagać dalszą biodegradację poprzez zmniejszenie presji toksycznej oddziałującej na mikroorganizmy oraz poprawę warunków środowiskowych panujących w strefie korzeniowej. Pobieranie zanieczyszczeń organicznych przez rośliny zachodzi przede wszystkim wraz z wodą na drodze dyfuzji prostej. Należy jednak podkreślić, że ze względu na hydrofobowy charakter większości składników ropy naftowej, ich transport do tkanek roślinnych jest znacznie ograniczony (Wojtowicz et al., 2023a; 2026a, 2026b). Wynika to z wysokiej lipofilności oraz dużej masy cząsteczkowej tych związków, które utrudniają ich przemieszczanie się w roztworze glebowym i przenikanie przez bariery biologiczne. Jak wykazano w licznych badaniach, stopień pobierania lipofilowych związków organicznych przez rośliny jest dodatnio skorelowany z zawartością lipidów w epidermie korzeni, co wskazuje na istotną rolę właściwości fizjologicznych roślin w procesie akumulacji zanieczyszczeń (Kamath et al., 2004; Khan et al., 2013; Pawlik & Piotrowska-Seget, 2015; Wang et al., 2016; Wayens et al., 2009a, 2009b, 2009c, 2009d). Złożoność

mechanizmów zachodzących z udziałem roślin znalazła odzwierciedlenie w wyodrębnieniu kilku podstawowych rodzajów fitoremediacji (rys. 6). Do najważniejszych z nich zalicza się fitoekstrakcję, fitodegradację, ryzofiltrację, fitostabilizację, fitoewaporację, fitostymulację oraz epuwalizację (Ijaz et al., 2016). Każdy z wymienionych procesów opiera się na odmiennych mechanizmach usuwania, transformacji lub unieruchamiania zanieczyszczeń, jednak ich wspólnym celem jest zwiększenie efektywności oczyszczania środowiska glebowego.



Rysunek 6 Schemat przedstawiający poszczególne rodzaje fitoremediacji (Obraz wygenerowany przez OpenAI i poddany obróbce przez autora) .

Projektując proces bioremediacji z wykorzystaniem techniki fitoremediacji lub fitoremediacji wspomaganej bioaugmentacją, należy uwzględnić, cechy jakimi powinny się charakteryzować rośliny wykorzystywane w badaniach. Wskazuje się, że rośliny fitoremediujące powinny wykazywać dużą tolerancję na wysokie stężenia ksenobiotyków, zdolność do akumulacji lub biodegradacji różnych rodzajów zanieczyszczeń, szybki wzrost, wysoką produkcją biomasy oraz odporność na choroby i trudne warunki środowiskowe (Cooney et al., 1985; Barathi et al., 2001). W zależności od rodzaju zanieczyszczeń obecnych w glebie w procesach fitoremediacji wykorzystuje się różne grupy roślin, spośród których największe znaczenie mają hiperakumulatory oraz naftofity.

- **Hiperakumulatory** to rośliny wykorzystywane do oczyszczania gleb zanieczyszczonych metalami ciężkimi, najczęściej na terenach poprzemysłowych. Charakteryzują się zdolnością do pobierania, transportu

i gromadzenia w swoich tkankach bardzo dużych ilości metali bez wywoływania efektu toksycznego. Zgodnie z przyjętą definicją hiperakumulacji za hiperakumulatory można uznać rośliny zdolne do gromadzenia w swoich tkankachzanieczyszczeń w ilości: około 100 mg/kg s.m. dla selenu, kadmu i talu, 300 mg/kg s.m. dla miedzi, kobaltu i chromu, 1000 mg/kg s.m. dla niklu, arsenu i ołowiu, 3000 mg/kg s.m. dla cynku oraz 10000 mg/kg s.m. dla manganu (Baker i Brooks, 1989; Reeves, 2003; Reeves et al., 2018).

- **Naftofity** (rośliny wspomagające biodegradację węglowodorów) to rośliny wykorzystywane do oczyszczania gleb zanieczyszczonych związkami ropopochodnymi, głównie na terenach skażonych wyciekami ropy naftowej i działalnością przemysłu naftowego. Charakteryzują się wysoką tolerancją na obecność substancji ropopochodnych w glebie oraz zdolnością do wspierania ich rozkładu poprzez stymulowanie aktywności mikroorganizmów w strefie korzeniowej (Kluk & Steliga, 2017).

W oczyszczaniu gruntów z terenów dołów urobkowych stosuje się odpowiednio dobrane gatunki traw (Poaceae), roślin motylkowych (Fabaceae), roślin zielnych oraz drzew, które mogą wspomagać biodegradację węglowodorów ropopochodnych. Do najczęściej opisywanych w literaturze gatunków traw zalicza się turzycę (*Carex hirta*), wiechlinę łąkową (*Poa pratensis*), życicę trwałą (*Lolium perenne*) (Pawluśkiewicz et al. 2022; Wojtowicz et al., 2026b), kłosówkę wełnistą (*Holcus lanatus*) (Rabêlo et al. 2021), kostrzewę czerwoną (*Festuca rubra*), kostrzewę trzcinową (*Festuca arundinacea*) (Borowik et al. 2019; Steliga & Kluk 2020), kupkówkę pospolitą (*Dactylis glomerata*) (Gałązka & Gałązka 2015), sorgo dwubarwne (*Sorghum bicolor*) (Bangs et al. 2003) oraz kukurydzę zwyczajną (*Zea mays*) (Baoune et al. 2019; Gałązka et al. 2010; Liao et al. 2015; Wojtowicz et al. 2023a). W grupie roślin motylkowych szczególne znaczenie pod kątem wykorzystania w remediacji gruntów zanieczyszczonych substancjami ropopochodnymi przypisuje się koniczynie białej (*Trifolium repens*) (Xi et al. 2018), koniczynie łąkowej czerwonej (*Trifolium pratense*) (Furtak et al. 2020), lucernie siewnej (*Medicago sativa*) (Li et al. 2020), komonicy zwyczajnej (*Lotus corniculatus*) (Pawlik et al. 2017; Wojtowicz & Steliga 2023b; Yousaf et al. 2010) oraz nostrzykowi żółtemu (*Melilotus officinalis*) (Steliga & Kluk 2021), natomiast w grupie roślin zielnych babce zwyczajnej (*Plantago major*), pokrzywie zwyczajnej (*Urtica dioica*) (Małachowska-Jatusz et al. 2012), jeżówce purpurowej (*Echinacea purpurea*) (Heidari et al. 2018; Liu et al. 2012; Wojtowicz et al. 2023b) oraz dziwaczkowi peruwiańskiemu (*Mirabilis jalapa*) (Peng et al. 2009). Coraz częściej w procesie fitoremediacji i fitoremediacji wspomaganej bioaugmentacją, gruntów zanieczyszczonych węglowodorami ropopochodnymi uwzględnia się również gatunki drzewiaste o szybkim wzroście i rozbudowanym systemie korzeniowym, takie jak topole (*Populus L.*) i wierzby (*Salix L.*), które zapewniają wysoką produkcję biomasy oraz dobrą stabilizację gruntu, co przekłada się na wysoką efektywność oczyszczania (Doty 2006; Moore et al. 2006; Hall et al. 2011).

Kolejną, stosunkowo nową strategią wspomagania procesów biodegradacyjnych jest wzbogacanie gleby nanomateriałami, które ze względu na swoje właściwości fizykochemiczne budzą rosnące zainteresowanie naukowców zajmujących się

zagadnieniami z zakresu remediacji gruntów (Mohanapragash et al., 2025; Souza et al., 2020). W literaturze naukowej szczególną uwagę pod względem możliwości wykorzystania w procesie bioremediacji gruntów zanieczyszczonych zarówno związkami nieorganicznymi, jak i organicznymi poświęca się nanocząstkom metali oraz tlenków metali takim jak: tlenek cynku (ZnO) (Pérez-Hernández et al.,2023; Wojtowicz et al., 2026b), magnetyt (Fe_3O_4) (Baragaño et al., 2020), tlenek glinu (Al_2O_3), tlenek krzemu (SiO_2) tlenek tytanu (TiO_2) (Naderi Peikam & Jalali, 2019), a także nowoczesnym materiałom węglowym (Chemng et al., 2019), w tym grafenowi (GO) (Özlü & Elif Fırat, 2025), tlenkowi grafenu (Zhou et al., 2024; Wojtowicz et al., 2026b), nanorurkom węglowym (CNTs) (Moses et al., 2023), oraz różnego rodzaju nanopolimerom (Nwadinigwe & Ugwu, 2018). Wskazuje się, że dzięki dużej powierzchni właściwej i wysokiej reaktywności nanomateriały mogą wchodzić w interakcje z zanieczyszczeniami na poziomie molekularnym, co czyni je potencjalnie skutecznymi narzędziami w usuwaniu szkodliwych substancji ze środowiska (Jiang et al., 2025; Rai et al., 2020; Wang et al., 2026). Istnieją również doniesienia literaturowe wskazujące, że niektóre nanomateriały, np. nanocząstki żelaza zerowartościowego (nZVI) mogą wpływać na mobilność zanieczyszczeń organicznych w środowisku glebowym poprzez modyfikację procesów sorpcji oraz reakcji utleniania i redukcji, co w konsekwencji może pośrednio oddziaływać na ich dostępność w procesach biodegradacji (Liu et al., 2024). Z kolei nanocząstki tlenków metali, takie jak TiO_2 i ZnO, mogą pełnić rolę fotokatalizatorów, prowadząc do generowania reaktywnych form tlenu (ROS) pod wpływem promieniowania, co może inicjować utlenianie związków organicznych, w tym węglowodorów, oraz ich transformację do produktów o prostszej budowie chemicznej (Kessler et al., 2022). W literaturze podkreśla się, że procesy te wpisują się w szerszy potencjał nanomateriałów do inicjowania i przyspieszania przemian zanieczyszczeń organicznych w środowisku glebowym, co może zwiększać efektywność ich dalszej biodegradacji (Chen et al., 2025; Gomes, 2025). Jak wspomniano wcześniej w kontekście biodegradacji węglowodorów ropopochodnych istotną rolę przypisuje się również nanomateriałom węglowym, takim jak nanorurki węglowe (CNT) oraz tlenek grafenu (GO), które mogą pełnić funkcję zarówno adsorbentów zanieczyszczeń organicznych, jak i nośników enzymów lub mikroorganizmów, zwiększając ich stabilność oraz potencjał degradacyjny mikrobioty glebowej (Ramakrishna et al., 2018; Wang et al., 2019). Dodatkowo Azeez i in. wykazali, że niektóre nanomateriały mogą modyfikować formy chemiczne metali ciężkich, np. poprzez redukcję Cr(VI) do mniej toksycznego Cr(III), co prowadzi do zmniejszenia mobilności i biodostępności chromu (Azeez et al., 2021).

Ciekawym zagadnieniem związanym z wykorzystaniem nanomateriałów w kontekście wspomagania bioremediacji gruntów zanieczyszczonych węglowodorami ropopochodnymi jest integracja nanotechnologii z fitoremediacją, określana w literaturze jako nanofitoremediacja. Dostępne dane literaturowe wskazują, że nanomateriały mogą znacznie wspomóc efektywność procesów fitoremediacyjnych, poprzez przyspieszenie degradacji zanieczyszczeń do form łatwiej przyswajalnych przez rośliny, poprawę pobierania i akumulacji zanieczyszczeń przez rośliny fitoremediujące, łagodzenie stresu środowiskowego oraz dostarczanie substancji wspierających wzrost roślin (Chen et al., 2025, Gomez, 2025).

Jak wspomniano wcześniej, dostępne dane literaturowe wskazują, że wprowadzenie nanomateriałów do środowiska glebowego może istotnie wpłynąć na poprawę biodostępności zanieczyszczeń, a także ich przekształcanie do form łatwiej przyswajalnych przez rośliny. W przeglądzie literaturowym przeprowadzonym przez Al-Khayri i in. zwrócono jednak uwagę, że korzyści wynikające z zastosowania nanomateriałów w procesach fitoremediacyjnych nie ograniczają się wyłącznie do zwiększenia dostępności zanieczyszczeń dla roślin. Autorzy podkreślają istotną rolę nanomateriałów w stymulowaniu wzrostu roślin fitoremediacyjnych poprzez dostarczanie cennych składników odżywczych takich jak żelazo czy cynk bezpośrednio do strefy korzeniowej, a także poprzez stymulację aktywności enzymatycznej mikroorganizmów zasiedlających ryzosferę (Alazaiza et al., 2025; Al-Khayri et al., 2023, Ma et al., 2023). Dodatkowo wskazano, że w wyniku wprowadzenia nanocząstek do skażonej gleby, może dochodzić do wzmożonej aktywności enzymów antyoksydacyjnych takich jak peroksydaza askorbinianowa i dysmutaza ponadtlenkowa, co chroni komórki roślinne przed uszkodzeniami oksydacyjnymi i sprzyja utrzymaniu prawidłowego przebiegu procesów metabolicznych w warunkach stresu środowiskowego (Alazaiza et al., 2025, Al-Khayri et al., 2023, Ma et al., 2023). W tym miejscu warto podkreślić, że współczesny rozwój nanotechnologii wykracza daleko poza wykorzystanie nanomateriałów jako biernych dodatków poprawiających właściwości środowiska glebowego. Coraz większe znaczenie pod kątem możliwości wykorzystania w procesach bioremediacyjnych zyskuje perspektywa zaprojektowania tzw. zaawansowanych systemów kontrolowanego dostarczania substancji biologicznie czynnych, opartych na nanonośnikach, które umożliwią precyzyjną enkapsulację oraz stopniowe uwalnianie składników odżywczych, fitohormonów, promotorów wzrostu czy pożytecznych mikroorganizmów ryzosferowych w miejscu do tego wyznaczonym (Ren et al., 2022; Sattar et al., 2022; Ulhassan et al., 2022). Dzięki możliwości modyfikacji właściwości fizykochemicznych nanocząstek możliwe może być również dostosowanie tempa i miejsca uwalniania substancji aktywnych do potrzeb roślin oraz warunków panujących w ryzosferze, co zwiększa potencjał ich efektywnego wykorzystania i ogranicza straty mogące wyniknąć z ich wymywania lub niekontrolowanej dyfuzji (Gomez, 2025). Takie ukierunkowane systemy dostarczania substancji biologicznie czynnych mogą w przyszłości istotnie zwiększać skuteczność biodegradacji i pobierania zanieczyszczeń przez rośliny, jednocześnie ograniczając ryzyko nadmiernego obciążenia środowiska związanego z niekontrolowaną aplikacją substancji wspomagających.

1.5. Charakterystyka gleb z dołów urobkowych

Szczególną kategorię terenów wymagających działań remediacyjnych stanowią obszary związane z historycznym wydobyciem i eksploatacją ropy naftowej, na których występują tzw. doły urobkowe. W miejscach tych w przeszłości deponowano odpady wiertnicze, które w swym składzie zawierały przede wszystkim zwierciny, zużyte płuczki bentonitowe oraz zanieczyszczenia ropopochodne, pochodzące z nawierconych warstw roponośnych. Odpady te częściowo zostały zmieszane z ziemią, torfem itp., gdyż w początkowym okresie rozwoju górnictwa nafty i gazu wiedza ekologiczna była na niskim

poziomie, a co za tym idzie, świadomość zagrożeń stwarzanych przez gromadzone odpady zarówno wśród pracowników branży, jak i zwykłych ludzi była niedostateczna. Substancje ropopochodne stanowią jedno z głównych źródeł skażenia gleby/odpadów zdeponowanych w starych dołach urobkowych, powodując degradację życia biologicznego na terenie składowania i nieprzydatność użytkową, także ze względu na niekorzystne zmiany parametrów wytrzymałościowych gruntu na obszarze składowiska (Steliga, 2009).

Obecność tych substancji w matrycy glebowej prowadzi przede wszystkim do zmian w ilości oraz składzie chemicznym materii organicznej w glebie. Wzrost zawartości węgla organicznego wynikający z obecności frakcji ropopochodnych zaburza stosunek C:N:P, który dla prawidłowego funkcjonowania mikroorganizmów oraz przebiegu biodegradacji zanieczyszczeń powinien wynosić 100:10:1 (Nwankwegu et al., 2016; Steliga et al., 2012; Wu et al., 2017). Takie zaburzenie ogranicza dostępność azotu i fosforu dla mikroorganizmów glebowych i roślin, spowalniając naturalne procesy biologiczne zachodzące w glebie. Obecność zanieczyszczeń ropopochodnych w glebie może również wpływać na jej właściwości fizyczne .Wypełnianie przestrzeni porowej przez substancje oleiste zmniejsza pojemność wodną gleby i utrudnia wymianę gazową pomiędzy glebą a atmosferą. Ograniczona dostępność tlenu w profilu glebowym, w połączeniu ze zwiększonym zapotrzebowaniem mikroorganizmów na tlen, destabilizuje procesy mikrobiologiczne i biochemiczne zachodzące w środowisku glebowym. W rezultacie dochodzi do zaburzeń funkcjonowania mikrobioty oraz zmian w liczebności i różnorodności mikroorganizmów odpowiedzialnych za obieg materii w glebie. Obecność związków ropopochodnych w glebie może prowadzić także do zaburzeń procesów sorpcyjnych, wymiany jonowej czy zmiany odczynu gleby, a w niektórych przypadkach nawet wtórnym zakwaszeniem środowiska glebowego, co dodatkowo pogarsza warunki siedliskowe dla roślin i mikroorganizmów.

Migracja substancji ropopochodnych w głąb profilu glebowego oraz ich przemieszczanie się wraz z wodami opadowymi może prowadzić do wtórnego skażenia otaczających gruntów, a także wód powierzchniowych i podziemnych. Wielorakość i specyficzne własności składników zanieczyszczeń ropopochodnych oraz produktów ich biologicznych i fizykochemicznych przekształceń w środowisku przyrodniczym powodują duże trudności w rozpoznaniu ekologicznych następstw zanieczyszczenia środowiska, wyborze technologii oczyszczania, a także metod diagnozowania i monitorowania postępu oczyszczania środowiska.

Rekultywacja terenów starych dołów urobkowych zdegradowanych przez skażenie substancjami ropopochodnymi wymaga opracowania nowych, prostych i uzasadnionych ekonomicznie do zastosowania w warunkach kopalnianych technologii w celu usunięcia lub ograniczenia ich ujemnego wpływu na środowisko. Likwidacja polegająca na biodegradacji węglowodorów ropopochodnych metodą „*in-situ*" wydaje się być w tym przypadku najkorzystniejszym rozwiązaniem, ponieważ zmniejsza do minimum zagrożenia związane ze skażeniem środowiska ropą naftową.

Jednym z obszarów, na którym znajdują się liczne doły urobkowe, jest teren kopalni Grabownica, zlokalizowanej w południowo-wschodniej Polsce, gdzie zachowane obiekty tego typu pochodzą z okresu intensywnych prac wiertniczych prowadzonych w

latach 1925–1950. Odpady wiertnicze zgromadzone w dołach urobkowych powstały podczas wykonywania płytkich otworów o głębokości poniżej 1000 m ppt., realizowanych z wykorzystaniem prostych technik wiercenia charakterystycznych dla obszaru Podkarpacia. W początkowej fazie prac stosowano metodę udarową, natomiast wraz z pogłębianiem otworów wykorzystywano metodę obrotową z użyciem płuczki bentonitowej, co w istotny sposób wpływało na skład powstających odpadów. W konsekwencji materiał zdeponowany w dołach urobkowych, oprócz urobku skalnego, zawierał również składniki płuczki wiertniczej oraz ropę naftową pochodzącą z przewierconych warstw roponośnych. W tamtych latach procesie wiercenia stosowano wyłącznie niemodyfikowane płuczki iłowe, przygotowywane na bazie wody i minerałów ilastych, najczęściej bentonitu. Z tego względu odpady wiertnicze zdeponowane w dołach urobkowych kopalni Grabownica można było scharakteryzować jako uwodniony, nieorganiczny materiał skalny o wysokiej zawartości frakcji ilastej, zawierający domieszki piasku, kamieni oraz osadów sedymentacyjnych. Materiał ten był jednocześnie zanieczyszczony substancjami ropopochodnymi, które występowały zarówno w postaci rozpuszczonej i zmieszanej, jak i zaadsorbowanej na powierzchni ziaren gruntu.


Odpady wiertnicze/gleba w dołach urobkowych przeznaczone do oczyszczania charakteryzowały się bardzo wysoką zawartością zanieczyszczeń ropopochodnych w zakresie od 143 283 do 341 247 mg/kg s.m., przy czym dla poszczególnych dołów urobkowych wynosiła ona odpowiednio: G-6 (39 458 – 143 283 mg/kg s.m.), G-44 (85 650 – 139 240 mg/kg s.m.), G-70 (85 650 – 189 240 mg/kg s.m.) oraz G-11 (124 584 – 341 247 mg/kg s.m.) (Rys.6d). Z uwagi na złożony charakter zanieczyszczeń oraz wysoki stopień degradacji środowiska glebowego, rekultywacja terenów po dawnych dołach urobkowych stanowiła poważne wyzwanie technologiczne i środowiskowe.

Prowadzone przez wiele lat badania laboratoryjne w skali półtechnicznej w warunkach *ex-situ* oraz prace pilotażowe realizowane w warunkach przemysłowych pozwoliły na opracowanie i wdrożenie przez INiG – PIB (Zakład Technologii Eksploatacji Płynów Złożowych oraz Zakład Mikrobiologii) nowoczesnej, etapowej technologii oczyszczania gleby i ziemi z terenów dołów urobkowych zanieczyszczonej substancjami ropopochodnymi metodą *in-situ*. Weryfikacja opracowanej technologii etapowego odpadów ze starych dołów urobkowych z zanieczyszczeń ropopochodnych w warunkach przemysłowych potwierdziła jej użyteczność i przydatność oraz skuteczność bioremediacji zanieczyszczeń ropopochodnych występujących w odpadach wiertniczych glebie/w wysokich stężeniach. Obejmowała ona następujące etapy (Rys. 7):(Steliga et.al., 2012, 2018).

- remediację wstępną polegającą na drenażu melioracyjno-odciekowym, co pozwoliło na obniżenie zawartości zanieczyszczeń ropopochodnych o 45,2-78,2%, czyli do poziomu poniżej 50 000 mg/kg s.m. umożliwiającego realizowanie dalszych etapów oczyszczania odpadów zgromadzonych w dołach urobkowych;
- modyfikację struktury gleb polegająca na zmieszaniu jej w różnych proporcjach z „czystą ziemią” celem zwiększenia biodostępności mikroorganizmów do substancji biogennych i zanieczyszczeń ropopochodnych, które określono

opierając się na dyfraktogramach rentgenowskich frakcji ilastej, analizie zmian aktywności dehydrogenazowej;


- bioremediacji podstawowej stymulowanej przez biowentylację (napowietrzanie) oraz wzbogacanie oczyszczanego środowiska w składniki biogenne, w dawkach wyznaczonych na podstawie badań laboratoryjnych;
- bioaugmentację polegającą na wprowadzeniu do ekosystemu oczyszczanego terenu aktywnego biopreparatu sporządzonego na bazie wyizolowanych, wyselekcjonowanych i namnożonych niepatogennych gatunków bakterii i grzybów autochtonicznych, którą realizowano w kilku seriach w zależności do oczyszczanego terenu dołu urobkowego.




Teren dołu urobkowego G-44



Teren dołu urobkowego G-6



Teren dołu urobkowego G-11



Teren dołu urobkowego G-70

Rysunek 7 Ilustracja wybranych terenów dołów urobkowych: a) przed rozpoczęciem prac oraz w trakcie: b) drenażu melioracyjno-odciekowego; c) biopremediacji podstawowej; d) inokulacji biopreparatami mikrobiologicznymi

W wyniku prac bioremediacyjnych prowadzonych przez okres 3 lat na terenach powyższych dołów urobkowych osiągnięto istotne obniżenie stężeń zanieczyszczeń ropopochodnych. Wartości stężeń zanieczyszczeń w oczyszczanych gruntach kształtowały się na poziomie: G-44 (12 365 – 22 125 mg/kg s.m.), G-6 (4 320 – 6 062 mg/kg s.m.), G-11 (7 475 – 22 125 mg/kg s.m.) oraz G-70 (5 176 – 9 176 mg/kg s.m.). Ze względu na początkowo wysoką zawartość zanieczyszczeń ropopochodnych osiągnięcie zakładanego poziomu oczyszczenia terenu dołów urobkowych wymaga kontynuacji procesu oczyszczania, natomiast w celu jego głębszego doczyszczenia wskazane jest zastosowanie bardziej zaawansowanych technik bioremediacyjnych.

1.6. Motywacja, hipoteza i cele pracy

Niniejsza rozprawa doktorska została przygotowana w formie spójnego tematycznie zbioru artykułów opublikowanych w czasopismach naukowych. Zaproponowana tematyka pracy wpisuje się w aktualne kierunki rozwoju metod oczyszczania środowiska i jest zgodna przyjętą na lata 2020-2025 polityką INiG – PIB, która za jeden ze strategicznych kierunków rozwoju firmy przyjęła doskonalenie technik rekultywacji i remediacji dołów urobkowych i terenów poeksploatacyjnych oraz bioremediacji gleb zanieczyszczonych substancjami ropopochodnymi.

Jak wspomniano wcześniej, górnictwo nafty i gazu należy do gałęzi przemysłu, które wiążą się ze znaczną ingerencją w strukturę litosfery oraz powstawaniem znacznych ilości odpadów wiertniczych. Wśród nich istotną grupę stanowią odpady zanieczyszczone węglowodorami ropopochodnymi, które w przypadku niewłaściwego zagospodarowania mogą stanowić jedno z głównych źródeł skażenia gleb na terenach związanych z eksploatacją ropy naftowej i gazu ziemnego. Substancje ropopochodne charakteryzują się zdolnością do przemieszczania w profilu glebowym, stosunkowo długim czasem utrzymywania się w środowisku oraz właściwościami toksycznymi, co sprawia, że ich obecność w gruncie prowadzi do długotrwałego zaburzenia funkcjonowania ekosystemów glebowych. Dodatkowo ich wielorakość oraz zróżnicowane właściwości fizykochemiczne powodują, że zarówno dobór odpowiednich metod oczyszczania jak i ocena ich oddziaływania na środowisko stanowią istotne wyzwanie przed którym stoi obecnie branża górnictwa nafty i gazu.

Przykładem obszaru górniczego, na którym INiG-PIB prowadził działania rekultywacyjne, był teren złoża Grabownica, na którym w wyniku wieloletniej działalności wiertniczej powstały liczne doły urobkowe wykorzystywane w przeszłości do deponowania odpadów wiertniczych zanieczyszczonych węglowodorami ropopochodnymi. Długotrwałe składowanie tych odpadów bezpośrednio w obrębie dołów urobkowych doprowadziło do trwałego skażenia gruntów na tym obszarze oraz stworzyło zagrożenie dla środowiska gruntowo-wodnego, co uzasadniało podjęcie działań rekultywacyjnych ukierunkowanych na ograniczenie stopnia zanieczyszczenia i przywrócenie właściwości użytkowych zdegradowanych terenów. Pomimo dostępności wielu metod bioremediacyjnych wykorzystywanych do oczyszczania gleb zanieczyszczonych tego typu związkami, w przypadku gleb pochodzących z terenów dołów urobkowych na obszarze kopalni Grabownica ich skuteczność okazała się niewystarczająca. Opracowana i zastosowana przez INiG – PIB technologia etapowego oczyszczania odpadów wiertniczych zanieczyszczonych związkami ropopochodnymi zdeponowanych w starych dołach urobkowych metodą *in-situ* umożliwiła znaczne obniżenie stężeń węglowodorów ropopochodnych, jednakże po 3-letnim okresie prowadzenia prac bioremediacyjnych nie osiągnięto zakładanego poziomu zanieczyszczeń. Stężenia zanieczyszczeń ropopochodnych w końcowej fazie oczyszczania terenów dołów urobkowych kształtowały się w zakresie 4 025 – 22 125 mg/kg s.m. W związku z tym istniała potrzeba opracowania metod umożliwiających głębsze doczyszczenie gruntów z dołów urobkowych ukierunkowanych na zwiększenie efektywności degradacji zarówno węglowodorów alifatycznych, jak i aromatycznych w glebie, a także na obniżenie jej toksyczności.

W tym kontekście postawiono ***hipotezę, że głębsze doczyszczenie gruntów z dołów urobkowych można będzie osiągnąć poprzez modyfikację istniejących technik bioremediacyjnych i opracowanie zintegrowanych strategii zwiększających efektywność biodegradacji węglowodorów ropopochodnych.***

Celem naukowym pracy była identyfikacja i analiza kluczowych parametrów umożliwiających kompleksową ocenę efektywności zaproponowanych strategii bioremediacji gruntów z dołów urobkowych.

Celem utylitarnym było opracowanie skutecznych i przyjaznych dla środowiska metod oczyszczania gleb zanieczyszczonych węglowodorami ropopochodnymi, umożliwiających przywrócenie właściwości użytkowych zdegradowanych terenów. Badania prowadzone na rzeczywistych próbkach gleb z dołów urobkowych zanieczyszczonych zestarzałymi węglowodorami ropopochodnymi pozwoliły na uzyskanie wiarygodnych wyników, co zwiększa możliwość ich wykorzystania w przyszłości w warunkach przemysłowych.

Cele szczegółowe:

- Ocena wpływu zastosowania dodatków (γ-PGA oraz wytypowanych biosurfaktantów) modyfikujących proces inokulacji biopreparatem opracowanym na bazie bakterii autochtonicznych w celu zwiększenia efektywności biodegradacji węglowodorów ropopochodnych w gruncie z terenu dołu urobkowego.

- Określenie wpływu wzbogacenia biopreparatu bakteryjnego w wybrane szczepy grzybów autochtonicznych na intensyfikację procesów biodegradacji węglowodorów alifatycznych oraz wielopierścieniowych węglowodorów ropopochodnych.

- Wprowadzenie fitoremediacji z wykorzystaniem roślin uprawnych w połączeniu z bioaugmentacją jako strategii umożliwiającej głębsze doczyszczenie gruntów z terenów dołów urobkowych.

- Zastosowanie fitoremediacji z wykorzystaniem roślin dziko rosnących, wspomaganej inokulacją biooreparatem na bazie bakterii autochtonicznych w celu głębszego doczyszczenia gruntów historycznie skażonych węglowodorami ropopochodnymi z terenu dołu urobkowego.

- Scharakteryzowanie społeczności bakteryjnej i orzęskowej w glebie po przeprowadzonych procesach fitoremediacyjnych z wykorzystaniem roślin dziko rosnących.

- Ocena wpływu wybranych nanomateriałów na efektywność procesów bioremediacyjnych wspomaganych bioaugmentacją oraz określenie ich potencjału aplikacyjnego w fitoremediacji gruntu pochodzącego z dołu urobkowego.

- Ocena wpływu współwystępowania polichlorowanych bifenyli (PCB) na przebieg oraz efektywność biodegradacji węglowodorów ropopochodnych.

1.7. Zakres pracy

Praca doktorska została podzielona na następujące części:

Rozdział 1 stanowi wstęp pracy doktorskiej, w którym przedstawiono zagadnienia obejmujące: ogólne informacje na temat zanieczyszczenia środowiska glebowego

węglowodorami ropopochodnymi, charakterystykę zanieczyszczeń ropopochodnych, przegląd strategii prowadzenia remediacji gruntów zanieczyszczonych węglowodorami ropopochodnymi, opis procesów biodegradacji tych związków, omówienie czynników wpływających na efektywność biodegradacji oraz metod wspomagania tego procesu, a także charakterystykę gleb z dołów urobkowych. Rozdział ten przedstawia również motywację do podjęcia zagadnienia, formułuje hipotezę badawczą oraz określa cele i zakres pracy.

Rozdział 2 obejmuje charakterystykę materiałów badawczych, w tym badanych gleb, stosowanych biopreparatów, roślin wykorzystywanych w procesach fitoremediacji oraz substancji zastosowanych w celu zwiększenia efektywności biodegradacji węglowodorów ropopochodnych. W rozdziale 2 omówiono również metody badawcze obejmujące analizy respirometryczne, chromatograficzne, mikrobiologiczne oraz toksykologiczne.

Rozdział 3 przedstawia potencjał modyfikacji procesu inokulacji biopreparatem na bazie bakterii autochtonicznych polegający na wprowadzeniu dodatkowo do gleby γ-PGA, w celu poprawy efektywności bioremediacji gleby pochodzącej z dołu urobkowego, charakteryzującej się wysokim stężeniem zanieczyszczeń ropopochodnych. Zawiera on rozbudowane badania monitoringowe prowadzone w dwóch wariantach wzbogacenia gleby tj. inokulacji biopreparatem oraz biopreparatem w połączeniu z γ-PGA, obejmujące analizy respirometryczne, chromatograficzne i toksykologiczne, na których podstawie określono skuteczność zaproponowanych strategii bioremediacji.

Rozdział 4 rozbudowuje zagadnienie związane z możliwością poprawy efektywności biodegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego poprzez modyfikację procesu inokulacji biopreparatem na bazie bakterii autochtonicznych z wykorzystaniem wybranych biosurfaktantów. Zawiera on kompleksowe wyniki badań laboratoryjnych w układach z sześcioma różnymi biosurfaktantami w dwóch stężeniach, obejmujące analizy respirometryczne, chromatograficzne i toksykologiczne. W rozdziale oceniono skuteczność zaproponowanej strategii bioremediacji gruntu z dołu urobkowego, wskazano biosurfaktant o największym potencjale do zwiększenia biodegradacji zanieczyszczeń w badanej glebie oraz zbadano wpływ stężenia biosurfaktanta w układzie na efektywność procesu oczyszczania.

Rozdział 5 składa się z dwóch części. W pierwszej części przedstawiono wyniki badań biodegradacji gleby z dołu urobkowego poddanej inokulacji biopreparatem na bazie bakterii lub bakterii i grzybów autochtonicznych. W oparciu o przeprowadzone analizy chromatograficzne i toksykologiczne oceniono potencjał wzbogacenia biopreparatu na bazie bakterii autochtonicznych w wybrane szczepy grzybów autochtonicznych w poprawie efektywności oczyszczania gleby zanieczyszczonej węglowodorami ropopochodnymi. W drugiej części omówiono możliwość zastosowana fitoremediacji w celu głębszego doczyszczenia gleb z dołów urobkowych. Omówiono ocenę efektywności fitoremediacji i fitoremediacji wspomaganej bioaugmentacją (biopreparat lub biopreparat z dodatkiem γ-PGA), z wykorzystaniem dwóch gatunków roślin: *Zea mays* i *Echinacea purpurea* na podstawie przeprowadzonych analiz

chromatograficznych i toksykologicznych. W rozdziale 5 zamieszczono również analizę morfologiczną roślin oraz zbadano ilość TPH i WWA w korzeniach i pędach *Echinacea purpurea.*

Rozdział 6 rozbudowuje zagadnienia związane z możliwością zastosowania fitoremediacji wspomaganej bioaugmentacją w celu głębszego doczyszczenia gleb z dołów urobkowych. W rozdziale tym omówiono badania z wykorzystaniem roślin dziko rosnących na terenie naturalnego wycieku ropy naftowej *Scirpus sylvaticus* oraz *Cirsium oleraceum.* Poza standardowymi analizami chromatograficznymi i toksykologicznymi gleby poddanej oczyszczaniu, rozdział zawiera analiz materiału roślinnego obejmującą analizę biomasy, tempa wzrostu roślin oraz zawartości TPH i WWA w korzeniach i pędach. W niniejszym rozdziale przestawiono charakterystykę społeczności bakteryjnej w zremediowanej glebie pod kątem obecności bakterii w ryzosferze przyspieszających degradacje zanieczyszczeń i zwiększających tolerancję roślin na niekorzystne warunki środowiskowe oraz oceny zmian w zbiorowiskach orzęsków mogących pełnić role bioindykatorów glebowych.

Rozdział 7 przedstawia potencjał wykorzystania nanomateriałów GO-NP (nanocząstki tlenku grafenu) i ZnO-NP (nanocząstki tlenku cynku) w procesie bioremediacji gleb z dołów urobkowych. Rozdział składa się z dwóch części. W pierwszej części omówiono wyniki badań biodegradacji z wykorzystaniem inokulacji gleby biopreparatem na bazie bakterii autochtonicznych oraz wzbogacenia gleby w badane nanomateriały w czterech różnych stężeniach. W drugiej części zaprezentowano wyniki badań fitoremediacji z wykorzystaniem *Lollium perenne* wspomaganej bioaugmentacją i nanomateriałami. W artykule przedstawiono rozbudowane wyniki badań respirometrycznych, chromatograficznych i toksykologicznych na podstawie których oceniono skuteczności zaproponowanej strategii bioremediacji pod kątem oczyszczania gleb z dołu urobkowego.

Rozdział 8 zawiera ocenę wpływu współwystępowania polichlorowanych bifenyli (PCB) na przebieg oraz efektywność biodegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego. Omawia wyniki badań biodegradacji prowadzonych w warunkach laboratoryjnych na próbkach sterylnych zanieczyszczonych wyłącznie PCB lub węglowodorami ropopochodnymi oraz dwoma rodzajami zanieczyszczeń tj. PCB i węglowodorami ropopochodnymi. Zawiera również wyniki badań prowadzonych w warunkach półtechnicznych z monitoringiem przebiegu biodegradacji zanieczyszczeń w glebie oraz oceną ekotoksykologiczną gleby w trakcie oczyszczania.

Rozdział 9 stanowi podsumowanie wraz z najważniejszymi wnioskami wynikającymi z wszystkich prac przeprowadzonych w ramach realizacji niniejszej rozprawy doktorskiej.

2. Materiały i metody

2.1. Materiały badawcze

Do badań zaprezentowanych w artykułach załączonych do niniejszej pracy doktorskiej wykorzystano próbki gleb pobrane z terenów historycznie skażonych węglowodorami ropopochodnymi, zlokalizowanych w południowo-wschodniej Polsce, na obszarze kilku dołów urobkowych znajdujących się blisko siebie (doły G6, G11, G44, G70), a także z miejsca skażonego w wyniku naturalnego wycieku ropy naftowej (gleba OS- Artykuł IV) (Tabela 2), po zakończeniu etapowej bioremediacji gruntu (rozdział 1.6). Próbki gleb pobierano z różnych interwałów głębokościowych (0-0,25; 0,25-1; 0,25-1,5 m ppt.), odpowiednio oznakowywano, umieszczano w hermetycznie zamkniętych pojemnikach transportowych i niezwłocznie przewożono do laboratorium Zakładu Technologii Eksploatacji Płynów Złożowych INiG – PIB. W warunkach laboratoryjnych próbki poddawano mieszaniu i homogenizacji w celu uzyskania reprezentatywnego materiału testowego przeznaczonego do dalszych analiz oraz badań biodegradacyjnych.

Po przeprowadzeniu analiz fizyczno-chemicznych, gleby przeznaczone do badań wzbogacano w składniki biogenne, których odpowiednia zawartość była niezbędna dla prawidłowego wzrostu i aktywności mikroorganizmów. Jako optymalny stosunek azotu do fosforu (N:P) w badanych glebach przyjęto 10:1, który został określony na podstawie wcześniejszych badań prowadzonych w Zakładzie Technologii Eksploatacji Płynów Złożowych INiG – PIB (Steliga et al., 2012). Do suplementacji gleb wykorzystano nawóz mineralny „Azofoska" o składzie: 13,6% N całkowitego, 5,5% azotu azotanowego, 8,1% azotu amonowego, 6,4% P_2O_5 rozpuszczalnego, 19,1% K_2O w postaci K_2SO_4, 4,4% MgO w postaci rozpuszczalnego $MgSO_4$ oraz mikroelementy (0,17% Fe, 0,27% Mn, 0,18% Cu, 0,045% Zn, 0,09% Mo), którego dawki obliczano indywidualnie dla każdej z gleb przeznaczonej do badań.

Dodatkowo w celu korekty odczynu gleb stosowano wapno nawozowe w ilości 1,0–1,5 g/kg, aż do uzyskania optymalnego pH mieszczącego się w przedziale 7,5–7,6. Obecność wapnia nie tylko stabilizowała odczyn gleby, ale także zwiększała przeżywalność mikroorganizmów poprzez ochronę ich błon komórkowych przed dezintegrującym wpływem węglowodorów alifatycznych. Ponadto wapń pochodzący z węglanu wapnia sprzyjał tworzeniu struktury granulowanej gleby poprzez koagulację koloidów, co poprawiało warunki powietrzno-wodne i tym samym stwarzało korzystne środowisko dla procesów biologicznego oczyszczania gleb pochodzących z terenów dołów urobkowych (Steliga, 2009).

Tabela 2 Charakterystyka gleb wykorzystanych w badaniach bioremediacji omówionych w artykułach I-VII.

Oznaczenie gleby w artykule	Lokalizacja (dół urobkowy / miejsce skażenia)	Węglowodory ropopochodne		Dodatkowe informacje o zanieczyszczeniach	Artykuł
		TPH [mg/kg s.m.]	WWA [mg/kg s.m.]		
Soil AB	dół urobkowy G11/ (N: 49°40′31.23″, E: 22°04′30.86″)	19 774,23	27,03	obecność BTEX (17,35 mg/kg s.m.)	[I]
Soil G6	dół urobkowy G6 (N: 49°40′16.22″, E: 22°04′42.99″)	6 062,59	12,05		[II,VI]
Soil G6	dół urobkowy G6 (inna część dołu urobkowego)/ (N: 49°40′16.22″, E: 22°04′42.99″)	4 527,70	10,48		[III]
Soil DW	dół urobkowy G70 (N: 49°40′52.03″, E: 22°04′21.92″)	5 176,03	17,71		[IV]
Soil OS	wyciek ropy naftowej	4 025,65	10,30		[IV]
Soil G6	dół urobkowy G6 (inna część dołu urobkowego) (N: 49°40′16.22″, E: 22°04′42.99″)	4 320,70	7,19		[V]
Soil A	gleba wzorcowa zanieczyszczona PCB	-	-	Obecność PCB (13 100 µg/kg s.m.)	[VII]
Soil B	dół urobkowy G44/ (N 54°14′55″, E 46°68′51″)	12 515	–		[VII]
Soil C	dół urobkowy G44/ (N 54°14′55″, E 46°68′51″)	12 365	–	współwystępowanie PCB (13 100 µg/kg s.m.)	[VII]
Soil D	dół urobkowy G44/ (N 54°14′55″, E 46°68′51″)	22 125	–	współwystępowanie PCB (8752 µg/kg s.m.)	[VII]

Do bioaugmentacji gleby stosowano pojedyncze szczepy bakterii (Artykuł VII) oraz biopreparaty opracowane na bazie bakterii (Artykuły I-VII) lub bakterii i grzybów (Artykuły III-IV) autochtonicznych wyizolowanych z terenów dołów urobkowych znajdujących się na terenie kopalni Grabownica. Szczep *Rhodococcus* sp. IN306 wykorzystany w badaniach z artykułu VII został wyizolowany z próbki gleby pobranej na pustyni Negev (Izrael) i wykorzystano go do badań ze względu na duże podobieństwo do

szczepu *Rhodococcus jostii* RHA, znanego ze swej zdolności do rozkładu polichlorowanych bifenyli. (Tabela 3). Wszystkie szczepy (z wyjątkiem jednego) wykorzystane do konstrukcji biopreparatów pochodziły z kolekcji Zakładu Mikrobiologii INiG – PIB i zostały przetestowane pod kątem zdolności do degradacji węglowodorów ropopochodnych. Wykorzystanie w badaniach biopreparatów na bazie mikroorganizmów autochtonicznych wynikało z ich naturalnej adaptacji do obecności węglowodorów ropopochodnych oraz lokalnych warunków glebowych, co sprzyjało zwiększeniu efektywności procesów biodegradacji.

Tabela 3 Biopreparaty wykorzystywane w badaniach bioremediacji gleb z terenów dołów urobkowych omówionych w artykułach I-VII.

Oznaczenie konsorcjum/szczepu w artykule	Skład biopreparatów	Dodatkowe informacje	Artykuł
biopreparat A1	*Dietzia* sp. IN133, *Gordonia* sp. IN138, *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN119, *Rhodococcus* sp. IN136, *Pseudomonas* sp. IN132	biopreparat opracowany na bazie bakterii autochtonicznych	[I]
konsorcjum bakteryjne MC/ biopreparat B1	*Dietzia* sp. IN118, *Gordonia* sp. IN101, *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN119, *Rhodococcus globerulus* IN113, *Raoultella* sp. IN109	biopreparat opracowany na bazie bakterii autochtonicznych	[II–IV, VI]
biopreparat B2	bakterie konsorcjum B1 oraz grzyby: *Aspergillus sydowii*, *Aspergillus versicolor*, *Candida* sp., *Cladosporium halotolerans*, *Penicillium chrysogenum*	biopreparat opracowany na bazie bakterii autochtonicznych wzbogacone w wybrane grzyby autochtoniczne biopreparat bakteryjno-grzybowy	[III-IV]
konsorcjum bakteryjne MC	bakterie należące do rodzajów *Dietzia*, *Gordonia*, *Mycolicibacterium*, *Pseudomonas*, *Raoultella* oraz *Rhodococcus*	biopreparat opracowany na bazie bakterii autochtonicznych	[V]
Mycolicibacterium frederiksbergense IN53	*Mycolicibacterium frederiksbergense* IN53	szczep autochtoniczny	[VII]
Rhodococcus erythropolis IN129	*Rhodococcus erythropolis* IN129	szczep autochtoniczny	[VII]
Rhodococcus sp. IN306	*Rhodococcus* sp. IN306	szczep wyizolowany z gleby pustynnej skażonej węglowodorami ropopochodnymi	[VII]
kultura mieszana M1	*Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306	biopreparat opracowany na bazie trzech szczepów w stosunku 1:1:1	[VII]


Procesy biodegradacyjne były modyfikowane przez zastosowanie różnorodnych substancji chemicznych mających za zadanie zwiększyć efektywność oczyszczania gleb zanieczyszczonych węglowodorami ropopochodnymi. Do modyfikacji procesu inokulacji wykorzystano komercyjnie dostępny kwas γ-poliglutaminowy (γ-PGA), zawierający 10% (w/w) γ-PGA. γ-PGA jest naturalnym produktem niektórych bakterii i archeonów, w tym przede wszystkim gatunków rodzaju *Bacillus*, i wykorzystywany był ze względu na właściwości biosurfaktantowe (Artykuły I-IV). Poza tym stosowano biosurfaktanty, takie jak rhamnolipidy i surfaktyna, komercyjne preparaty JBR 425 i JBR 320 (mieszaniny rhamnolipidów) oraz autorską mieszaninę PSR, opracowaną na bazie γ-PGA, rhamnolipidów i surfaktyny w równych proporcjach objętościowych (Artykuł II). Badania biodegradacji wspomagano również poprzez wzbogacenie gleby w nanocząstki tlenku grafenu (GO-NP) oraz tlenku cynku (ZnO-NP), które wykazują zdolność stymulowania aktywności mikroorganizmów degradujących węglowodory oraz zwiększania biodostępności zanieczyszczeń ropopochodnych (Artykuł VI).

Do fitoremediacji stosowano dwie kategorie roślin fitoremediacyjnych, mianowicie gatunki uprawne, takie jak *Zea mays* (Artykuł III), *Echinacea purpurea* (Artykuł IV), *Lollium perenne* (Artykuł VI), jak i rośliny dziko rosnące *Scirpus sylvaticus* oraz *Cirsium oleraceum* (Artykuł V) pobrane z terenu skażonego ropą naftową zlokalizowanego w pobliżu dołów urobkowych.

Szczegółowa charakterystyka zastosowanych w badaniach biopreparatów, substancji wspomagających oraz roślin fitoremediacyjnych została przedstawiona w publikacjach stanowiących podstawę niniejszej rozprawy doktorskiej (Artykuły I–VII).

2.2. Metody badawcze

Badania bioremediacji prowadzono w dwóch skalach eksperymentalnych: laboratoryjnej oraz półtechnicznej. Eksperymenty w skali laboratoryjnej realizowano w układach zamkniętych z wykorzystaniem zestawu OxiTop, który umożliwiał prowadzenie analiz w ściśle kontrolowanych warunkach procesu. Badania w skali półtechnicznej obejmowały doświadczenia biodegradacji prowadzone metodą pryzmowania *ex situ* (Rys. 8a) oraz badania fitoremediacji wspomaganej bioaugmentacją realizowane metodą wazonową (Rys. 8b) w warunkach zbliżonych do rzeczywistych. W badaniach stosowano różne strategie wspomagania procesów bioremediacji, w szczególności bioaugmentację z wykorzystaniem biopreparatów opracowanych na bazie mikroorganizmów autochtonicznych oraz fitoremediację, a także dodatkowe czynniki stymulujące procesy degradacji zanieczyszczeń takie jak biosurfaktanty, czy nanomateriały.




Rysunek 8 Schemat stanowisk badawczych do prowadzenia eksperymentów a) biodegradacji, b) fitoremediacji w skali półtechnicznej (opracowanie własne na podstawie Wojtowicz i in., 2022, Wojtowicz i in. 2023b poddane obróbce przez Open AI).

Do oceny efektywności procesów bioremediacji wykorzystywano analizy respirometryczne oraz analizy chromatograficzne umożliwiające oznaczanie zawartości węglowodorów ropopochodnych w próbkach. Uzupełnieniem oceny skuteczności prowadzonych procesów była analiza potencjalnego oddziaływania badanych gleb na organizmy żywe. W tym celu wykonywano testy ekotoksykologiczne z wykorzystaniem organizmów bioindykatorowych należących do wszystkich poziomów troficznych tj. producentów, konsumentów i reducentów. Szczegółową metodykę zastosowanych analiz przedstawiono w kolejnych podrozdziałach niniejszej pracy oraz artykułach załączonych do pracy.

2.2.1. Analizy respirometryczne

Analizy respirometryczne wykonywano w celu oceny intensywności procesów biodegradacji zanieczyszczeń węglowodorowych w glebie. Metoda polegała na pomiarze zużycia tlenu przez mikroorganizmy glebowe w trakcie rozkładu związków organicznych, co umożliwiało ocenę aktywności metabolicznej mikroorganizmów oraz przebiegu procesów biodegradacji (Malińska, 2016). Badania prowadzono z wykorzystaniem zestawu respirometrycznego OxiTop Control firmy WTW (Rys. 9a), który umożliwia bezpośredni pomiar zużycia tlenu w układzie zamkniętym na podstawie zmian ciśnienia w przestrzeni gazowej naczynia pomiarowego. Próbki gleby umieszczano w naczyniach pomiarowych, do których wprowadzano biopreparat mikrobiologiczny, stosowany samodzielnie lub w połączeniu z dodatkowymi czynnikami wspomagającymi procesy biodegradacji. Równolegle przygotowywano próbki kontrolne. Naczynia pomiarowe szczelnie zamykano główkami pomiarowymi systemu OxiTop, a następnie inkubowano w warunkach kontrolowanej temperatury przez cały czas trwania eksperymentu (Steliga & Uliasz, 2014; Steliga & Wojtowicz, 2016;.Wojtowicz et al., 2020b).



Rysunek 9 Zestaw Oxi-Top wykorzystywany do badań respirometrycznych.

W trakcie eksperymentu główki pomiarowe rejestrowały zmiany ciśnienia w układzie zamkniętym w regularnych odstępach czasu. Zebrane dane były przesyłane za pomocą interfejsu IR do kontrolera OxiTop OC 110 (Rys. 9b), a następnie poddawano je obróbce graficznej i statystycznej przy użyciu programu Achat OC. Przetwarzanie mierzonego ciśnienia na wartość zużytego tlenu (mO_2) wykonywano według wzoru (Steliga & Uliasz, 2014):

$$\Delta O_2 = \frac{M(O_2) \cdot V_g \cdot R \cdot T_m}{R \cdot T_0 \cdot \alpha} \cdot \Delta p$$

gdzie: $M(O_2)$ – masa molowa tlenu [kg/mol], Vg – objętość wolnego gazu [m^3], R – stała gazowa [$J \cdot mol^{-1} \cdot K^{-1}$], Tm – wartość pomiarowa temperatury [K], T_0 – temperatura odniesienia (273,15 K), α – współczynnik absorpcji (0,03103), Δp – spadek ciśnienia w próbie [Pa].

Wyniki przedstawiano w postaci krzywych zależności zużycia tlenu od czasu trwania eksperymentu, co umożliwiało ocenę dynamiki procesów biodegradacji w analizowanych układach reakcyjnych.

2.2.2. Analizy chromatograficzne

Analizy chromatograficzne wykonano w celu oceny skuteczności bioremediacji gleb i gruntów pobranych z terenu dołów urobkowych. Ocenę przeprowadzono na podstawie porównania stężeń wybranych zanieczyszczeń przed i po zastosowaniu zaproponowanych strategii oczyszczania. Ponadto w przypadku badań fitoremediacji wykorzystano je do oceny akumulacji zanieczyszczeń w roślinach, a tym samym określenia stopnia ich pobierania i gromadzenia w korzeniach i pędach zastosowanych roślin fitoremediujących. Badania prowadzono z wykorzystaniem techniki chromatografii gazowej (GC), wykorzystywanej do analizy TPH, n-alkanów, BTEX i PCB, oraz wysokosprawnej chromatografii cieczowej (HPLC), stosowanej w oznaczaniu WWA. Dla wszystkich analizowanych grup związków (TPH, WWA, BTEX, PCB) stosowano autorskie metodyki analityczne opracowane w Zakładzie Technologii Eksploatacji Płynów Złożowych INiG – PIB (Steliga et al., 2012; Wojtowicz, 2018; 2022; Wojtowicz & Jakubowicz 2019), które umożliwiły na:

- sumaryczne oznaczenie zawartości TPH, WWA, BTEX i PCB w badanej glebie/gruncie oraz obliczenie stopnia ich biodegradacji w wyniku zaproponowanej strategii oczyszczania,
- identyfikację poszczególnych n-alkanów, węglowodorów aromatycznych oraz kongenerów PCB obecnych w badanej glebie/gruncie oraz określenie zmian ich stężeń w trakcie lub po zakończeniu oczyszczania, co umożliwiało ocenę stopnia ich biodegradacji, a tym samym efektywności zaproponowanych technik bioremediacyjnych,
- identyfikację węglowodorów z grupy izoprenoidów (pristan i fitan), należących do związków trudno biodegradowalnych, co pozwalało na określenie stopnia biodegradacji węglowodorów ropopochodnych (n-alkanów) na podstawie zmian wartości wskaźników biodegradacji n-C_{17}/Pr oraz n-C_{18}/F (Rosa et al., 2007; Steliga, 2009),
- opracowanie matematycznego modelu biodegradacji zanieczyszczeń ropopochodnych oraz wyznaczenie stałych szybkości biodegradacji zanieczyszczeń w badanej glebie/gruncie (Steliga et al., 2020),
- sumaryczne oznaczenie zawartości TPH i WWA w korzeniach oraz pędach roślin stosowanych w badaniach fitoremediacji.

Ogólne warunki przygotowania próbek oraz parametry analiz chromatograficznych zestawiono w tabeli 4, natomiast pełne opisy metodyki chromatograficznego oznaczania zanieczyszczeń ropopochodnych w materiale badawczym przedstawiono w publikacjach stanowiących podstawę niniejszej rozprawy doktorskiej (Artykuły I-VII).

Tabela 4 Zestawienie najważniejszych parametrów prowadzenia oznaczeń z wykorzystaniem analiz chromatograficznych


Oznaczane związki	Przygotowanie próbki	Technika / Aparatura	Kolumna chromatograficzna	Detektor	Warunki analizy	Artykuł
TPH (n-alkany, pristan, fitan)	• ekstrakcja rozpuszczalnikowa • ekstrakcja dichlorometanem (3 × 20 mL) z zastosowaniem ultradźwięków, • oczyszczanie ekstraktu na kolumnach SPE Florisil	GC, Clarus 500 (Perkin Elmer)	RTX-1 (30 m × 0,53 mm, Restek)/	FID	gaz nośny He 20 mL/min; temp. inżektora 290 °C; temp. detektora 320 °C; program pieca: 30 °C (2 min), 30–105 °C (10 °C/min), 105–285 °C (5 °C/min), 285 °C (5 min)	I-VII
WWA	• ekstrakcja metodą QuEChERS • ekstrakcja mieszaniną woda/ acetonitryl/ mieszanina ekstrakcyjna $MgSO_4$/NaCl • oczyszczanie dSPE ($MgSO_4$ + PSA, JO3930 Clean-up Kit, • filtracja przez filtr strzykawkowy	HPLC, Vanquish Core (Thermo Scientific)	NUCLEODUR C18 PAH (125 × 4 mm, 3 µm, Macherey-Nagel)	UV-VIS + FLD	faza ruchoma: metanol / acetonitryl; przepływ 1,5 mL/min; gradient: 20–50% B (1,5 min), 50–100% B (1 min), 100% B (1 min), powrót do 0% B (3 min), 100% A (3 min)	I-VI
BTEX	• ekstrakcja z fazy nadpowierzchniowej • próbka gleby termostatowana przed poborem fazy gazowej	GC-HS, Clarus 500 GC-FID z przystawką TurboMatrix 16 (Perkin Elmer)	RT-TCEP (60 m × 0,25 mm, Restek)	FID	temp. inżektora 200 °C; temp. detektora 280 °C; program pieca: 60 °C (5 min), 60–100 °C (5 °C/min), 100 °C (10 min)	I
PCB	• ekstrakcja rozpuszczalnikowa • ekstrakcja mieszaniną n-heksan/dichlorometan (1:1); • oczyszczanie ekstraktu na kolumnach krzemionkowych; • dodatek miedzi w celu usunięcia związków siarki	GC, Clarus 500 (Perkin Elmer)	RTX-XLB (30 m × 0,32 mm, Restek)	ECD	gaz nośny He 3,5 mL/min; temp. inżektora 280 °C; temp. detektora 320 °C; program pieca: 120 °C (2 min), 120–280 °C (15 °C/min), 280 °C (5 min), 280–320 °C (28 °C/min), 320 °C (10 min)	VII

2.2.3. Ocena ekotoksykologiczna

Kompleksowa ocena efektywności zaprezentowanych w artykułach I-VII zmodyfikowanych strategii biologicznego oczyszczania gleb zanieczyszczonych węglowodorami ropopochodnymi wymagała przeprowadzenia oceny ekotoksykologicznej. Miało to na celu sprawdzenie czy w wyniku zaproponowanych strategii bioremediacji nie dochodzi do wtórnego zanieczyszczenia oczyszczanych gleb np. w wyniku powstawania wtórnych metabolitów pośrednich nierzadko wykazujących większe właściwości toksyczne niż pierwotne zanieczyszczenie. Takie związki, mogły nie zostać wykryte podczas rutynowych analiz fizyczno-chemicznych oraz chromatograficznych, dlatego w celu potwierdzenia bezpieczeństwa środowiskowego opracowanych w ramach pracy doktorskiej metod bioremediacji gruntów z terenów dołów urobkowych, zdecydowano się zastosować pakiet testów toksykologicznych opartych na analizie bioindykacyjnej. Analiza bioindykacyjna wykorzystuje organizmy żywe jako wskaźniki zmian zachodzących w środowisku, a ich reakcja odzwierciedla łączne oddziaływanie substancji toksycznych oraz ich interakcje z czynnikami środowiskowymi. W każdym z załączonych w pracy artykułów weryfikacji przyjętej strategii bioremediacji dokonano z zastosowaniem komercyjnych testów Phytotoxkit™, Ostracodtoxkit F™ oraz Microtox® SPT. Dodatkowo w artykułach I i VII ocenę ekotoksykologiczną rozszerzono o wyniki testu oceny mutagenności Ames, natomiast w artykułach IV-V oraz VII o wyniki testu oceny ryzyka środowiskowego MARA.

2.2.3.1. Test fitotoksyczności Phytotoxkit™


Oceny toksyczności gleby na poziomie producentów dokonano za pomocą testu fitotoksyczności Phytotoxkit™ (Rys. 10a), który należy do testów oceny toksyczności ostrej i umożliwia ocenę wpływu badanej gleby na kiełkowanie oraz zahamowanie wzrostu korzeni roślin bioindykatorowych (*Sorghum saccharatum*, *Lepidium sativum*, *Sinapis alba*) (Baran i Tarnawski, 2013; Blinova et al., 2013; Mamindy-Pajany et a., 2011). Test prowadzono zgodnie z instrukcją dostarczoną przez producenta i przebiegał następująco. Na dolną część polistyrenowej płytki nanoszono nawodnioną próbkę gleby, a następnie przykrywano ją warstwą papierowego filtra. Następnie w jednej linii, w równych odstępach, umieszczano po 10 nasion danej rośliny testowej, płytkę zamykano pokrywką i umieszczano w cieplarce zapewniając stałą temperaturę (25°C) przez cały czas trwania testu (72 godziny). Po zakończeniu inkubacji za pomocą skanera wykonano zdjęcia płytek na tle ramki kalibracyjnej oraz przeprowadzono analizę wyników obejmującą obliczenie procentu wykiełkowanych nasion, pomiar długości korzeni każdej rośliny testowej oraz obliczenie procentowego zahamowania wzrostu korzeni względem próbki kontrolnej (Rys. 10b). (Steliga et al., 2018; www.Tigret.eu; Wojtowicz et al., 2023a, 2023b; 2025)



Rysunek 10 a) Składniki testu Phytotoxkit™ oraz b) przykładowy obraz wzrostu roślin testowych na próbce kontrolnej (www.Tigret.eu)

2.2.3.2. Test bezpośredniego kontaktu Ostracodtoxkit F™


Oceny toksyczności gleb na poziomie konsumentów dokonano za pomocą testu Ostracodtoxkit F™ (Rys 11a), który służy do oceny toksyczności chronicznej w układzie bezpośredniego kontaktu z badaną próbką. Test wykorzystuje jako organizmy wskaźnikowe skorupiaki z gatunku *Heterocypris incongruens* (Rys. 11b) i polega na ocenie toksyczności gleby poprzez analizę przeżywalności oraz zahamowania wzrostu małżoraczków (Niyommaneerat et al., 2017; Rouvalis et al., 2013; Steliga et al., 2018). Procedura wykonywania testu przebiegała zgodnie z instrukcją dostarczoną przez producenta. Test prowadzono w polistyrenowych, 6-dołkowych mikropłytkach, przy czym do każdego dołka wprowadzano zawiesinę glonów jako standardową pożywkę (2 ml) oraz po 10 osobników testowych. Do szeregu A mikropłytki wprowadzano 0,5 g gleby kontrolnej, natomiast do pozostałych rzędów po 0,5 g badanej próbki gleby. Inkubację prowadzono przez 6 dni w cieplarce zapewniając stałą temperaturę 25°C. Po zakończeniu testu oceniano przeżywalność organizmów poprzez określenie liczby żywych osobników w każdym dołku mikropłytki oraz analizowano ich wzrost na podstawie pomiaru długości (Steliga et al., 2018; www. Tigret.eu; Wojtowicz et al., 2023a; 2023b; 2025).



Rysunek 11 a) Składniki testu Ostracodtoxkit oraz b) powiększony obraz organizmów wskaźnikowych Heterocypris incongruens (www.Tigret.eu)

2.2.3.3. *Microtox® Solid Phase Test*

Do oceny toksyczności na poziomie reducentów zastosowano test Microtox® Solid Phase (SPT) (Rys 12a), wykorzystujący liofilizowane bakterie luminescencyjne *Vibrio fischeri* (Rys. 12b) jako organizmy wskaźnikowe. Zmiana intensywności luminescencji bakterii stanowiła miarę ich aktywności metabolicznej i ulegała obniżeniu w obecności substancji toksycznych (Foucault et al., 2013; Lima et al., 2011; Oleszczuk et al., 2014). Oznaczenie przeprowadzono w układzie Microtox® Solid Phase, umożliwiającym bezpośredni kontakt bakterii z badaną próbką gleby. W tym celu w naczyniach testowych umieszczano próbki gleby, a następnie dodawano roztwór chlorku sodu aby uzyskać warunki osmotyczne odpowiednie dla bakterii luminescencyjnych. Następnie do układu dodano wykorzystane w teście bioindykatory i prowadzono inkubację zgodnie ze standardową procedurą SDI z użyciem analizatora Delta Tox, wykonując trzy powtórzenia dla każdej próbki. Po 15 minutach inkubacji odczytywano procent efektu toksycznego (PE), który stanowił podstawę oceny toksyczności próbek. W przypadku próbek wykazujących istotny efekt toksyczny wykonywano dodatkową analizę w układzie rozcieńczeń zgodnie z procedurą testu Microtox®, w celu wyznaczenia wartości EC_{50}. W tym etapie badano spadek luminescencji w kolejnych rozcieńczeniach próbki, a wyniki opracowywano z wykorzystaniem oprogramowania SDI. Na podstawie uzyskanych danych wyznaczano wartość EC_{50}, odpowiadającą stężeniu próbki powodującemu 50% efektu toksycznego, którą następnie przeliczano na jednostki toksyczności TU, co umożliwiało klasyfikację próbek według przyjętej skali toksyczności środowiskowej (Steliga et al., 2018; www. Tigret.eu; Wojtowicz et al., 2023a; 2023b; 2025).




Rysunek 12 a) Aparat DeltaTox do wykonywania oznaczeń (polowa wersja systemu Microtox®) oraz b) luminescencja komórek bakterii Vibrio fischeri (prezentacja-Cohiba)

2.2.3.4. *MARA (Microbial Assay for Risk Assessment)*

Do oceny chronicznej toksyczności próbek środowiskowych zastosowano test MARA (Microbial Assay for Risk Assessment), który umożliwia jednoczesną ocenę

reakcji wielu organizmów wskaźnikowych w ramach jednego oznaczenia, przy niewielkim zapotrzebowaniu na próbkę oraz bez konieczności stosowania zaawansowanej aparatury laboratoryjnej (Steliga, 2009; Steliga at al., 2020). Oznaczenie wykonywano w 96-dołkowych mikropłytkach zawierających liofilizowane mikroorganizmy rozmieszczone w kolumnach 1–11, natomiast kolumna 12 pełniła funkcję kontroli negatywnej. Do poszczególnych rzędów wprowadzano różne stężenia badanej próbki, a następnie płytki inkubowano przez 18 godzin w temperaturze 30°C. Po zakończeniu inkubacji oceniano stopień zahamowania wzrostu mikroorganizmów na podstawie redukcji barwnika tetrazoliowego (z ang. *tetrazolium red*), a następnie wykonywano skanowanie płytki i analizę obrazu z wykorzystaniem dedykowanego oprogramowania MARA. Na tej podstawie wyznaczano wartości EC_{50} dla poszczególnych szczepów oraz obliczano średnią wartość MTC (*Mean Toxicity Concentration*), określającą ogólną toksyczność próbki. Profil toksyczności (*fingerprint*) stanowił zestawienie reakcji wszystkich mikroorganizmów testowych i umożliwiał porównawczą ocenę charakteru toksycznego badanych próbek (Steliga, 2009; Steliga at al., 2020; www.Tigret.eu (Rys. 13).




Rysunek 13 Schemat przeprowadzenia procedury testu MARA (Steliga et al., 2020)

2.2.3.5. Ames

Do oceny gleby pod kątem obecności związków mutagennych zastosowano test Muta-Chromoplate™ Kit, stanowiący zmodyfikowaną wersję testu Amesa. Metoda ta opiera się na zjawisku mutacji powrotnych u szczepów *Salmonella typhimurium*, które

pierwotnie wykazują auksotrofię względem histydyny na skutek mutacji w genach odpowiedzialnych za jej syntezę. Najczęściej stosowane szczepy testowe, takie jak TA98 i TA100, umożliwiają wykrywanie różnych typów mutacji, w tym przesunięć ramki odczytu oraz substytucji par zasad, a w kontroli dodatniej wykorzystuje się mutageny, takie jak 2-nitrofluoren i azydek sodu. W obecności czynników mutagennych może dochodzić do rewersji tych mutacji, co prowadzi do odzyskania zdolności syntezy histydyny i umożliwia wzrost komórek w środowisku jej pozbawionym. Liczba takich zdarzeń stanowi miarę aktywności mutagennej badanej próbki (Steliga, 2009). W odróżnieniu od klasycznego testu Amesa, który prowadzony jest na podłożu stałym i opiera się na zliczaniu kolonii rewertantów, test Muta-Chromoplate™ wykorzystuje podłoże płynne oraz płytki 48-studzienkowe, zgodnie z zasadą testu fluktuacyjnego. Ocena wyniku nadal odnosi się do częstości występowania rewertantów, jednak zamiast bezpośredniego liczenia kolonii analizuje się liczbę studzienek, w których zaszła mutacja. Takie podejście zwiększa czułość oznaczenia oraz ogranicza wpływ zmienności związanej ze wzrostem kolonii na podłożu stałym. Dodatkową zaletą jest uproszczona, ustandaryzowana procedura, umożliwiająca wykonywanie oznaczeń także w mniej wyspecjalizowanych laboratoriach. Wynik testu oceniany jest na podstawie zmiany barwy podłoża. Kolor żółty wskazuje na wystąpienie mutacji powrotnej, natomiast purpurowy oznacza jej brak. Analiza liczby studzienek, w których zaszła mutacja indukowana badaną próbką, w porównaniu z częstością mutacji spontanicznych, umożliwia statystyczną ocenę mutagenności przy użyciu oprogramowania dostarczonego przez producenta (Steliga 2009; Steliga et al., 2020) (Rys. 14).



Rysunek 14 Schemat procedury testu Ames (Steliga, 2009; Steliga et al., 2020).

2.2.4. Analizy mikrobiologiczne

2.2.4.1. Izolacja i selekcja bakterii zdolnych do rozkładu związków ropopochodnych

Mikroorganizmy izolowano z próbek gleby pobranych z dołów urobkowych przed rozpoczęciem pierwszych prac remediacyjnych, wykorzystując podłoże mineralne z dodatkiem agaru (15 g/l) oraz ropy naftowej jako źródła węgla. W celu uzyskania możliwie dużej różnorodności mikrobiologicznej analizowano po 10 prób (min. 100 g) pobranych z różnych miejsc każdego obiektu. Wybrane kolonie przenoszono na podłoże płynne o analogicznym składzie, co pozwalało wyeliminować mikroorganizmy wykorzystujące zanieczyszczenia obecne w agarze lub resztkowe związki organiczne z prób środowiskowych. Następnie prowadzono wielokrotny posiew redukcyjny (3–7 powtórzeń) w celu uzyskania czystych kultur, które włączano do banku szczepów Zakładu Mikrobiologii INiG – PIB. Wstępna charakterystyka obejmowała ocenę podstawowych cech fizjologicznych, takich jak ruchliwość, zdolność wzrostu w warunkach tlenowych i beztlenowych, zakres temperatur (4–40°C), tolerancja pH oraz zasolenia (do 10% NaCl). Badania szczegółowe obejmowały zdolność wykorzystywania ropy naftowej oraz wybranych węglowodorów alifatycznych i aromatycznych jako jedynego źródła węgla. Następnie analizowano zdolność wzrostu na n-oktadekanie oraz innych reprezentatywnych węglowodorach o zróżnicowanej strukturze i właściwościach fizykochemicznych (izo-nonadekan, toluen, ksylen, naftalen, antracen, fenantren, fluoren, fluoranten, piren). Dobór warunków inkubacji dostosowywano do charakteru badanych związków (lotność, stan skupienia), stosując m.in. hodowle w atmosferze węglowodorów lub ich bezpośredni dodatek do podłoża. Do dalszych badań wybierano szczepy zdolne do degradacji zarówno węglowodorów alifatycznych, jak i aromatycznych, ze szczególnym uwzględnieniem szerokiego spektrum działania (Steliga, 2009; Steliga i in., 2012; Brzeszcz, 2017).

Identyfikację mikroorganizmów przeprowadzono zarówno metodami klasycznymi, jak i molekularnymi. Cechy charakterystyczne poszczególnych szczepów ustalono na podstawie (Steliga, 2009; Brzeszcz, 2017):

- morfologii kolonii: kształt i barwa kolonii wyrosłych na podłożu agarowym),
- obserwacji mikroskopowych: kształt komórek, konidioforów i konidiów, ocena ruchliwości, barwienie metodą Grama, Ziehl-Nielsena, barwienie spor Dornera w modyfikacji Snydera, barwienie otoczek,
- wzrostu na selektywnych podłożach,
- profili biochemicznych (testy API Coryne i 32GN bioMerieux).

W celu uzyskania potwierdzenia wyników dla wyizolowanych mikroorganizmów przeprowadzono sekwencjonowanie genu (kodującego) 16S rRNA u bakterii. DNA genomowy izolowano na podstawie zmodyfikowanej metody Marmura wykorzystującej ekstrakcję roztworem fenol/chloroform/alkohol izoamylowy (stosunek objętościowy 25:24:1). Następnie zastosowano metodę enzymatycznej amplifikacji (PCR-I – ang. *polymerase chain reaction*, czyli *reakcji łańcuchowej polimerazy*) fragmentów DNA przy użyciu starterów flankujących ok. 500 nukleotydowy fragment badanego genu i oceniano

długość fragmentów DNA na żelu agarozowym. Kolejny etap stanowiła enzymatyczna reakcja sekwencjonowania (PCR-II) z wykorzystaniem zestawu do sekwencjonowania BigDye Terminator v 3.0 Ready Reaction Cycle Sequencing Sequencing Kit firmy Amersham Biosystems (Brzeszcz, 2017). Analizę sekwencjonowania wykonano przy użyciu sekwenatora kapilarnego ABI Prizm® 3100 Genetic Analyser, a sekwencje składano w całość przy pomocy programu ze strony http://arep.med.harvard.edu/labgc/adnan/projects/Utilities/revcomp.html. Identyfikacja szczepów bakteryjnych była możliwa dzięki bazie danych sekwencji 16S rDNA MicroSeq, jak również dzięki programowi BLAST, znajdującemu się na stronie National Center of Biotechnology Information pod adresem http://www.ncbi.nlm.nih.gov/ BLAST (Brzeszcz, 2017; Steliga, 2009; Steliga et al., 2012).

2.2.4.2. Identyfikacja grzybów metodą MAlDI-TOF

Identyfikację wyizolowanych szczepów grzybów przeprowadzono z wykorzystaniem spektrometrii mas z desorpcją/ jonizacją laserową wspomaganą matrycą (MALDI-TOF MS) w Jagiellońskim Centrum Innowacji (Life Science Park, ul. Bobrzyńskiego 14, 30–348 Kraków).

Do przygotowania próbek wykorzystano kolonie pochodzące ze świeżej, nocnej hodowli. Materiał biologiczny zawieszano w 300 µl wody podwójnie destylowanej, a następnie dodawano 900 µl etanolu i całość dokładnie mieszano. Po odwirowaniu próbki przez 2 minuty przy 9000 × g usuwano supernatant. Kolejnym etapem było przeprowadzenie ekstrakcji białek poprzez dodanie do osadu grzybów 40 µl 70% roztworu wodnego kwasu mrówkowego, dokładne wymieszanie próbki oraz dodanie 40 µl acetonitrylu. Po ponownym wirowaniu przy 9000 × g przez 2 minuty pobierano 1 µl supernatantu zawierającego ekstrakt grzybowy i nanoszono go do dołka 96 dołkowej płytki MALDI Biotarget, pozostawiając do wyschnięcia w temperaturze pokojowej. Następnie próbkę pokrywano 2 µl roztworu matrycy MALDI zawierającej kwas α-cyjano-4-hydroksycynamonowy i ponownie suszono na powietrzu. Pomiary prowadzono z wykorzystaniem systemu MALDI BioTyper 2.0 Microflex LT firmy Bruker, zgodnie z zaleceniami producenta. Uzyskane widma białkowe importowano do oprogramowania MALDI BioTyper 2.0, gdzie poddawano je analizie z zastosowaniem standardowego algorytmu dopasowania do widm referencyjnych zgromadzonych w bazie danych BioTyper. W procesie identyfikacji uwzględniano parametry obejmujące stosunek masy do ładunku oraz intensywność pików, a uzyskany wynik dopasowania stanowił podstawę do uszeregowania wyników identyfikacji poszczególnych szczepów (Wojtowicz et al. 2023a, 2023b).

2.2.4.3. Sposób przygotowania biopreparatów

Na podstawie badań przeprowadzonych z wykorzystaniem klasycznych oraz molekularnych technik analitycznych wytypowano najbardziej obiecujące szczepy mikroorganizmów, które następnie wykorzystano do przygotowania biopreparatów przeznaczonych do badań biodegradacji. Kluczowym etapem było dobranie takich mikroorganizmów, które nie tylko są zdolne do efektywnej degradacji substancji ropopochodnych, lecz także nie wykazują względem siebie reakcji antagonistycznych. W tym celu z każdego obszaru dołów urobkowych wyizolowano możliwie szerokie

spektrum mikroorganizmów. Izolaty te, jak wspomniano wcześniej, były przechowywane w kolekcji Zakładu Mikrobiologii INiG – PIB i stanowiły podstawę do przygotowania biopreparatów wykorzystywanych w procesie inokulacji.

Do przygotowania biopreparatów każdy szczep hodowano osobno lub w bulionie wzbogaconym (BD Difco™ lub Biomaxima S.A) dodatkiem octanu sodu (2 g/L) – bakterie (Steliga et al., 2020; Wojtowicz et al., 2022; 2023b), albo w płynnym podłożu Czapek-Dox (BD Difco™) – grzyby (Wojtowicz et al., 2023b). Szczepy inkubowano w temperaturze pokojowej (25°C) na wytrząsarce (150 obr./min.), aż do otrzymania liczebności na poziomie $1 \cdot 10^8$ – $1 \cdot 10^9$ jtk/mL (od 24 -72 godzin). Liczebność weryfikowano poprzez wysianie mikroorganizmów na takie samo podłoże tylko z dodatkiem 15 g agaru. Następnie łączono jednakowe objętości wybranych szczepów w jedną całość i tak otrzymane biopreparaty utrzymywano w bulionie wzbogaconym z dodatkiem octanu sodu (2 g/L) - gdy były obecne tylko bakterie i sacharozy (2 g/L) – jeśli obecne były też grzyby (Brzeszcz, 2017; Steliga et al., 2012;).

2.2.4.4. Sekwencjonowanie nowej generacji (NGS – ang. Next-Generation Sequencing)

Izolację DNA, sekwencjonowanie 16S rDNA i całą analizę społeczności bakteryjnej gleby po fitoremediacji omówioną w artykule V wykonano w firmie Genomed.S.A, (ul. Ponczowa 12, 02-971 Warszawa).

Analiza metagenomiczna populacji bakterii i archeonów została przeprowadzona na bazie hiperzmiennego regionu V3-V4 genu 16S rRNA. Do amplifikacji wybranego regionu i przygotowania biblioteki zostały użyte selektywne sekwencje starterów 341F oraz 785R (Klindworth et al., 2013). PCR przeprowadzono z użyciem Q5 Hot Start High-Fidelity 2X Master Mix, warunki reakcji zgodnie z zaleceniami producenta. Sekwencjonowanie odbyło się na aparacie MiSeq, w technologii paired-end (PE), 2x300 nt, z użyciem zestawu v3 Illuminy. Automatyczna wstępna analiza danych została przeprowadzona na sekwenatorze MiSeq z użyciem oprogramowania MiSeq Reporter (MSR) v2.6. Analiza składała się z dwóch etapów:

1. automatyczne demultipleksowanie próbek,
2. generowanie plików fastq, zawierających surowe odczyty.

Analiza bioinformatyczna zapewniająca klasyfikację odczytów do poziomu gatunku została przeprowadzona pakietem oprogramowania QIIME 2 w oparciu o bazę sekwencji referencyjnych Silva 138. Został również wykorzystany pakiet DADA2 (Callahan et al., 2016), który pozwolił na wyszczególnienie sekwencji pochodzenia biologicznego od tych nowo powstałych w procesie sekwencjonowania. Pakiet ten posłużył również do wyodrębnienia unikalnych sekwencji biologicznego pochodzenia, czyli tzw. sekwencji ASV (ang. *amplicon sequence variant*). Analiza składała się z etapów:

1. kontrola jakości odczytów:
    - analiza profilu błędu poszczególnych próbek oraz dynamiczne generowanie parametrów do kontroli jakości (narzędzie FIGARO)

- przeprowadzenie kontroli jakości w oparciu o współczynnik maksymalnych oczekiwanych błędów próbki (ang. maximum expected errors)

2. wstępna obróbka danych przy użyciu narzędzia Cutadapt: usunięcie sekwencji adaptorowych i odrzucenie zbyt krótkich odczytów (minimalna długość 30 nt.)
3. wyselekcjonowanie unikalnych sekwencji ASV (przy użyciu pakietu DADA2) poprzez: odfiltrowanie sekwencji zawierających błędy powstałe w trakcie procesu sekwencjonowania (ang. denoising), łączenie sparowanych odczytów (połączenie odpowiadających sobie odczytów forward i reverse), dereplikację (czyli scalenie identycznych, unikalnych sekwencji przy zachowaniu liczby ich wystąpień oraz profilu jakości) i filtrowanie chimer (pozbycie się konstruktów powstałych z nieprawidłowego złożenia sekwencji w trakcie PCR)
4. przyporządkowanie taksonomii do sekwencji ASV na podstawie referencyjnej bazy SILVA ver. 138 (Quast et al., 2013) , stosując podejście hybrydowe:
   - porównanie sekwencji ASV względem bazy w poszukiwaniu identycznych sekwencji referencyjnych (vsearch)
   - nietypowe sekwencje pozostałe po poprzednim kroku zostały zaklasyfikowane w oparciu o metody wykorzystujące uczenie maszynowe (sklearn).

Analizy społeczności mikroorganizmów wykonano wykorzystując pakiet phyloseq R ver. 1.38.0 (McMurdie et al. 2013), a uzyskane sekwencje zamieszczono w bazie NCBI jako Bioproject PRJNA1224491 dostępne pod adresem internetowym https://www.ncbi.nlm.nih.gov/bioproject?term=PRJNA1224491&cmd=DetailsSearch.

2.2.4.5. Analiza mikroskopowa liczebności i różnorodności pierwotniaków glebowych

Analizę liczebności i różnorodności orzęsek w glebie zaprezentowaną w artykule V przeprowadzono na Uniwersytecie Jagiellońskim w Krakowie.

Do badań wykorzystano próbki gleby o masie 10 g, które umieszczano w szalkach Petriego o średnicy 10 cm i nasycano 10 ml wody dejonizowanej. Tak przygotowane próbki inkubowano w komorze klimatycznej SANYO MLR-350 w temperaturze 20 ± 1°C w warunkach ciemności. Oznaczenia mikroskopowe wykonywano w 3, 6, 9 oraz 12 dniu inkubacji, stosując metodę Finlaya (Finay et al. 2000) z niewielkimi modyfikacjami. Wodę międzyporową pobierano poprzez delikatne uciskanie powierzchni gleby, a następnie 1 ml uzyskanego ekstraktu przenoszono do komory zliczającej Sedgewick Rafter S52 (Graticules Optics). Liczebność orzęsek oraz innych organizmów określano pod mikroskopem Nikon Eclipse E 80 przy powiększeniu 200×, analizując 200–400 µl wody glebowej na próbkę. Wyniki przeliczano na gram suchej gleby, uwzględniając całkowitą objętość wody glebowej w próbce. Różnorodność gatunkową oceniano na podstawie skumulowanych krzywych różnorodności gatunkowej w kolejnych dniach

inkubacji, a organizmy oznaczano do możliwie najniższego poziomu taksonomicznego przy użyciu kluczy Foissnera (Foissner & Berger 1996; Foissner et al.., 1991; 1992, 1999).

2.2.5. Analiza statystyczna

Wyniki przeprowadzonych analiz uzupełniono o analizę statystyczną, którą prowadzono z wykorzystaniem programu Statistica ver. 14.0 firmy StatSoft Inc. (obecnie TIBCO Software Inc.). W opracowaniu wyników uwzględniono wartości odchylenia standardowego (SD), względnego odchylenia standardowego (RSD, %) oraz współczynnika korelacji Pearsona. Uzyskane dane poddano następnie analizie rozkładu normalnego i jednoczynnikowej analizie wariancji (ANOVA). W przypadku uzyskania wyników istotnych statystycznie wykonano test post hoc Tukey'a do porównania poszczególnych grup. We wszystkich analizach za poziom istotności statystycznej przyjęto wartość $p < 0,05$. (Steliga et al., 2012, Wojtowicz et al., 2022; 2023a; 2023b; 2026b).

3. Wpływ γ-PGA na efektywność biodegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego wspomaganej bioaugmentacją- Artykuł I

Biodegradacja z zastosowaniem biopreparatów opartych na mikroorganizmach autochtonicznych jest jedną z podstawowych i powszechnie stosowanych metod oczyszczania gleb historycznie zanieczyszczonych węglowodorami ropopochodnymi, która w warunkach odpowiedniej biostymulacji oraz kontroli parametrów środowiskowych pozwala na znaczną redukcję stężeń zanieczyszczeń. W przypadku gleb z terenów dołów urobkowych, w celu uzyskania zakładanego poziomu oczyszczania, postanowiono zmodyfikować proces inokulacji poprzez wprowadzenie do gleby biopreparatu w połączeniu z γ-PGA, co przyczyniło się do zwiększenia biodostępności zanieczyszczeń i poprawy efektywności biodegradacji.

W artykule I przedstawiono kompleksowe badania bioremediacji gruntu z terenu dołu urobkowego G11 o wysokim stopniu skażenia węglowodorami ropopochodnymi (TPH=19 774,23 mg/kg s.m., BTEX=17,35 mg/kg s.m., WWA=27,03 mg/kg s.m.), obejmujące analizy mikrobiologiczne, respirometryczne oraz chromatograficzne gleby pod kątem zawartości węglowodorów alifatycznych, węglowodorów monoaromatycznych i wielopierścieniowych węglowodorów aromatycznych, a także analizy toksykologiczne z wykorzystaniem pakietu testów toksykologicznych Phytotoxkit™, Ostracodtoxkit F™, Microtox® SPT i Ames.

Badania zaprezentowane w artykule I prowadzono warunkach laboratoryjnych oraz w warunkach półtechnicznych (metoda pryzmowania *ex-situ*), co zwiększało ich znaczenie aplikacyjne oraz możliwość wykorzystania uzyskanych wyników w przyszłości w praktyce terenowej. Do badań biodegradacji w warunkach laboratoryjnych wykorzystano autorski biopreparat A1, opracowany na bazie szczepów bakterii wyizolowanych z gleby przeznaczonej do oczyszczania. Izolacja mikroorganizmów z badanego środowiska stanowiła podstawę konstrukcji biopreparatu i miała na celu zapewnienie jego wysokiej adaptacyjności oraz aktywności w warunkach silnego skażenia. Inokulacja biopreparatem A1 została zmodyfikowana poprzez dodatek γ-PGA, czyli kwasu γ-poliglutaminowego, co stanowiło element zastosowanej w badaniach modyfikacji technologicznej. Modyfikacja procesu inokulacji biopreparatem A1 poprzez dodatek γ-PGA miała charakter nowatorski w kontekście bioaugmentacji gleb zanieczyszczonych węglowodorami ropopochodnymi i nie była wcześniej opisywana w literaturze przedmiotu. Dobór γ-PGA jako czynnika modyfikacyjnego wynikał z jego potencjalnych właściwości biosurfaktantowych, co miało za zadanie wpłynąć na poprawę biodostępności zanieczyszczeń i w konsekwencji zwiększyć efektywność biodegradacji. Ponadto γ-PGA poprzez silne właściwości higroskopijne, umożliwił utrzymanie dobrego nawilżenia gleby i ograniczenie możliwości jej przesuszenia w trakcie oczyszczania.


Na podstawie przeprowadzonych analiz respirometrycznych możliwe było określenie stężenia γ-PGA, które w najlepszy sposób wpływało na wzrost aktywności mikrobiologicznej w układzie reakcyjnym, i wskazanie go do dalszych badań. Badania porównawcze prowadzone na rzeczywistych próbkach gleby z terenu dołu urobkowego

poddanych inokulacji biopreparatem A1 oraz zmodyfikowanej inokulacji biopreparatem A1 poprzez dodatek γ-PGA obejmujące monitoring stężeń zanieczyszczeń ropopochodnych oraz toksyczność gleby w trakcie oczyszczania, pozwolił na ocenę efektywności zastosowanych strategii bioremediacji oraz wpływu zaproponowanej modyfikacji na poprawę skuteczności biodegradacji zanieczyszczeń. Wyniki uzyskane w artykule I stanowiły podstawę do dalszych badań ukierunkowanych na zwiększenie efektywności biodegradacji węglowodorów ropopochodnych w glebach z dołów urobkowych, poprzez modyfikację procesów inokulacyjnych opracowanych na bazie bakterii autochtonicznych innymi substancjami wspomagającymi (Artykuł II).

3.1. Omówienie wyników zaprezentowanych w artykule I

Oceny efektywności zaproponowanych w artykule I strategii bioremedicji gleby zanieczyszczonej węglowodorami ropopochodnymi dokonano na podstawie przeprowadzonych analiz respirometrycznych, chromatograficznych i toksykologicznych.


Analizy respirometryczne prowadzone z wykorzystaniem systemu OxiTop, umożliwiły precyzyjny pomiar zużycia tlenu w trakcie eksperymentu, co było skorelowane ze wzrostem aktywności mikrobiologicznej w środowisku reakcyjnym (Rys. 15). Wyniki analiz respirometrycznych wykazały, że w glebie z terenu dołu urobkowego aktywność mikrobiologiczna była ograniczona, o czym świadczyła wartość zużycia tlenu po zakończeniu eksperymentu wynosząca zaledwie 62,1 mg O_2/dm^3. Zasadne zatem było zastosowanie technik wspomagających biodegradację. W wyniku inokulacji gleby biopreparatem A1 zużycie tlenu w układzie reakcyjnym po 60 dniach testu wzrosło do 1 582 mg O_2/dm^3, co potwierdziło skuteczność bioaugmentacji biopreparatem na bazie bakterii autochtonicznych. Jeszcze większe wartości zużycia tlenu odnotowano w układach inokulowanych biopreparatem A1 w połączeniu z γ-PGA. W zależności od stosunku objętościowego biopreparatu A1 do γ-PGA w procesie inokulacyjnym wartość zużycia tlenu w układzie po 60 dniach testu wynosiła 2 040 mg O_2/dm^3 (2:1) i 3 769 mg O_2/dm^3 (1:1). Analizy respirometryczne potwierdziły skuteczność zaproponowanych technik wspomagania biodegradacji, wskazując jednocześnie, że modyfikacja procesu inokulacji biopreparatem opracowanym na bazie bakterii autochtonicznych poprzez zastosowanie γ-PGA może istotnie wpłynąć na wzrost efektywności biodegradacji zanieczyszczeń ropopochodnych. Poza tym analizy respirometryczne wskazały, że efektywność biodegradacji zanieczyszczeń zależała od stosunku objętościowego biopreparatu A1 do γ-PGA w procesie inokulującym. W naszych badaniach wykazano, że najbardziej optymalny był stosunek 1:1, dla którego przeprowadzono dalsze badania w skali półtechnicznej (metoda pryzmowania *ex-situ*) obejmujące analizy chromatograficzne i toksykologiczne.



Rysunek 15 Zależność ilości zużytego tlenu [mg/dm^3] od czasu w glebie AB inokulowanej biopreparatem A1 oraz biopreparatem A1 i roztworem γ-PGA.

Przeprowadzone analizy chromatograficzne umożliwiły zidentyfikowanie i określenie zawartości poszczególnych zanieczyszczeń ropopochodnych w badanej glebie przed, w trakcie oraz po zakończeniu eksperymentu. Badania porównawcze prowadzone metodą *ex-situ* na glebie poddanej procesowi inokulacji biopreparatem A1 lub biopreparatem A1 i γ-PGA pozwoliły na określenie w jaki sposób zaproponowana modyfikacja techniki inokulacji wpłynęła na obniżenie zawartości oznaczonych węglowodorów ropopochodnych w badanej w glebie.

Zaprezentowane w artykule I wyniki analiz chromatograficznych wykazały wyraźne obniżenie zawartości węglowodorów ropopochodnych w glebie w wyniku zaproponowanych strategii oczyszczania. Inokulacja gleby biopreparatem A1 spowodowała obniżenie stężenia TPH o 66,03%, BTEX o 80,08% i WWA o 38,86% (po 6 miesiącach badań). Modyfikacja inokulacji biopreparatu A1 poprzez wprowadzenie do gleby dodatkowo γ-PGA wpłynęła na istotną poprawę efektywności biodegradacji tych węglowodorów ropopochodnych wynoszącej odpowiednio 79,21%; 90,19% i 51,18%. Wyniki te wskazują na skuteczność zaproponowanego sposobu modyfikacji procesu inokulacji biopreparatem A1, wyrażonej poprawą efektywności biodegradacji TPH o 13,18%, BTEX o 10,11% i WWA o 12,32%. Ponadto wyniki analiz chromatograficznych dostarczyły cennych informacji na temat podatności poszczególnych węglowodorów ropopochodnych wchodzących w skład tych zanieczyszczeń na biodegradację w wyniku zastosowania analizowanych technik wspomagających (Rys. 16).



Rysunek 16 Zestawienie stężeń n-alkanów oznaczonych w glebie z dołu urobkowego poddanej inokulacji a) biopreparatem A1, b) biopreparatem A1 oraz γ-PGA w trakcie prowadzenia procesu biodegradacji (metoda ex-situ).

Przeprowadzone w ramach badań analizy chromatograficzne wykazały, że niezależnie od sposoby inokulacji największą podatność na rozkład biologiczny w glebie AB wykazują węglowodory alifatyczne o długości łańcucha węglowego n-C_9 – n-C_{21}. Węglowodory z zakresu n-C_{22} – n-C_{30} również ulegały biodegradacji w zadowalającym stopniu, natomiast węglowodory alifatyczne posiadające powyżej 30 atomów węgla w cząsteczce były najbardziej oporne na rozkład biologiczny. Uzyskane wyniki były tożsame z wynikami innych badaczy i potwierdziły, że skuteczność biodegradacji węglowodorów alifatycznych maleje wraz ze wzrostem długości łańcuch węglowego w cząsteczce węglowodoru. Warto zauważyć, że skuteczność biodegradacji wszystkich oznaczonych grup węglowodorów alifatycznych była wyższa w glebie poddanej inokulacji biopreparatem A1 oraz γ-PGA aniżeli samym biopreparatem, wskazując na przewagę strategii modyfikowanej (Tabela 5). Ponadto identyfikacja węglowodorów z grupy izoprenoidów (Pr i F), należących do związków trudno biodegradowalnych, umożliwiła ocenę skuteczności bioremediacji na podstawie zmian wartości wskaźników n-C_{17}/Pr i n-C_{18}/F. Po 6 miesiącach badań wartości współczynników n-C_{17}/Pr oraz n-C_{18}/F zmniejszyły się z 3,0 do 0,48 i 0,53 w wariancie inokulowanym biopreparatem A1 oraz do 0,20 i 0,21 w wariancie inokulacji biopreparatem A1 z dodatkiem γ-PGA, co wskazuje na wysoką skuteczność zastosowanych strategii bioremediacyjnych.


Tabela 5 Porównanie efektywności biodegradacji wybranych grup węglowodorów ropopochodnych w glebie inokulowanej biopreparatem A1 oraz biopreparatem A1 i γ-PGA

Węglowodory	Efektywność biodegradacji w glebie inokulowanej biopreparatem A1 [%]	Efektywność biodegradacji w glebie inokulowanej biopreparatem A1 i γ-PGA [%]
TPH	66,03	79,21
Węglowodory niezidentyfikowane	59,49	71,84
$\sum nC_9–nC_{21}$	86,52 - 94,59	96,57 - 99,55
$\sum nC_{22}–nC_{30}$	44,62 - 77,77	80,35 - 88,35
$\sum nC_{31}–nC_{40}$	25,13 - 40,60	46,43 - 65,73
BTEX	80,08	90,19
benzen	91,75	95,89
toluen	95,54	98,89
etylobenzen	93,15	97,69
ksleny	71,06 – 78,69	85,14 – 89,88
∑WWAs	38,86	51,18
WWA dwupierścieniowe	71,58	81,06
WWA trzypierścieniowe	63,73 – 64,44	68,63 – 72,22
WWA czteropierścieniowe	32,33 – 42,78	44,65 – 56,24
WWA pięciopierścieniowe	30,06	40,30 – 43,17
WWA sześciopierścieniowe	24,62	35,48 – 37,83


Badania zawartości węglowodorów monoaromatycznych przeprowadzone z wykorzystaniem autorskiej metodyki chromatograficznego oznaczania węglowodorów z grupy BTEX umożliwiły identyfikację i ilościowe oznaczenie benzenu, toluenu, etylobenzenu i izomerów ksylenu w oczyszczanej glebie (Rys. 17). Analizy chromatograficzne wykazały, że w wyniku inokulacji gleby zarówno samym biopreparatem A1, jak i biopreparatem A1 oraz γ-PGA podatność na biodegradację malała w kolejności : toluen > etylobenzen > benzen > p-ksylen > m-ksylen > o-ksylen, natomiast stopnie ich biodegradacji były większe w glebie poddanej zmodyfikowanej bioaugmentacji (Tabela 5). Wskazuje to, że wprowadzenie do układu reakcyjnego γ-PGA wpłynęło na poprawę efektywności biodegradacji nie tylko węglowodorów alifatycznych, ale również węglowodorów monoaromatycznych. Może to świadczyć o wpływie γ-PGA na poprawę biodostępności zanieczyszczeń dla mikroorganizmów glebowych.

Analizy chromatograficzne gleby pod kątem zawartości WWA, wykazały zależność pomiędzy ilością pierścieni aromatycznych w cząsteczce węglowodoru, a jego podatnością na rozkład mikrobiologiczny. Oznaczone zawartości 15 zidentyfikowanych WWA w glebie przed, w trakcie i po zakończeniu oczyszczania wskazują, że zarówno w wyniku inokulacji gleby biopreparatem A1 jak i inokulacji biopreparatem A1 zmodyfikowanej poprzez dodatek γ-PGA podatność na biodegradację maleje w kolejności: WWA dwupierścieniowe > WWA trójpierścieniowe > WWA czteropierścieniowe > WWA pięciopierścieniowe > WWA sześciopierścieniowe (Rys. 18). Ponadto analiza WWA potwierdziła skuteczność γ-PGA pod kątem intensyfikacji procesów biodegradacyjnych, wyrażonej wyższymi wartościami stopni biodegradacji wszystkich zidentyfikowanych wielopierścieniowych węglowodorów aromatycznych w

glebie inokulowanej biopreparatem A1 wzbogaconej w γ-PGA aniżeli samym biopreparatem (Tabala 5). Wskazuje to na skuteczność zastosowanej techniki modyfikacji procesu inokulacji, pod kątem poprawy efektywności biodegradacji różnych grup węglowodorów ropopochodnych obecnych w zanieczyszczonej glebie.



Rysunek 17 Zestawienie stężeń węglowodorów z grupy BTEX w glebie z dołu urobkowego poddanej inokulacji a) biopreparatem A1, b) biopreparatem A1 oraz γ-PGA w trakcie prowadzenia procesu biodegradacji (metoda ex-situ).



Rysunek 18 Zestawienie stężeń WWA oznaczonych w glebie z dołu urobkowego poddanej inokulacji a) biopreparatem A1, b) biopreparatem A1 oraz γ-PGA w trakcie prowadzenia procesu biodegradacji (metoda ex-situ).

Uzupełnieniem analiz chromatograficznych przedstawionych w artykule I było obliczenie stałych szybkości biodegradacji zanieczyszczeń ropopochodnych w glebie oczyszczanej techniką inokulacji biopreparatem A1 oraz po jej zmodyfokowaniu poprzez dodatek preparatu γ-PGA. Wykazano, że wartości stałych szybkości biodegradacji były wyższe w układzie z dodatkiem γ-PGA i wynosiły odpowiednio 0,0083 d^{-1} dla TPH, 0,0112 d^{-1} dla BTEX oraz 0,0046 d^{-1} dla WWA, natomiast w układzie bez γ-PGA wynosiły 0,0058 d^{-1} dla TPH, 0,0072 d^{-1} dla BTEX i 0,0037 d^{-1} dla WWA, co potwierdziło korzystny wpływ dodatku γ-PGA na kinetykę biodegradacji zarówno węglowodorów alifatycznych, jak i aromatycznych.

Ocenę skuteczności zastosowanych strategii bioremediacyjnych uzupełniono o monitoring toksykologiczny którego celem było określenie wpływu zanieczyszczeń ropopochodnych (TPH, BTEX, WWA) oraz pośrednich metabolitów powstających w wyniku rozkładu mikrobiologicznego na biocenozę glebową w oparciu o przeprowadzone testy toksykologiczne. W artykule zaprezentowano wyniki testów Phytotoxkit^{TM}, Ostracodtoxikit F^{TM}, Microtox SPT oraz Ames, co umożliwiło kompleksową analizę toksyczności całego układu, z uwzględnieniem interakcji pomiędzy wszystkimi składnikami próbki. Wyniki testów toksykologicznych jednoznacznie wykazały, że zarówno inokulacja samym biopreparatem jak i z dodatkiem γ-PGA istotnie przyczyniły się do poprawy jakości gleby, jednocześnie potwierdzając przewagę strategii zmodyfikowanej inokulacji poprzez dodatek γ-PGA (Tabela 6). Ponadto wyniki analiz toksykologicznych dowiodły, że modyfikacja inokulacji poprzez dodatek preparatu γ-PGA jest bezpieczna i nie wpływa negatywnie na organizmy żywe.

Tabela 6 Wyniki testów toksykologicznych przeprowadzonych na próbce wyjściowej oraz próbkach zremediowanych techniką inokulacji biopreparatem A1 i biopreparatem A1 i γ-PGA

Testowany organizm	**Parametr**	**Gleba wyjściowa AB**	**Gleba inokulowana biopreparatem A1**	**Gleba inokulowana biopreparatem A1 i γ-PGA**
Sorghum saccharatum	kiełkowanie [%]	80	100	100
	średnia długość korzenia [mm]	11,4	40,8	44,8
	zahamowanie wzrostu [%]	76,25	15,00	6,67
Lepidium Sativum	kiełkowanie [%]	60	100	100
	średnia długość korzenia [mm]	11,6	56,4	62,9
	zahamowanie wzrostu [%]	83,19	18,26	8,84
Sinapis alba	kiełkowanie [%]	70	100	100
	średnia długość korzenia [mm]	20,2	68,4	75,5
	zahamowanie wzrostu [%]	74,75	14,5	5,63
Heterocypris incongruens	średnia śmiertelność [%]	81,67	18,33	10,00
	średnia hamowania wzrostu [%]	70,6	10,11	4,99
Vibrio fischeri	*TU*	32,3	9,4	4,3
Salmonella typhimirium TA-100	*Wskaźnik mutagenności*	6,15	2,15	1,54

3.2 Podsumowanie artykułu I

W artykule I przedstawiono rezultaty badań dotyczących bioremediacji gleby z dołu urobkowego z zastosowaniem inokulacji biopreparatem A1 opracowanym na bazie niepatogennych szczepów bakterii autochtonicznych oraz zmodyfikowanej inokulacji biopreparatem A1 poprzez dodatek γ-PGA. Analizy prowadzono w warunkach laboratoryjnych i półtechnicznych, co umożliwiło kompleksową ocenę wpływu zastosowanych strategii bioremediacyjnych na przebieg biodegradacji węglowodorów ropopochodnych oraz stan biocenozy glebowej. Na tej podstawie sformułowano najważniejsze wnioski wynikające z badań zaprezentowanych w artykule I:

- wykonane analizy respirometryczne wykazały, że modyfikacja inokulacji biopreparatem A1 poprzez dodatek roztworu γ-PGA spowodowała wzrost aktywności biologicznej w układzie reakcyjnym, co wskazywało na poprawę efektywności biodegradacji zanieczyszczeń ropopochodnych w badanej glebie,
- spośród testowanych wariantów inokulacji największym potencjałem pod kątem wspomagania procesów biodegradacji w glebie z dołu urobkowego wykazał wariant o optymalnym stosunku objętościowym biopreparatu A1 do γ-PGA wynoszącym 1:1, który wytypowano do badań porównawczych w skali półtechnicznej,
- przeprowadzenie badań w warunkach półtechnicznych na samodzielnie skonstruowanych stanowiskach badawczych pozwoliło na analizę zmian stężeń węglowodorów alifatycznych, BTEX oraz WWA w glebie inokulowanej biopreparatem A1 oraz biopreparatem A1 w połączeniu z γ-PGA, a także na porównanie wpływu zaproponowanej modyfikacji zabiegu inokulacji na zwiększenie efektywności biodegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego,
- biodegradacja zanieczyszczeń ropopochodnych bardziej intensywnie zachodziła w glebie inokulowanej biopreparatem i roztworem γ-PGA aniżeli samym biopreparatem. Stopień biodegradacji TPH, BTEX i WWA w glebie inokulowanej biopreparatem A1 w połączeniu z γ-PGA po zakończeniu oczyszczania wynosił odpowiednio 79,21%, 90,19% i 52,18%, natomiast w glebie inokulowanej samym biopreparatem odpowiednio 66,03%, 80,08% i 38,86%,
- wykonanie analiz chromatograficznych z wykorzystaniem autorskich procedur badawczych, umożliwiło jakościowe i ilościowe oznaczenie poszczególnych węglowodorów wchodzących w skład zanieczyszczeń ropopochodnych w trakcie prowadzenia procesu oczyszczania badanej gleby. Pozwoliło to na określenie podatności na biodegradację poszczególnych grup węglowodorów, określenie współczynników biodegradacji n-C_{17}/Pr i n-C_{18}/F będących miarą efektywności zaproponowanych strategii oczyszczania oraz wyznaczenie stałych szybkości biodegradacji,
- rozszerzenie badań o monitoring toksykologiczny w trakcie procesu oczyszczania pozwoliło na ocenę wpływu zastosowanych strategii bioremediacji na biocenozę

glebową oraz potwierdzenie bezpieczeństwa stosowania zaproponowanej modyfikacji zabiegu inokulacji.

Wyniki badań zaprezentowanych w artykule I wskazały, że modyfikacja inokulacji biopreparatem A1 opracowanym na bazie bakterii autochtonicznych poprzez wzbogacenie gleby roztworem γ-PGA wpłynęła na poprawę efektywności biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego. Pozytywne wyniki badań biodegradacji zanieczyszczeń w połączeniu z kompleksową oceną ekotoksykologiczną wskazują, że zaproponowana strategia bioremediacyjna może przyczynić się do głębszego doczyszczenia gruntów na obszarach dołów urobkowych i zostać w przyszłości wykorzystana w praktyce terenowej. Wyniki badań zaprezentowane w artykule I przyczyniły się do realizacji założonych celów pracy doktorskiej oraz stanowiły podstawę do dalszego rozwijania strategii bioremediacyjnej polegającej na modyfikacji procesów inokulacyjnych z zastosowaniem biopreparatów na bazie bakterii autochtonicznych, co zostało podjęte i rozwinięte w kolejnych publikacjach.

4. Rola biosurfaktantów w poprawie efektywności biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego – Artykuł II


W artykule II rozwinięto koncepcję możliwości modyfikacji inokulacji biopreparatem opracowanego na bazie bakterii autochtonicznych w poprawie efektywności bioremediacji gleby z dołu urobkowego G6, charakteryzującej się podwyższonym stężeniem węglowodorów ropopochodnych (TPH=6 062,59 mg/kg s.m., WWA=12,05 mg/kg s.m.). Zaproponowano możliwość modyfikacji zabiegu inokulacji biopreparatem poprzez dodatkowe zastosowanie biosurfaktantów, które bazując na dostępnych danych literaturowych, jako naturalne związki powierzchniowo czynne, mogą przyczynić się do obniżenia napięcia międzyfazowego, zwiększenia rozpuszczalności węglowodorów oraz ułatwienia desorpcji zanieczyszczeń z matrycy glebowej. Właściwości te wskazują, że obecność biosurfaktantów w glebie może znacząco poprawić biodostępność zanieczyszczeń ropopochodnych dla mikroorganizmów odpowiedzialnych za ich biodegradację, a tym samym zwiększyć efektywność procesu oczyszczania.

W artykule II przedstawiono kompleksowe badania bioremediacji gleby z dołu urobkowego G6 poddanej w zależności od układu eksperymentalnego bioaugmentacji biopreparatem na bazie bakterii autochtonicznych MC, zmodyfikowanej bioaugmentacji biopreparatem MC poprzez dodatek wybranego biosurfaktanta (autorska mieszanina PSR, preparaty JBR 425, JBR 320, ramnolipidy, surfaktyna lub γ-PGA) w dwóch stężeniach biosurfaktanta w mieszaninie inokulującej: 1g/dm^3 i 5 g/dm^3 lub wzbogaceniu wybranymi biosurfaktantami w stężeniach 1g/dm^3 i 5 g/dm^3 (próbki kontrolne). Badania prowadzone w warunkach laboratoryjnych umożliwiły jednoczesne zbadanie kilku wariantów inokulacji i określenie ich skuteczności. Weryfikacji efektywności zaproponowanych strategii bioremediacji dokonano na podstawie analiz respirometrycznych, chromatograficznych i toksykologicznych z wykorzystaniem testów Phytotoxkit^{TM}, Ostracodtoxikit F^{TM} oraz Microtox SPT. Przeprowadzone badania umożliwiły określenie wpływu zaproponowanych modyfikacji procesu inokulacji na efektywność biodegradacji TPH, WWA, a także poszczególnych węglowodorów wchodzących w skład tych zanieczyszczeń, wytypowanie najbardziej skutecznego biosurfaktanta w poprawie skuteczności biologicznego rozkładu węglowodorów ropopochodnych oraz określenie wpływu stężenia biosurfaktanta w mieszaninie inokulującej na efektywność bioremediacji gleby. Ponadto przeprowadzona ocena ekotoksykologiczna gleby po zakończeniu eksperymentu umożliwiła określenie wpływu zastosowanych strategii bioremediacyjnych na biocenozę glebową oraz bezpieczeństwa ich stosowania.

4.1. Omówienie wyników zaprezentowanych w artykule II


Wyniki badań laboratoryjnych przedstawionych w artykule II wykazały zróżnicowaną efektywność biodegradacji oznaczonych w glebie węglowodorów ropopochodnych (TPH i WWA), w zależności od rodzaju i stężenia zastosowanego biosurfaktanta w mieszaninie inokulującej. Omówione w artykule wyniki analiz

respirometrycznych, wykazały niewielkie zużycie tlenu w próbce nieinokulowanej, które po 90 dniach testu wynosiło zaledwie 15,51 mg O_2/dm^3, co wskazywało na ograniczoną biodegradację samoistną zanieczyszczeń w badanej glebie z dołu urobkowego, podobnie jak w badaniach omówionych w artykule I. Zastosowanie techniki wspomagającej jaką była inokulacja biopreparatem MC spowodowała wyraźny wzrost zużycia tlenu w układzie reakcyjnym 1 944,8 mg O_2/dm^3, potwierdzając skuteczność biopreparatów na bazie bakterii autochtonicznych w poprawie efektywności biodegradacji węglowodorów ropopochodnych. W próbkach wzbogaconych wyłącznie biosurfaktantami zużycie tlenu po zakończeniu eksperymentu było niższe i w zależności od stężenia substancji powierzchniowo czynnej wynosiło od 1 052,2 do 1 446,2 mg O_2/dm^3 (przy stężeniu 1g/dm^3) oraz od 1 273,2 do 1 749,8 mg O_2/dm^3 (przy stężeniu 5g/dm^3). Zdecydowanie największy wzrost aktywności biologicznej wykazano w układach inokulowanych biopreparatem MC w połączeniu z wytypowanymi do badań biosurfaktantami. W zależności od stężenia substancji powierzchniowo czynnej w mieszaninie inokulującej wartości zużycia tlenu po zakończeniu eksperymentu w testowanych układach reakcyjnych kształtowały się w granicach od 3 332,2 do 3 885,3 mg O_2/dm^3 (przy stężeniu 1g/dm^3) oraz od 4 258,3 mg O_2/dm^3 do 5 164,8 mg O_2/dm^3 (przy stężeniu 5g/dm^3) (Rys. 19). Ponadto warto zauważyć, że spośród wszystkich zastosowanych w badaniach biosurfaktantów najlepsze wyniki eksperymentów uzyskano dla inokulacji zmodyfikwanej poprzez dodatek mieszaniny PSR opracowanej na bazie różnych typów biosurfaktantów. Nieznacznie gorsze wyniki uzyskano dla komercyjnie dostępnych preparatów JBR 425 i JBR 320, natomiast najmniej skuteczna pod kątem wspomagania biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego G6 okazała się surfaktyna.



Rysunek 19 Zależność ilości zużytego tlenu [mg/dm^3] od czasu w glebie G6 inokulowanej biopreparatem MC, biosurfaktantami oraz biopreparatem z biosurfaktantami przy stężeniu biosurfaktantów w mieszaninie inokulującej a) 1 g/ dm^3, b) 5 g dm^3 (liczba powtórzeń n = 5, p < 0,05).


Przeprowadzone w ramach badań analizy chromatograficzne potwierdziły, że modyfikacja inokulacji biopreparatem MC poprzez wprowadzenie do układu biosurfaktantów, może istotnie wpłynąć na poprawę efektywności biodegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego. Stopnie biodegradacji TPH i WWA w próbce nieinokulowanej wynosiły odpowiednio 0,2% i 0,05%, co potwierdziło wyniki uzyskane w trakcie analiz respirometrycznych, jednoznacznie wykazując, że w glebie z dołu urobkowego biodegradacja samoistna praktycznie nie zachodziła. W związku z powyższym w celu bioremediacji gleby G6 konieczne było zastosowanie technik wspomagających. Inokulacja gleby biopreparatem MC spowodowała istotny wzrost efektywności biodegradacji TPH (44,33%) i WWA (36,97%) w układzie, natomiast zastosowanie inokulacji biopreparatem MC zmodyfikowanej poprzez dodatek wybranych biosurfaktantów do układów reakcyjnych doprowadziło do uzyskania jeszcze lepszego efektu oczyszczania. Oznaczone wartości stopni biodegradacji TPH w tych glebach kształtowały się w przedziale od 53,52% do 65,90% (przy stężeniu 1g/dm^3) oraz od 60,64% do 77,30% (przy stężeniu 5g/dm^3), natomiast WWA w przedziale od 43,67% do 61,53% (przy stężeniu 1g/dm^3) oraz od 52,64% do 70,32% (przy stężeniu 5g/dm^3). Dla porównania biodegradacja TPH i WWA w próbkach inokulowanych wyłącznie biosurfaktantami nie przekraczała 20% (Rys. 20).




Rysunek 20 Zestawienie stopni biodegradacji TPH i WWA w badanych układach reakcyjnych przy stężeniu biosurfaktantów 5 g/dm³ (liczba powtórzeń n = 7-10, p < 0,05).

Szczegółowa analiza n-alkanów, potwierdziła że modyfikacja inokulacji biopreparatem MC poprzez dodatek wytypowanych dawek substancji powierzchniowo czynnych wpływa na poprawę efektywności biodegradacji wszystkich n-alkanów. W każdej z analizowanych próbek najłatwiej biodegradacji ulegały węglowodory o długości łańcucha węglowego n-C_{10}–n-C_{21} oraz n-C_6–n-C_9. Nieznacznie gorsze wyniki biodegradacji uzyskano dla węglowodorów z zakresu C_{22}–n-C_{30}, natomiast zdecydowanie najbardziej oporne na degradację były węglowodory ciężkie n-C_{31}–n-C_{36} (Rys. 21). Analogiczne wyniki uzyskano dla wielopierścieniowych węglowodorów aromatycznych oznaczonych w badanych próbkach gleby. Szczegółowa analiza chromatograficzna

wykazała, że modyfikacja inokulacji biopreparatu MC poprzez dodatek wybranych biosurfaktantów prowadzi do poprawy efektywności biodegradacji wszystkich WWA, jednakże najbardziej podatne na biodegradację były naftalen oraz węglowodory trójpierścieniowe. Węglowodory zawierające cztery i więcej pierścieni aromatycznych w cząsteczce trudniej ulegały biodegradacji, co jest zgodne z wynikami omówionymi w artykule I (Rys.22).



Rysunek 21 Zestawienie stopni biodegradacji n-alkanów w badanych układach reakcyjnych z uwzględnieniem długości łańcucha węglowego przy stężeniu biosurfaktantów a) 1 g/dm³, b) 5 g/dm³ (liczba powtórzeń n = 7-10, p < 0,05).




Rysunek 22 Zestawienie stopni biodegradacji WWA w badanych układach reakcyjnych z uwzględnieniem liczby pierścieni aromatycznych w cząstece przy stężeniu biosurfaktantów 5 g/dm³ (liczba powtórzeń n = 7-10, p < 0,05).

W celu weryfikacji skuteczności procesu biodegradacji zanieczyszczeń ropopochodnych w badanej glebie obliczono wartości współczynników biodegradacji n-C_{17}/Pr i n-C_{18}/F przed i po zakończeniu eksperymentu.


W wyniku zaproponowanych technik wspomagających biodegradację uzyskano obniżenie wartości współczynnika n-C_{17}/Pr z 2,716 do 1,247 (inokulacja biopreparatem MC), 0,604 – 0,959 (inokulacja biopreparatem wzbogacona o dodatek biosurfaktanta o stężeniu 1 g/dm^3), 0,164 – 0,688 (inokulacja biopreparatem wzbogacona o dodatek biosurfaktanta o stężeniu 5 g/dm^3), 2,108 – 2,286 (inokulacja wyłącznie biosurfaktantem). Wartości drugiego współczynnika biodegradacji C_{18}/F obniżyły się z 2,575 do 1,303 (inokulacja biopreparatem), 0,665 – 1,004 (inokulacja biopreparatem wzbogacona o dadatek biosurfaktanta o stężeniu 1 g/dm^3), 0,173 – 0,719 (inokulacja biopreparatem wzbogacona o dodatek biosurfaktanta o stężeniu 5 g/dm^3), 2,109 – 2,392 (inokulacja wyłącznie biosurfaktantem). Wyraźnie niższe wartości obliczonych współczynników biodegradacji uzyskane w wyniku zmodyfikowanej techniki inokulacji potwierdziły założenie, że zastosowanie związków powierzchniowo czynnych w procesie inokulacji biopreparatem pozwala na poprawę efektywności biodegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego.

Bezpieczeństwo stosowania zaproponowanych w artykule strategii bioremediacyjnych oraz ich wpływ na spadek toksyczności badanego gruntu zweryfikowano poprzez przeprowadzenie pakietu testów toksykologicznych (Phytotoxkit^{TM}, Ostracodtoxikit F^{TM} oraz Microtox SPT) gleby przed i po zakończeniu eksperymentu.

Ocena ekotoksykologiczna gleby potwierdziła skuteczność zaproponowanej strategii bioremediacji w poprawie jakości badanego gruntu, czego skutkiem były obniżone wartości współczynników toksyczności TU w każdym z przeprowadzonych testów toksykologicznych (Rys. 23, Rys. 24). Największy spadek wartości współczynników toksyczności odnotowano w próbkach gleby inokulowanych biopreparatem w połączeniu z biosurfaktantami zastosowanymi w stężeniu 5 g/dm³, dla których wartości TU po zakończeniu eksperymentu kształtowały się w przedziale 5,20–6,35 dla kiełkowania nasion oraz 5,01–6,69 dla zahamowanie wzrostu korzenia w teście Phytotoxkit™, 4,35–6,42 w teście Ostracodtoxkit F™ oraz 4,91–7,05 o w teście Microtox SPT. Wyniki te korelowały z wynikami analiz respirometrycznych i chromatograficznych zaprezentowanymi w artykule, potwierdzając zależność pomiędzy stopniem toksyczności gleby z dołu urobkowego, a stężeniem zawartych w niej zanieczyszczeń ropopochodnych. Ponadto obniżone wartości współczynników toksyczności po zakończeniu eksperymentu wskazują, że w wyniku zaproponowanej modyfikacji procesu inokulacji w oczyszczanej glebie nie powstawały toksyczne metabolity pośrednie, co jest niezwykle istotne z punktu widzenia bezpieczeństwa środowiskowego.



Rysunek 23 Wyniki testu fitotoksyczności Phytotoxkit™ przeprowadzonego po zakończeniu biodegradacji w badanych układach reakcyjnych przy stężeniu biosurfaktantów 5 g/L (liczba powtórzeń n = 7-10, p < 0,05).



Rysunek 24 Wyniki testów a) Ostracodtoxkit b) Microtox przeprowadzonych po zakończeniu biodegradacji w badanych układach reakcyjnych (liczba powtórzeń n = 7-10, p < 0,05).

4.2 Podsumowanie artykułu II

W artykule II przedstawiono wyniki badań efektywności biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego w wyniku inokulacji biopreparatem MC oraz zmodyfikowanego procesu inokulacji poprzez dodatkowe wprowadzenie do układu biosurfaktantów. Zaproponowana modyfikacja procesu

inokulacji miała na celu poprawę biodostępności zanieczyszczeń, czego konsekwencją było zwiększenie skuteczności biologicznego rozkładu węglowodorów alifatycznych i WWA. Badania prowadzone w warunkach laboratoryjnych umożliwiły jednoczesne przetestowanie kilku biosurfaktantów jako dodatków modyfikujących proces bioaugmentacji i wytypowanie tego, który charakteryzował się największym potencjałem do intensyfikacji bidegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego. Wyniki uzyskane na podstawie przeprowadzonych analiz respirometrycznych, chromatograficznych i toksykologicznych umożliwiły kompleksową ocenę zastosowanych strategii wspomagania bioremediacji oraz sformułowanie najważniejszych wniosków wynikających z badań zaprezentowanych w artykule:

- przeprowadzone analizy respirometryczne wykazały wzrost aktywności biologicznej w badanych układach reakcyjnych wzbogaconych różnymi mieszaninami inokulacyjnymi, przy czym skuteczność oddziaływania była zależna od ich składu. Najniższą aktywność odnotowano dla układów z dodatkiem samych biosurfaktantów, wyższą dla biopreparatu MC, natomiast najwyższą dla inokulacji biopreparatem MC oraz biosurfaktantami,
- uzyskane wyniki jednoznacznie potwierdziły, że modyfikacja inokulacji biopreparatem MC poprzez dodatek biosurfaktantów była najbardziej efektywna pod względem zwiększenia intensywności procesów biodegradacji węglowodorów ropopochodnych w badanej glebie;
- spośród analizowanych biosurfaktantów najwyższą skuteczność pod kątem wspomagania procesu bidegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego wykazała autorska mieszanina PSR, składająca się z trzech różnych typów biosurfaktantów. Zastosowanie pozostałych biosurfaktantów również prowadziło do istotnej poprawy efektywności bioremediacji gleby, przy czym ich skuteczność malała w kolejności: JBR 425 > JBR 320 > γ-PGA > ramnolipidy > surfaktyna,
- badania prowadzone przy dwóch stężeniach biosurfaktantów w mieszaninie inokulacyjnej wykazały, że efektywność oczyszczania gleby była uzależniona od dawki zastosowanego biosurfaktanta w układzie,
- wykonanie analiz chromatograficznych z wykorzystaniem autorskich procedur badawczych umożliwiło jakościowe i ilościowe oznaczenie poszczególnych węglowodorów ropopochodnych obecnych w zanieczyszczonej glebie, co pozwoliło na ocenę podatności na biodegradację poszczególnych frakcji węglowodorów oraz wyznaczenie wskaźników n-C_{17}/Pr i n-C_{18}/F, stanowiących miarę efektywności zastosowanych strategii remediacyjnych,
- modyfikacja inokulacji biopreparatem MC poprzez dodatek biosurfaktantów prowadziła do zwiększenia efektywności biodegradacji wszystkich oznaczonych w glebie węglowodorów ropopochodnych, co wskazuje na istotną poprawę biodostępności zanieczyszczeń dla mikroorganizmów,
- weryfikacja skuteczności zaproponowanej metody z wykorzystaniem testów toksykologicznych potwierdziła efektywność inokulacji wspomaganej biosurfaktantami w poprawie jakości gleby zanieczyszczonej TPH i WWA oraz

wykazała, że proces ten jest bezpieczny środowiskowo i nie prowadzi do powstawania wtórnych metabolitów o właściwościach toksycznych.

Wyniki badań laboratoryjnych zaprezentowanych w artykule II wykazały, że modyfikacja inokulacji biopreparatu na bazie bakterii autochtonicznych poprzez dodatek biosurfaktantow wpłynęła na poprawę efektywności bioremediacji gleby z dołu urobkowego. Kompleksowe wyniki obejmujące analizy respirometryczne, chromatograficzne i ocenę ekotoksykologiczna gleby po zakończeniu eksperymentu, wskazują że zaproponowana strategia bioremediacji będzie mogła zostać w przyszłości wdrożona w warunkach przemysłowych. Warto zaznaczyć, że opracowana strategia została już zweryfikowana w warunkach półtechnicznych, gdzie do inokulacji gleby z dołu urobkowego wykorzystano połączenie biopreparatu MC z autorską mieszaniną biosurfaktantów PSR. Wyniki tych badań zostały omówione w artykule „*Wpływ mieszaniny biosurfaktantów na efektywność bioremediacji gruntu zanieczyszczonego węglowodorami ropopochodnymi*" (Wojtowicz & Steliga, 2024), i potwierdziły skuteczność zaproponowanej strategii bioremediacyjnej w warunkach zbliżonych do rzeczywistych. Badania zaprezentowane w artykule II przyczyniły się do realizacji założonych celów niniejszej rozprawy doktorskiej, a także stanowiły podstawę do rozszerzenia zakresu prowadzonych prac o dalsze modyfikacje biopreparatów na bazie bakterii autochtonicznych. Ponadto bazując na obserwacjach uzyskanych w ramach realizacji badań omówionych w artykułach I-II, zdecydowano się opracować zintegrowane strategie bioremediacyjne łączące ze sobą bioaugmentację z fitoremediacją, co miało na celu głębsze doczyszczenie gleb z dołów urobkowych.

5. Wpływ wzbogacenia biopreparatu bakteryjnego w wybrane autochtoniczne szczepy grzybów na efektywność biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego oraz zastosowania fitoremediacji w celu jej głębszego doczyszczenia – Artykuły III i IV

W artykule III zaprezentowano możliwość zwiększenia efektywności biodegradacji węglowodorów ropopochodnych w glebie z dołu urobkowego poprzez wzbogacenie biopreparatu opracowanego na bazie bakterii autochtonicznych w wybrane szczepy grzybów autochtonicznych. Bazując na dostępnej literaturze przedmiotu wskazującej, że niektóre szczepy grzybów mają zdolność do degradacji węglowodorów ropopochodnych, założono, że rozszerzenie składu biopreparatu o wybrane szczepy grzybów wyizolowanych z gleby przeznaczonej do oczyszczania pozwoli na rozszerzenie spektrum działania biopreparatu oraz wspomoże biodegradację zarówno węglowodorów alifatycznych jak i WWA. W artykule zaprezentowano badania prowadzone w warunkach półtechnicznych na dwóch próbkach gleby z dołu urobkowego, które w zależności od układu eksperymentalnego inokulowano biopreparatem bakteryjnym (B1) lub biopreparatem bakteryjno-grzybowym (B2). Badania prowadzone w dwóch układach eksperymentalnych z wykorzystaniem analiz chromatograficznych i toksykologicznych, umożliwiły porównanie efektywności biodegradacji węglowodorów alifatycznych i WWA obecnych w zanieczyszczonej glebie w wyniku inokulacji różnymi biopreparatami.

Istotnym elementem badań zaprezentowanym w artykułach III i IV było wprowadzenie kolejnej techniki bioremediacyjnej jaką była fitoremediacja. Fitoremediacja polega na wykorzystaniu roślin i ich naturalnych procesów fizjologicznych oraz biochemicznych do usuwania, przekształcania lub unieszkodliwiania zanieczyszczeń obecnych w glebie. W przypadku historycznego skażenia gleby węglowodorami ropopochodnymi skuteczność fitoremediacji często bywa ograniczona ze względu na hydrofobowy charakter zanieczyszczeń i ich niską biodostępność. Połączenie w naszych badaniach fitoremediacji z bioaugmentacją pozwoliło na pełniejsze wykorzystanie potencjału tej techniki bioremediacyjnej i uzyskanie wyraźnie lepszych efektów oczyszczania gleb z terenów dołów urobkowych.

W artykułach III i IV badania fitoremediacji prowadzono z wykorzystaniem roślin uprawnych *Zea mays* (Artykuł III) i ozdobnych *Echinacea purpurea* (Artykuł IV). W artykule III przedstawiono wyniki badań fitoremediacji oraz fitoremediacji wspomaganej inokulacją biopreparatem B2 lub biopreparatem B2 oraz γ-PGA. Badania prowadzone w trzech układach badawczych oparte na wynikach analiz chromatograficznych i toksykologicznych gleb oraz analiz morfologicznych roślin fitoremediacyjnych, pozwoliły na porównanie efektywności zaproponowanych strategii bioremediacyjnych oraz potwierdziły słuszność łączenia kilku technik bioremediacyjnych. W artykule IV przedstawiono wyniki badań fitoremediacji z wykorzystaniem *Echinacea purpurea*,

prowadzonych na dwóch typach gleb, tj. glebie z dołu urobkowego oraz glebie z terenu naturalnego wycieku ropy naftowej. Badania prowadzone w kilku układach badawczych obejmowały fitoremediację oraz fitoremediację wspomaganą bioaugmentacją z zastosowaniem biopreparatów B1 i B2, a także ich wersji wzbogaconych w γ-PGA oraz wariant z glebą wzbogaconą wyłącznie w γ-PGA. Analogicznie do badań zaprezentowanych w artykule III, badania fitoremediacji w artykule IV również prowadzono w oparciu o wyniki analiz chromatograficznych i toksykologicznych gleb oraz analiz morfologicznych roślin uczestniczących w procesie fitoremediacji. Dodatkowo oznaczano zawartość węglowodorów (TPH i WWA) w korzeniach i pędach *Echinacea purpurea*, co pozwoliło na ocenę stopnia ich akumulacji w roślinie. Przedstawione w artykułach III i IV badania stanowią spójne rozwinięcie wcześniejszych prac i potwierdzają zasadność przyjętej strategii badawczej polegającej na wprowadzeniu fitoremediacji jako elementu wspomagającego procesy bioremediacji gleb z dołów urobkowych. Innowacyjność przeprowadzonych badań polegała na zastosowaniu połączenia fitoremediacji i bioaugmentacji, w tym z wykorzystaniem zmodyfikowanej strategii inokulacji biopreparatem oraz γ-PGA omówionej w artykule I, które w odniesieniu do rzeczywistych gleb z terenów dołów urobkowych nie było dotychczas szeroko stosowane.

5.1. Omówienie wyników badań biodegradacji zaprezentowanych w artykule III


Badania biodegradacji zaprezentowane w artykule III wykazały, że wzbogacenie biopreparatu opracowanego na bazie bakterii autochtonicznych w wybrane szczepy grzybów może wpłynąć na poprawę efektywności biodegradacji zarówno węglowodorów alifatycznych jak i wielopierścieniowych węglowodorów aromatycznych. Analizy chromatograficzne przeprowadzone na próbkach gleby poddanych inokulacji biopreparatem bakteryjnym (B1) lub biopreparatem bakteryjno grzybowym (B2) wykazały znaczącą różnicę w stopniach biodegradacji TPH i WWA w trakcie oczyszczania. W glebie inokulowanej biopreparatem B1 po trzech miesiącach bioremediacji uzyskano obniżenie stężenia TPH o 31,85% i WWA o 27,41%, natomiast w glebie zaszczepionej biopreparatem B2 wartości te wynosiły odpowiednio 41,67% (TPH) oraz 34,73% (WWA). Wynika z tego, że wzbogacenie biopreparatu w wybrane szczepy grzybów wpłynęło na poprawę efektywności biodegradacji TPH o 9,82% i WWA o 7,32%, co wskazuje na wysoki potencjał biopreparatów bakteryjno-grzybowych do zastosowania w bioremediacji gleb z terenów dołów urobkowych (Tabela 7).

Tabela 7 Porównanie efektywności biodegradacji wybranych grup węglowodorów ropopochodnych w glebie inokulowanej biopreparatem B1 i B2


Węglowodory	Efektywność biodegradacji w glebie inokulowanej biopreparatem B1 [%]	Efektywność biodegradacji w glebie inokulowanej biopreparatem B2 [%]
TPH	31,85	41,67
Węglowodory niezidentyfikowane	31,44	37,97
nC_6–nC_9	53,96 – 63,99	78,50 – 65,17
nC_{10}–nC_{21}	34,78 – 64,58	54,08 -78,00
nC_{22}–nC_{30}	10,25 – 32,44	22,05 – 47,63
nC_{31}–nC_{40}	9,23 – 10,67	14,77 – 18,65
∑WWAs	27,41	34,73
WWA dwupierścieniowe	34,49	43,81
WWA trzypierścieniowe	24,40 – 30,82	30,12 – 38,05
WWA czteropierścieniowe	15,22 – 21,71	18,79 – 26,80
WWA pięciopierścieniowe	11,42 – 11,46	16,56 – 16,61
WWA sześciopierścieniowe	7,96 – 8,37	14,77 – 15,27

Szczegółowa analiza n-alkanów potwierdziła korelacje pomiędzy długością łańcucha węglowego, a podatnością węglowodorów alifatycznych na biodegradację (Rys. 25). Zarówno w wyniku zaszczepienia gleby biopreparatem B1 jak i biopreparatem B2 podatność węglowodorów na biodegradację malała w następującej kolejności: nC_{10}–nC_{21} > nC_6–nC_9 > nC_{22}–nC_{30} > nC_{31}–nC_{36}, co jest zgodne z wcześniej omówionymi badaniami (artykuł I i II). Warto zauważyć, że obecność grzybów w biopreparacie wpłynęła na poprawę efektywności biodegradacji wszystkich węglowodorów alifatycznych, jednak największy wpływ miała na biodegradację węglowodorów o długości łańcucha węglowego n-C_{22}–n-C_{30} oraz węglowodorów ciężkich o długości łańcucha węglowego n-C_{31}–n-C_{36} (Tabela 7). Obserwacje te były zgodne z dostępnymi danymi literaturowymi, które wymieniają *Penicillium* sp., *Candida* sp., *Aspergillus* sp. jako dobrych degraderów węglowodorów alifatycznych o długości łańcucha węglowego C_{10}–C_{36}. O zwiększonej efektywności biodegradacji związków ropopochodnych po wzbogaceniu biopreparatu bakteryjnego wybranymi szczepami grzybów świadczyły również niższe wartości wskaźników n-C_{17}/Pr i n-C_{18}/F. Po zastosowaniu bioaugmentacji biopreparatem bakteryjno-grzybowego wskaźniki te obniżyły się odpowiednio do wartości 1,32 i 1,38, natomiast po bioaugmentacji biopreparatem bakteryjnym osiągnęły wartości 1,69 i 1,80.

Szczegółowa analiza WWA również wykazała poprawę efektywności biodegradacji wszystkich oznaczonych wielopierścieniowych węglowodorów aromatycznych w glebie, w wyniku inokulacji biopreparatem opracowanym na bazie autochtonicznych bakterii i grzybów (Tabela 7). Szybkość usuwania WWA w glebie zaszczepionej zarówno biopreparatem B1 jak i biopreparatem B2, była zgodna z wynikami uzyskanymi w pozostałych artykułach i zmniejszała się w następującej kolejności: 2-pierścieniowy WWA > 3-pierścieniowe WWA > 4-pierścieniowe WWA > 5-pierścieniowe WWA > 6-pierścieniowe WWA (Rys. 26).




Rysunek 25 Zestawienie stężeń n-alkanów oznaczonych w glebie z dołu urobkowego poddanej inokulacji a) biopreparatem B1, b) biopreparatem B2 trakcie prowadzenia procesu biodegradacji (metoda ex-situ).



Rysunek 26 Zestawienie stężeń WWA oznaczonych w glebie z dołu urobkowego poddanej inokulacji a) biopreparatem B1, b) biopreparatem B2 trakcie prowadzenia procesu biodegradacji (metoda ex-situ).

Efektywność oraz bezpieczeństwo stosowania zaprezentowanych w artykule III biopreparatów opracowanych na bazie autochtonicznych bakterii (B1) oraz bakterii i grzybów (B2) w bioremediacji gleb z dołów urobkowych zanieczyszczonych węglowodorami ropopochodnymi została sprawdzona poprzez przeprowadzenie testów toksykologicznych Phytotoxkit™, Ostracodtoxkit F™ oraz Microtox SPT. Zauważalny spadek toksyczności gleby po zakończeniu eksperymentu w każdym z parametrów zastosowanych testów toksykologicznych wskazuje na bezpieczeństwo stosowania opracowanych biopreparatów oraz zauważalną poprawę jakości gleby, w tym również w wariancie zawierającym grzyby, którego potencjalny wpływ na powstawanie produktów pośrednich nie był wcześniej weryfikowany.



Rysunek 27 Wyniki testów a) Phytotoxkit, b)Ostracodtoxkit, c) Microtox SPT przeprowadzonych na próbce wyjściowej oraz próbkach zremediowanych techniką inokulacji biopreparatem B1 i B2.


Uzyskane wyniki biodegradacji węglowodorów alifatycznych oraz wielopierścieniowych węglowodorów aromatycznych po inokulacji gleby biopreparatami B1 i B2 stanowiły podstawę do podjęcia badań nad ich zastosowaniem w fitoremediacji wspomaganej bioaugmentacją co omówiono w dalszej części pracy.

5.2. Omówienie wyników fitoremediacji wspomaganej bioaugmentacją zaprezentowanych w artykułach III i IV


W artykułach III i IV zaproponowano strategię poprawy efektywności biodegradacji węglowodorów ropopochodnych w glebie, polegającą na połączeniu fitoremediacji oraz bioaugmentacji. W badaniach wykorzystano rośliny uprawne *Zea mays* (artykuł III) oraz ozdobne *Echinacea purpurea* (artykuł IV), które zgodnie z dostępnymi danymi literaturowymi charakteryzują się stosunkowo wysoką tolerancją na obecność ksenobityków w glebie oraz szybkim tempem wzrostu i dużą produkcją biomasy.

Badania fitoremediacji z wykorzystaniem *Zea mays* zaprezentowane w artykule III były ukierunkowane na „doczyszczenie" skażonej gleby oraz wykazanie, że inokulacja odpowiednio opracowanym biopreparatem umożliwia znaczną poprawę efektywności fitoremediacji gleb zanieczyszczonych węglowodorami ropopochodnymi. Badania prowadzone w trzech układach badawczych przez okres sześciu miesięcy wykazały, że skuteczność oczyszczania gleby zależała od zastosowanej techniki wspomagania bioremediacji. W układzie, w którym zastosowano wyłącznie fitoremediację efektywność biodegradacji TPH po zakończeniu eksperymentu wynosiła zaledwie 22,81%. Połączenie fitoremediacji z inokulacją biopreparatem B2 wpłynęło na znaczną poprawę efektywność biodegradacji TPH (49,08%), natomiast dodatkowe wzbogacenie inokulacji biopreparatem B2 w γ-PGA skutkowało dalszym wzrostem efektywności biodegradacji TPH (65,35%). Analogiczne wyniki uzyskano dla WWA, mianowicie skuteczność biodegradacji WWA po sześciu miesiącach wynosiła odpowiednio 18,48% w glebie poddanej wyłącznie fitoremediacji, 40,74% w glebie poddanej fitoremediacji wspomaganej inokulacją biopreparatem B2 oraz 60,80% w glebie poddanej fitoremediacji oraz inokulacji biopreparatem B2 wzbogaconej w γ-PGA. Uzyskane wyniki jednoznacznie wskazują, że sama fitoremediacja nie jest skuteczną strategią oczyszczania gleb z dołów urobkowych natomiast połączenie jej z bioaugmentacją pozwala na istotną poprawę jej efektywności.

Szczegółowe analizy chromatograficzne gleby pod kątem zawartości węglowodorów alifatycznych i wielopierścieniowych węglowodorów aromatycznych (Rys. 28, Rys. 29), potwierdziły skuteczność zaproponowanych strategii łączących fitoremediację z bioaugmentacja, w celu doczyszczenia gruntu skażonego węglowodorami ropopochodnymi. Wszystkie oznaczone w glebie n-alkany oraz WWA najszybciej ulegały rozkładowi w glebie obsadzonej i inokulowanej biopreparatem B2 z dodatkiem γ-PGA, natomiast najwolniej w glebie wyłącznie obsadzonej. Szybkość usuwania poszczególnych węglowodorów była analogiczna do pozostałych badań zaprezentowanych w niniejszej rozprawie doktorskiej, mianowicie węglowodory alifatyczne ulegały degradacji w kolejności: nC_{10}–nC_{21} > nC_6–nC_9 > nC_{22}–nC_{30} > nC_{31}–nC_{36}, natomiast WWA w kolejności: 2-pierścieniowy WWA > 3-pierścieniowe WWA > 4-pierścieniowe WWA > 5-pierścieniowe WWA > 6-pierścieniowe WWA. Ponadto wyniki analiz chromatograficznych zaprezentowanych w artykule III potwierdziły, że łączenie kilku technik bioremediacyjnych (fitoremediacja, bioaugmentacja) oraz modyfikacje zabiegu inokulacji z wykorzystaniem biopreparatów na bazie niepatogennych autochtonicznych gatunków bakterii i grzybów poprzez wprowadzenie substancji wspomagających biodegradację takich jak np. γ-PGA (Artykuły I i II) pozwalają na uzyskanie największej skuteczności usuwania węglowodorów ropopochodnych z gleb historycznie zanieczyszczonych węglowodorami ropopochodnymi.




Rysunek 28 Zestawienie stężeń n-alkanów oznaczonych w glebie poddanej fitoremediacji z wykorzystaniem Zea mays



Rysunek 29 Zestawienie stężeń WWA oznaczonych w glebie poddanej fitoremediacji z wykorzystaniem Zea mays


Uzyskane wyniki analiz chromatograficznych korelowały z ogólnym stanem roślin przeprowadzających fitoremediację. Ocena morfologiczna roślin wykazała, że rozwój *Zea mays* zależał od sposobu wzbogacenia gleby. Rośliny pobrane z gleby nieinokulowanej charakteryzowały się niewielkimi rozmiarami od 30 do 50 cm oraz słabo rozwiniętymi systemami korzeniowymi. Rozwój roślin w glebie poddanej inokulacji biopreparatem B2 był zdecydowanie lepszy, natomiast nawiększy wzrost i rozwój systemów korzeniowych zaobserwowano dla *Zea mays* zasadzonych w glebie inokulowanej biopreparatem B2 oraz dodatkiem γ-PGA (Rys. 30). Połączenie analizy

morfologicznej z wynikami analiz chromatograficznych potwierdziło występowanie złożonych interakcji pomiędzy rośliną, a mikroorganizmami rozkładającymi węglowodory oraz zależności między obecnością zanieczyszczeń w glebie, a stanem roślin. Uzyskane wyniki wskazują, że mikroorganizmy mogą przekształcać zanieczyszczenia w formy łatwiej przyswajalne przez rośliny oraz obniżać ich stężenie do poziomu lepiej tolerowanego przez zastosowane w badaniach fitoremedianty. Z drugiej strony rośliny, poprzez prawidłowy wzrost oraz rozwój systemu korzeniowego, mogą poprawiać napowietrzenie gleby, wydzielać enzymy i związki organiczne wspomagające biodegradację zanieczyszczeń, poprawiać biodostępność zanieczyszczeń oraz stymulować aktywność i rozwój mikroorganizmów glebowych.



Rysunek 30 Zdjęcia Zea mays w glebie a) nieinokulowanej, b) inokulowanej biopreparatem B2, c) inokulowanej biopreparatem B2 i γ-PGA po 3 miesiącach uprawy

Ocena ekotoksykologiczna gleby po zakończeniu oczyszczenia techniką fitoremediacji oraz fitoremediacji wspomaganej bioaugmentacją biopreparatem B2 i biopreparatem B2 z dodatkiem γ-PGA, dowiodła, że wraz ze spadkiem stężenia zanieczyszczeń węglowodorów ropopochodnych w glebie maleje stopień jej toksyczności, co było zgodne z wynikami zaprezentowanymi w rozdziale 5.1. Ponadto sukcesywny spadek toksyczności gleby w każdym z parametrów testów Phytotoxkit™, OstracodtoxkitF™ oraz Microtox SPT wskazywał, że zaproponowana strategia oczyszczania gleb polegająca na połączeniu fitoremediacji z bioaugmentacją biopreparatem B2 i biopreparatem B2 z dodatkiem γ-PGA jest bezpieczna i pozwala na uzyskanie większego spadku toksyczności gleby niż w przypadku samej bioaugmentacji, przy jednoczesnym braku obserwacji efektów niepożądanych (Rys. 31).




Rysunek 31 Wyniki testów toksykologicznych przeprowadzonych na próbce gleby zanieczyszczonej węglowodorami ropopochodnymi poddanej fitoremediacji z wykorzystaniem Zea mays

Strategia bioremediacji gleby zanieczyszczonej węglowodorami ropopochodnymi z wykorzystaniem fitoremediacji wspomaganej bioaugmentacją została rozwinieta w artykule IV. Badania z wykorzystaniem *Echinacea purpurea* potwierdziły skuteczność łączenia kilku technik bioremediacyjnych, czyli fitoremediacji i inokulacji biopreparatami opracowanymi na bazie mikroorganizmów autochtonicznych oraz ich zmodyfikowanymi wersjami w poprawie efektywności biodegradacji zarówno węglowodorów alifatycznych jak i aromatycznych. Badania prowadzone z wykorzystaniem innej rośliny oraz dwóch rodzajów gleby (gleba z dołu urobkowego SD i wycieku ropy naftowej OS) wskazały że zaproponowana strategia bioremediacyjna jest uniwersalna i może zostać zastosowana w różnych warunkach uprawy.

Wyniki analiz chromatograficznych przeprowadzonych na próbkach gleb poddanych fitoremediacji z wykorzystaniem *Echinacea purpurea* jednoznacznie potwierdziły, że bez wprowadzenia do gleby mikroorganizmów wspomagających degradację węglowodorów ropopochodnych, efektywność oczyszczania gruntów z terenów dołów urobkowych i wycieków ropy naftowej jest niewielka. Po sześciu miesiącach badań stopnie biodegradacji TPH i WWA w glebie z dołu urobkowego poddanej wyłącznie fitoremediacji kształtowały się na poziomie 18,40% i 14,19%. Nieznacznie wyższe wartości stopni biodegradacji odnotowano w glebie obsadzonej i wzbogaconej w γ-PGA. Analogicznie do badań z wykorzystaniem *Zea mays,* inokulacja gleby z dołu urobkowego biopreparatami opracowanymi na bazie mikroorganizmów autochtonicznych prowadziła do wyraźnej poprawy efektywności fitoremediacji, przy czym lepsze efekty oczyszczania uzyskano dla biopreparatu bakteryjno grzybowego. Wartość stopnia biodegradacji TPH w glebie obsadzonej *Echinacea purpurea* i inokulowanej biopreparatem B1 po zakończeniu eksperymentu wynosiła 35,56%, natomiast obsadzonej i zaszczepionej biopreparatem B2 osiągnęła wartość 44,16%. Analogicznie, wartości stopni biodegradacji WWA wynosiły 30,82% w glebie obsadzonej i inokulowanej biopreparatem B1 oraz 36,73% w glebie obsadzonej i inokulowanej biopreparatem B2. Jeszcze wyższą efektywność biodegradacji zanieczyszczeń ropopochodnych odnotowano w układach inokulowanych biopreparatami oraz γ-PGA, co przełożyło się na spadek zawartości TPH i WWA w tych glebach o odpowiednio 46,71% i 39,22% (gleba obsadzona oraz inokulowana biopreparatem B1 i γ-


PGA) oraz 53,98% i 49,54% (gleba obsadzona oraz inokulowana biopreparatem B2 i γ-PGA) (Rys. 32). Badania przeprowadzone na glebie z terenu wycieku ropy naftowej wykazały podobne zależności pomiędzy sposobem fitoremediacji, a efektywnością biodegradacji węglowodorów ropopochodnych w glebie, co wskazuje na możliwość wykorzystania tej metody na różnych typach gleb zanieczyszczonych węglowodorami ropopochodnymi.



Rysunek 32Oznaczone zawartości a) TPH i b) WWA w glebie z dołu urobkowego poddanej fitoremediacji z wykorzystaniem Echinacea purpurea w różnych wariantach wzbogacenia gleby (system 1- brak inokulacji, system 2 – inokulacja biopreparatem B1, system 3 – inokulacja biopreparatem B1 i γ-PGA, system 4 – inokulacja biopreparatem B21, system 5 – inokulacja biopreparatem B2 i γ-PGA, system 6 – inokulacja γ-PGA).

W artykule IV ocenę efektywności fitoremediacji rozszerzono o analizę materiału roślinnego, obejmującą oznaczenie zawartości TPH i WWA w korzeniach i pędach *Echinacea purpurea* po zakończeniu oczyszczania. Bazując na nielicznych doniesieniach literaturowych wskazujących, że niewielka ilość zanieczyszczeń ropopochodnych może ulegać adsorpcji na powierzchni korzeni roślin przeprowadzających fitoremediację lub zostać zakumulowana w częściach nadziemnych, postanowiono doświadczalnie sprawdzić, czy istnieje korelacja pomiędzy efektywnością biodegradacji zanieczyszczeń w glebie, a transportem zanieczyszczeń do rośliny. Przeprowadzone analizy chromatograficzne wykazały obecność węglowodorów ropopochodnych (TPH i WWA) w korzeniach i pędach jeżówki purpurowej, z czego największe ich stężenia oznaczono w układach, w których stwierdzono największą efektywność biodegradacji zanieczyszczeń (Rys. 33). Świadczy to o zwiększonej biodostępności zanieczyszczeń w obecności biopreparatów oraz o zdolności mikroorganizmów w nich zawartych do rozkładu zanieczyszczeń ropopochodnych do lżejszych, łatwiej przyswajalnych przez rośliny frakcji. Dodatkowe wzbogacenie inokulacji biopreparatami o dodatek γ-PGA o potwierdzonych właściwościach surfaktantowych, sprzyjało zwiększonej adsorpcji

związków hydrofobowych. Ponadto warto zauważyć, że większa ilość zanieczyszczeń ropopochodnych została oznaczona w korzeniach roślin, co może wskazywać na rozkład zanieczyszczeń w tej części rośliny.




Rysunek 33 Oznaczone zawartości a) TPH b) WWA w korzeniach i pędach Echinacea purpurea poddanej analizowanych układów (system 1- brak inokulacji, system 2 – inokulacja biopreparatem B1, system 3 – inokulacja biopreparatem B1 i γ-PGA, system 4 – inokulacja biopreparatem B2, system 5 – inokulacja biopreparatem B2 i γ-PGA, system 6 – inokulacja γ-PGA).

Ocena toksykologiczna gleby z wykorzystaniem testów Phytotoxkit™, Ostracodtoxkit F™, Microtox SPT oraz MARA dopełniła kompleksowej oceny efektywności zaproponowanych technik bioremediacyjnych wyraźnie wskazując, że fitoremediacja wspomagana bioaugmentacją biopreparatem B2 z dodatkiem γ-PGA pozwoliła na uzyskanie najlepszego efektu oczyszczania oraz poprawy jakości gleb zanieczyszczonych węglowodorami ropopochodnymi. Potwierdzeniem tego były wartości współczynnika TU po zakończeniu eksperymentu wynoszące odpowiednio 2,6–

4 w teście Phytotoxkit™, 4,2 w teście Ostracodtoxkit F™, 4,4 w teście Microtox SPT oraz 4,0 w teście MARA, które stanowiły najniższe wartości spośród wszystkich wyników testów przeprowadzonych dla próbek gleb pochodzących z poszczególnych układów doświadczalnych (Rys. 34).

Wyniki testów toksykologicznych zaprezentowanych w artykule IV potwierdziły wyraźny spadek toksyczności gleb w układach obsadzonych i zaszczepionych biopreparatami na bazie mikroorganizmów autochtonicznych oraz ich modyfikowanymi wersjami, co wskazuje na potencjał zaproponowanych strategii oczyszczania do zastosowania w przyszłości w warunkach rzeczywistych.



Rysunek 34 Zestawienie wyników testów toksykologicznych gleby z dołu urobkowego poddanej fitoremediacji z wykorzystaniem Echinacea purpurea w różnych wariantach wzbogacenia gleby

5.3. Podsumowanie artykułów III i IV

W artykułach III i IV zaprezentowano dwie możliwości poprawy efektywności biodegradacji zanieczyszczeń ropopochodnych w glebach z terenu dołu urobkowego. W artykule III zwrócono uwagę na możliwość wzbogacenia biopreparatów na bazie bakterii autochtonicznych w wybrane szczepy grzybów autochtonicznych, wykazujących zdolność do rozkładu węglowodorów ropopochodnych. Ponadto w artykułach III i IV wskazano na możliwość wprowadzenia techniki fitoremediacji, która w połączeniu z bioaugmentacją odpowiednio dobraną mieszaniną inokulującą może istotnie wpłynąć na poprawę skuteczności oczyszczania gleby z dołu urobkowego. Kompleksowe badania prowadzone w warunkach półtechnicznych pozwoliły na sformułowanie najważniejszych wniosków wynikających z wyżej wymienionych artykułów:

- badania bioremediacyjne w skali półtechnicznej (Artykuł III) prowadzone na dwóch identycznych pryzmach glebowych umożliwiły porównanie skuteczności dwóch biopreparatów opartych na mikroorganizmach autochtonicznych: biopreparatu bakteryjnego B1 oraz biopreparatu bakteryjno-grzybowego B2. Wykazano, że wzbogacenie biopreparatu bakteryjnego o wyselekcjonowane szczepów grzybów autochtonicznych przyczyniło się do zwiększenia efektywności biodegradacji węglowodorów ropopochodnych, co znalazło odzwierciedlenie we wzroście stopnia biodegradacji TPH i WWA w glebie odpowiednio o 9,82% i 7,32% w porównaniu do biopreparatu bakteryjnego,
- wykonanie analiz chromatograficznych z wykorzystaniem autorskich procedur badawczych umożliwiło jakościowe i ilościowe oznaczenie poszczególnych węglowodorów ropopochodnych obecnych w badanej glebie oraz ocenę ich podatności na biodegradację. Wykazano, że obecność grzybów w biopreparacie przyczyniła się do zwiększenia efektywności biodegradacji wszystkich oznaczonych w glebie węglowodorów ropopochodnych, przy czym szczególnie wyraźną poprawę odnotowano w przypadku węglowodorów alifatycznych o dłuższych łańcuchach węglowych oraz WWA,
- przeprowadzony monitoring toksykologiczny wykazał, że inokulacja gleby biopreparatami B1 i B2 spowodowała wyraźne obniżenie toksyczności badanej gleby, co wskazuje na bezpieczeństwo ich stosowania oraz potencjał aplikacyjny w warunkach terenowych,
- badania fitoremediacji prowadzone z wykorzystaniem roślin *Zea mays* oraz *Echinacea purpurea* wykazały, że połączenie fitoremediacji z bioaugmentacją odpowiednio dobraną mieszaniną inokulacyjną prowadzi do istotnego zwiększenia efektywności bioremediacji gleb zanieczyszczonych węglowodorami ropopochodnymi. Wprowadzenie odpowiednio wyselekcjonowanych mikroorganizmów do strefy korzeniowej roślin fitoremediacyjnych intensyfikowało interakcje zachodzące pomiędzy mikroorganizmami rozkładającymi węglowodory, rośliną i glebą, czego efektem był wzrost biodostępności zanieczyszczeń i efektywności oczyszczania gleby. Jednocześnie wykazano, niską efektywność samej fitoremediacji w odniesieniu do gleb z dołów urobkowych zanieczyszczonych węglowodorami ropopochodnymi,
- badania z wykorzystaniem *Zea mays* wykazały, że sama fitoremediacja umożliwiła osiągnięcie biodegradacji TPH na poziomie 22,81% oraz WWA na poziomie 14,48%, natomiast jej połączenie z biopreparatem B2 zwiększyło efektywność oczyszczania odpowiednio o 26,99% i 22,26%. Dodatkowe wzbogacenie inokulacji biopreparatem B2 poporzez dodatek γ-PGA skutkowało dalszym wzrostem stopnia biodegradacji o 16,27% dla TPH oraz 20,06% dla WWA, co potwierdziło skuteczność γ-PGA w poprawie biodostępności zanieczyszczeń ropopochodnych i intensyfikacji procesów biodegradacyjnych,
- badania z wykorzystaniem *Echinacea purpurea* potwierdziły wysoką skuteczność fitoremediacji wspomaganej bioaugmentacją, przy czym efektywność wariantów inokulacji malała w kolejności: biopreparat B2 z dodatkiem γ-PGA > biopreparat

B1 z dodatkiem γ-PGA > biopreparat B2 > biopreparat B1 > γ-PGA > układ bez inokulacji, co jednoznacznie wskazuje na istotny wpływ składu mieszaniny inokulacyjnej na przebieg procesu,

- badania fitoremediacji zaprezentowane w artykule IV prowadzone na dwóch rodzajach gleby, tj. glebie z dołu urobkowego oraz glebie pochodzącej z naturalnego wycieku ropy naftowej umożliwiły weryfikację zaproponowanej strategii bioremediacyjnej w zróżnicowanych warunkach środowiskowych. Porównywalne wyniki badań potwierdziły możliwość zastosowania fitoremediacji wspomaganej bioaugmentacją w oczyszczaniu różnych typów gleb zanieczyszczonych węglowodorami ropopochodnymi,
- analiza morfologiczna roślin wykazała zależność pomiędzy efektywnością oczyszczania gleby, a kondycją roślin. Wykazano, że rozwój części nadziemnej i systemu korzeniowego zarówno *Zea mays*, jak i *Echinacea purpurea* był najlepszy w układach, w których odnotowano najwyższe stopnie biodegradacji zanieczyszczeń ropopochodnych,
- rozszerzenie zakresu analiz w artykule IV o oznaczenie zawartości TPH i WWA w korzeniach i pędach *Echinacea purpurea* wykazało, że niewielka część zanieczyszczeń może być pobierana przez rośliny. Przeprowadzone analizy chromatograficzne materiału roślinnego potwierdziły aktywny udział roślin w procesach usuwania zanieczyszczeń ropopochodnych z gleby, obejmujących zarówno mechanizmy akumulacji, jak i transformacji związków ropopochodnych. Ponadto wykazano, że ilość węglowodorów ropopochodnych oznaczonych w korzeniach i pędach *Echinacea purpurea* była zależna od zastosowanego wariantu inokulacji, co potwierdza wpływ bioaugmentacji na poprawę efektywności fitoekstrakcji zanieczyszczeń z gleby,
- weryfikacja skuteczności zaproponowanych strategii oczyszczania z wykorzystaniem analiz toksykologicznych wykazała poprawę jakości gleby we wszystkich analizowanych układach. Najniższe parametry toksyczności odnotowano dla gleb poddanych fitoremediacji wspomaganej bioaugmentacją z zastosowaniem biopreparatu B2 oraz dodatku γ-PGA, co potwierdza zasadność stosowania zintegrowanych metod remediacyjnych.

Wyniki badań omówionych w artykułach III i IV przyczyniły się do realizacji założonych celów niniejszej rozprawy doktorskiej oraz stanowiły podstawę do dalszych badań ukierunkowanych na wykorzystanie fitoremediacji wspomaganej bioaugmentacją w celu głębszego doczyszczenia gruntów z obszarów dołów urobkowych.

6. Zastosowanie roślin dziko rosnących w celu zwiększenia efektywności fitoremediacji gleb zanieczyszczonych węglowodorami ropopochodnymi – Artykuł V

Zadowalające wyniki badań uzyskanych w artykułach III i IV skłoniły do rozszerzenia zakresu prac nad wykorzystaniem fitoremediacji wspomaganej bioaugmentacją w oczyszczaniu gleb z terenów dołów urobkowych charakteryzujących się wysokim stopniem zanieczyszczenia węglowodorami ropopochodnymi. W badaniach przedstawionych w artykule V zaproponowano wykorzystanie fitoremediacji z zastosowaniem roślin dziko rosnących na terenie naturalnego wycieku ropy naftowej, zlokalizowanego w sąsiedztwie oczyszczanego dołu urobkowego. Założono, że gatunki przystosowane do wzrostu w warunkach długotrwałego skażenia mogą wykazywać zwiększoną tolerancję na obecność zanieczyszczeń, co istotnie może wpłynąć na poprawę efektywności procesów biodegradacyjnych w badanej glebie. W badaniach wykorzystano dwa gatunki roślin: *Scirpus sylvaticus* oraz *Cirsium oleraceum*. Na tamten moment przeprowadzone badania stanowiły jedną z pierwszych prób oceny skuteczności bioremediacji rzeczywistych gleb historycznie skażonych węglowodorami ropopochodnymi (TPH=4 320,7 mg/kg s.m., WWA= 7,19 mg/kg s.m.) z wykorzystaniem techniki inokulacji biopreparatem na bazie bakterii autochtonicznych w połączeniu z fitoremediacją z użyciem roślin dziko rosnących z gatunku *Scirpus sylvaticus* oraz *Cirsium oleraceum* pobranych z terenu naturalnego wycieku ropy naftowej, a także określenia interakcji zachodzących w badanych układach roślina–mikrobiota glebowa.

W artykule V przedstawiono kompleksowe badania fitoremediacji prowadzone w warunkach półtechnicznych z zastosowaniem metody wazonowej, obejmujące układy z roślinami *Scirpus sylvaticus* lub *Cirsium oleraceum*, zarówno w wariantach z bioaugmentacją, jak i bez jej zastosowania. Podejście to umożliwiło jednoczesne określenie skuteczności zaproponowanych strategii oczyszczania gleby, porównanie skuteczności oczyszczania gleby z wykorzystaniem różnych gatunków roślin oraz wykazanie wpływu bioaugmentacji na efektywność fitoremediacji. Weryfikację efektywności zaproponowanych strategii fitoremediacji przeprowadzono na podstawie analiz chromatograficznych, wagowych, toksykologicznych i mikrobiologicznych. Analiza chromatograficzna umożliwiła identyfikację oraz ilościowe oznaczenie węglowodorów wchodzących w skład zanieczyszczeń ropopochodnych oraz określenie stopnia biodegradacji tych związków po zakończeniu procesu oczyszczania. Dodatkowo pozwoliła na oznaczenie zawartości TPH i WWA w korzeniach i pędach roślin, co umożliwiło ocenę ich zdolności do akumulacji zanieczyszczeń. Analiza wagowa została wykorzystana do określenia biomasy roślin, natomiast analiza morfologiczna umożliwiła ocenę ogólnej kondycji roślin w poszczególnych wariantach badawczych. Uzupełnieniem były analizy toksykologiczne z zastosowaniem testów Phytotoxkit™, Ostracodtoxkit F™, Microtox SPT oraz MARA które pozwoliły na ocenę toksyczności gleby po zakończonych procesach fitoremediacji w odniesieniu do organizmów reprezentujących różne poziomy troficzne. Istotnym elementem badań były również analizy mikrobiologiczne, obejmujące ocenę zmian w strukturze zbiorowisk bakteryjnych,

istotnych z punktu widzenia procesów biodegradacyjnych. Uwzględnienie analizy bioróżnorodności mikroorganizmów w glebie po zakończeniu oczyszczania oraz włączenie do badań zbiorowisk orzęsków stanowiło istotny element nowości, rzadko dotychczas uwzględniany w badaniach nad bioremediacją gleb z terenów kopalnianych. Takie podejście pozwoliło nie tylko na ocenę efektywności usuwania zanieczyszczeń, lecz również na lepsze zrozumienie procesów zachodzących w glebie podczas jej rekultywacji, podkreślając wysoki potencjał tej strategii w praktycznych zastosowaniach.


6.1. Omówienie wyników zaprezentowanych w artykule V

Wyniki analiz chromatograficznych zaprezentowanych w artykule V wykazały, że efektywność fitoremediacji gleb zanieczyszczonych węglowodorami ropopochodnymi może zostać zwiększona dzięki zastosowaniu roślin naturalnie rosnących na terenach skażonych ropą naftową, najlepiej w pobliżu gruntów, które mają zostać poddane oczyszczaniu. W porównaniu do wyników zaprezentowanych w artykułach III i IV, efektywność biodegradacji węglowodorów ropopochodnych w glebach poddanych fitoremediacji z wykorzystaniem roślin dziko rosnących *Scirpus sylvaticus* i *Cirsium oleraceum* po sześciu miesiącach oczyszczania była zdecydowanie wyższa aniżeli w glebach odsadzonych roślinami uprawnymi *Zea mays* oraz ozdobnymi *Echinacea purpurea.* Efektywność biodegradacji TPH i WWA po zakończeniu eksperymentu wynosiła odpowiednio 36,24% i 31,99% dla fitoremediacji z udziałem *Scirpus sylvaticus* oraz 43,61% i 40,07% dla fitoremediacji z udziałem *Cirsium oleraceum*. Inokulacja gleby biopreparatem MC w połączeniu z fitoremediacją spowodowała jeszcze większą redukcję zanieczyszczeń ropopochodnych w glebie, która po zakończeniu eksperymentu w gruncie obsadzonym *Scirpus sylvaticus* wynosiła 89,47% (TPH) i 75,52% (WWA), natomiast w glebie obsadzonej *Cirsium oleraceum* 93,96% (TPH) i 79,55% (WWA). Uzyskane wyniki wykazały, że *Cirsium oleraceum* charakteryzuje się nieznacznie większym potencjałem do wykorzystania w fitoremediacji gleb zanieczyszczonych węglowodorami ropopochodnymi, aniżeli *Scirpus sylvaticus.*

Szczegółowa analiza chromatograficzna węglowodorów alifatycznych oraz WWA potwierdziła, że zastosowanie roślin dziko rosnących na terenach skażonych ropą naftową w fitoremediacji wspomaganej bioaumentacją, pozwoliło na uzyskanie wysokich wartości redukcji wszystkich węglowodorów oznaczonych w badanej glebie (Rys 35). Podatność węglowodorów ropopochodnych na rozkład mikrobiologiczny była zależna od długości łańcucha węglowego w cząsteczkach węglowodorów alifatycznych oraz ilości pierścieni aromatycznych w cząsteczkach węglowodorów aromatycznych. Węglowodory alifatyczne ulegały biodegradacji w kolejności: nC_{10}–nC_{21} > nC_6–nC_9 > nC_{22}–nC_{30} > nC_{31}–nC_{36}, natomiast WWA w kolejności: 2-pierścieniowy WWA > 3-pierścieniowe WWA > 4-pierścieniowe WWA > 5-pierścieniowe WWA > 6-pierścieniowe WWA. Wyniki te były zgodne z wynikami zaprezentowanymi w pozostałych artykułach będących podstawą niniejszej rozprawy doktorskiej.


Skuteczność fitoremediacji z wykorzystaniem roślin dziko rosnących została zweryfikowana za pomocą wskaźników biodegradacji, które w glebach obsadzonych *Scirpus sylvaticus* i inokulowanych biopreparatem MC zmalały z 2,64 do 0,13 (nC_{17}/Pr) oraz z 2,65 do 0,15 (nC_{18}/F). W glebie obsadzonej *Cirsium oleraceum* wartości

współczynników nC_{17}/Pr i nC_{18}/F po zakończeniu eksperymentu osiągnęły wartości odpowiednio 0,05 i 0,07. W glebach, które zostały poddane wyłącznie fitoremediacji wartości te były zdecydowanie wyższe, co potwierdza przewagę technik łączonych pod kątem efektywności usuwania zanieczyszczeń ropopochodnych .



Rysunek 35 Oznaczone zawartości n-alkanów i WWA w glebie a, d) wyjściowej i kontrolnej, b,e) poddanej fitoremediacji wspomaganej bioaugmentacją, c,f) poddanej wyłącznie fitoremediacji.


W artykule V rozszerzono badania o analizę materiału roślinnego, obejmującą analizę morfologiczną roślin, analizę biomasy, pomiar tempa wzrostu oraz określenie zawartości TPH i WWA w korzeniach i pędach roślin fitoremediacyjnych. Ocena morfologiczna wykazała lepszy rozwój korzeni i pędów roślin w glebach poddanych inokulacji biopreparatem, aniżeli w glebach wyłącznie obsadzonych, co przekładało się na większą produkcję biomasy i lepsze tempo wzrostu (Rys. 36). Sucha biomasa *Scirpus sylvaticus* i *Cirsium oleraceum* zasadzonych w glebach bioaugmentowanych była o 45,51% i 45,77% większa niż w glebach nie poddanych inokulacji. Lepszy rozwój roślin zasadzonych w glebach inokulowanych biopreparatem był skorelowany z wyższą efektywnością biodegradacji węglowodorów ropopochodnych oraz zwiększoną możliwością ich transportu z gleby do rośliny w trakcie fitoremediacji.



Rysunek 36 Biomasa korzeni i pędów: a) Scirpus sylvaticus, b) Cirsium oleraceum po zakończeniu procesów fitoremediacji w badanej glebie oraz przyrost pędów c) Scirpus sylvaticus, d) Cirsium oleraceum w czasie eksperymentu.

Analogicznie do badań zaprezentowanych w artykule IV, po zakończeniu eksperymentu oznaczono zawartość TPH i WWA w korzeniach i pędach roślin przeprowadzających fitoremediację (Rys. 37). Tutaj również większe ilości zanieczyszczeń ropopochodnych oznaczono w materiale roślinnym pobranym, z gleb poddanych bioaugmentacji, co dowodzi założenia, że inokulacja biopreparatem może wpływać na poprawę biodostępności zanieczyszczeń dla roślin fitoremediacyjnych. Poza tym większe ilości węglowodorów ropopochodnych oznaczono w korzeniach roślin, co potwierdza obserwacje z artykułu IV, w którym wskazano, że część zanieczyszczeń mogła ulec rozkładowi w tej części rośliny.

Badania przeprowadzone na dwóch gatunkach roślin różniących się cechami morfologicznymi i anatomicznymi pozwoliły na wykazanie korelacji pomiędzy typem rośliny, a zdolnością do pobierania i transportu zanieczyszczeń z gleby do jej tkanek. Ilość TPH i WWA oznaczona w korzeniach i pędach *Cirsium oleraceum* pobranych z gleb poddanych identycznym warunkom uprawy, była zdecydowanie większa niż w *Scirpus sylvaticus*, co można wiązać z różnicami w strategiach pobierania i translokacji zanieczyszczeń wynikającymi z odmiennych cech morfologicznych (szczególnie typu systemu korzeniowego) i fizjologicznych tych gatunków roślin.




Rysunek 37 Zawartość TPH i WWA w korzeniach i pędach a, c) Scirpus sylvaticus, b, d) Cirsium oleraceum przeprowadzających fitoremediację gleby z dołu urobkowego

Ocena skuteczności fitoremediacji z wykorzystaniem roślin dziko rosnących w badanych układach reakcyjnych, analogicznie do pozostałych artykułów omówionych w rozprawie doktorskiej, została uzupełniona o monitoring toksykologiczny z wykorzystaniem testów toksykologicznych Phytotoxkit™, Ostracodtoxkit F™, Microtox SPT oraz MARA (Rys. 38). Wyniki wszystkich testów wykazały poprawę jakości gleby w wyniku zaproponowanych strategii fitoremediacji, czego efektem był wzrost wartości współczynnika $EC_{50,}$ który w łatwy sposób można przeliczyć na wartość współczynnika toksyczności TU, co pozwala na porównanie wyników z pozostałymi badaniami omówionymi w niniejszej rozprawie doktorskiej.

Przeprowadzone analizy toksykologiczne wykazały, że gleba wyjściowa charakteryzowała się podwyższoną toksyczności, o czym świadczyły wartości współczynnika TU mieszczące się w przedziale 6,2-11,3 (w zależności od badanego parametru testu). Fitoremediacja gleby z wykorzystaniem roślin dziko rosnących wpłynęła na nieznaczną poprawę jakości gleby czego efektem były niewiele niższe

wartości TU po zakończeniu eksperymentu. Dopiero połączenie fitoremediacji z bioaugmentacją, pozwoliło na uzyskanie zadowalających wyników oczyszczania, czego dowodem było obniżenie wartości współczynników toksyczności TU w każdym z zastosowanych testów do wartości 2,1-2,4 dla gleby odsadzonej *Scirpus sylvaticus* i 1,3-1,7 dla gleby odsadzonej *Cirsium oleraceum*.




Rysunek 38 Zestawienie wyników testów toksykologicznych gleby z dołu urobkowego poddanej fitoremediacji z wykorzystaniem a) Scirpus sylvaticus b) Cirsium oleraceum w różnych wariantach wzbogacenia gleby

W niniejszym artykule w celu kompleksowej oceny procesu fitoremediacji oraz fitoremediacji wspomaganej bioagmentacją zdecydowano się rozszerzyć podstawowy zakres badań prezentowany w pozostałych artykułach (artykuły I-IV, VI-VII) o charakterystykę mikroorganizmów glebowych. Badania mikrobiologiczne wykazały zmiany w różnorodności mikrobiologicznej gleby po zakończeniu procesu oczyszczania, zróżnicowane w zależności od zastosowanego wariantu fitoremediacji i widoczne na poziomie rodzaju, rodziny i genu (Rys. 39). Zaobserwowano, że w wyniku fitoremediacji zanieczyszczonej gleby doszło do wyraźnego spadku liczebności archeonów metanogennych oraz typu *Desulfobacterota*, co może wskazywać na poprawę warunków tlenowych w zremediowanych glebach. Jednakże odnotowano również spadek liczebności niektórych bakterii tlenowych i fakultatywnie beztlenowych zwłaszcza z rodzajów *Agromyces, Cellulomonas, Galbitalea, Pseudoxanthomonas, Rhodoferax* i *Sulfuritalea*, co z kolei może sugerować, że zostały one wyeliminowane przez inne mikroorganizmy, lepiej przystosowane do warunków panujących w zremediowanych glebach lub uległy presji troficznej ze strony organizmów bakteriożernych np. z grupy proteozoa.

Istotnym wynikiem badań mikrobiologicznych, było wskazanie że w zremediowanych glebach najliczniej występowały mikroorganizmy zdolne do metabolizowania węglowodorów ropopochodnych, takie jak *Sphingomonas* (5,33–10,99%) i *Porticoccaceae* (2,51–6,28%), co potwierdziło skuteczność zaproponowanych strategii fitoremediacyjnych w poprawie efektywności biodegradacji węglowodorów ropopochodnych. Ponadto badania mikrobiologiczne wykazały, że w glebach zrekultywowanych pojawiły się bakterie nieobecne w glebie wyjściowej, ani w

zastosowanym biopreparacie. Może to sugerować, że część z nich stanowiła naturalny mikrobiom ryzosfery *Scirpus sylvaticus* i *Cirsium oleraceum*, w tym również bakterie zdolne do biodegradacji węglowodorów. Wykorzystanie w fitoremediacji roślin dziko rosnących na terenach zanieczyszczonych ropą naftową umożliwiło zatem wprowadzenie tych mikroorganizmów do oczyszczanej gleby, co mogło przyczynić się do wyższej efektywności biodegradacji węglowodorów ropopochodnych.




Rysunek 39 Wpływ fitoremediacji i bioaugmentacji na względną liczebność taksonów bakteryjnych: a) typ, b) rodzina (20 najliczniej występujących), c) rodzaj (20 najliczniej występujących).

Przeprowadzona w ramach badań mikrobiologicznych analiza RA (z ang. *Rarefaction analysis*) potwierdziła korzystny wpływ fitoremediacji z wykorzystaniem roślin z gatunku *Scirpus sylvaticus* i *Cirsium oleraceum* w połączeniu z inokulacją gleby biopreparatem MC na różnorodność mikrobiologiczną gleby względem próbki referencyjnej. Największy wzrost różnorodności społeczności mikroorganizmów glebowych zaobserwowano na poziomie rodzaju, gdzie liczba zidentyfikowanych rodzajów bakterii wzrosła z około 350 do ponad 500. Uzyskane wyniki stanowią uzupełnienie wcześniejszych obserwacji dotyczących zmian w strukturze społeczności bakteryjnej oczyszczanej gleby, wskazując, że fitoremediacja wspomagana

bioaugmentacją sprzyja wzbogaceniu społeczności mikrobiologicznych oraz poprawie funkcjonalności enzymatycznej gleby.

Ponadto zaprezentowana w artykule analiza RDA (z ang. *Redundancy analysis*) oparta na selekcji zmiennych środowiskowych, wykazała, że głównymi czynnikami kształtującymi strukturę mikrobioty w trakcie eksperymentu były zanieczyszczona gleba G6 i obecność konsorcjum MC, które wyjaśniały odpowiednio 21% i 18% zmienności składu taksonomicznego mikrobioty glebowej (Rys. 40).




Rysunek 40 Analiza RDA czynników środowiskowych (strzałki) i typów mikroorganizmów (okręgi) w przestrzeni dwuwymiarowej.

Istotnym elementem badań przedstawionych w artykule V było zbadanie różnorodności i liczebności orzęsków w glebach po zakończeniu oczyszczania, co umożliwiło nie tylko ocenę skuteczności remediacji za pomocą wskaźników mikrobiologicznych, lecz także identyfikację potencjalnych bioindykatorów regeneracji i stabilności ekosystemu glebowego.

Analiza społeczności orzęsków wykazała, wzrost liczebności oraz różnorodności gatunkowej protistów z typu Ciliophora w glebach zremediowanych. Wskazano że liczebność orzęsków wzrosła z 9 gatunków (gleba wyjściowa) do 21 gatunków w glebie obsadzonej *Scirpus sylvaticus* oraz 17 gatunków w glebie obsadzonej *Cirsium oleraceum.* Różnice pomiędzy glebą zanieczyszczoną a glebami po fitoremediacji zostały potwierdzone analizą wskaźników bioróżnorodności, obejmujących wskaźniki różnorodności (Shannona i Simpsona), równomierności (Shannon equitability, Simpson evenness oraz Buzas–Gibson evenness) oraz bogactwa gatunkowego (Margalef richness). (Rys. 41).

Przeprowadzone analizy społeczności orzęsków w glebie wykazały, że fitoremediacja wspomagana bioaugmentacją początkowo mogła ograniczać liczebność gatunków orzęsków w badanych glebach, jednak ostatecznie umożliwiła odbudowę populacji kluczowych bakteriożerców, podczas gdy sama fitoremediacja sprzyjała wyższej początkowej liczebności gatunków oraz większej różnorodności funkcjonalnej.



Rysunek 41 Zmiany wartości głównych wskaźników bioróżnorodności ekologicznej w czasie w glebie G6 (ciemnoniebieska linia) oraz w glebach obsadzonych Scirpus sylvaticus : SS-G6-A (pomarańczowy), SS-G6-B (szary) oraz Cirsium oleraceum CO-G6-A (żółty) i CO-G6-B (zielony) w różnych wariantach fitoremediacji.

6.2. Podsumowanie artykułu V

W artykule V przedstawiono możliwość wykorzystania roślin dziko rosnących na obszarach naturalnego wycieku ropy naftowej zlokalizowanego w sąsiedztwie przeznaczonego do oczyszczania gruntu z dołu urobkowego w procesie fitoremediacji i fitoremediacji wspomaganej bioaugmentacja. Takie podejście ułatwiało lepszą adaptację roślin przeprowadzających fitoremediację do zanieczyszczonego podłoża, co przełożyło się na wzrost efektywności degradacji zarówno węglowodorów alifatycznych, jak i

wielopierścieniowych węglowodorów aromatycznych oraz obniżenie toksyczności gleby uznawanego za bezpieczny dla biocenozy glebowej. Kompleksowe badania prowadzone w warunkach zbliżonych do rzeczywistych pozwoliły na sformułowanie najważniejszych wniosków które wynikały z artykułu V:

- badania zaprezentowane w artykule V wykazały, że fitoremediacja gleby z dołu urobkowego z wykorzystaniem roślin naturalnie przystosowanych do wzrostu na terenach skażonych ropą naftową pozwala na uzyskanie wyraźnie wyższej efektywność bioremediacji niż z wykorzystaniem roślin uprawnych i ozdobnych stosowanych w artykułach III i IV (*Zea mays*, *Echinacea purpurea*). Fitoremediacja z użyciem *Scirpus sylvaticus* pozwoliła na redukcję TPH i WWA w badanej glebie o odpowiednio 36,24% i 31,99%, natomiast zastosowanie *Cirsium oleraceum* umożliwiło obniżenie tych zanieczyszczeń o 43,61% i 40,07%. Fitoremediacja wspomagana bioaugmentacją z wykorzystaniem biopreparatu MC w układzie z *Scirpus sylvaticus* zwiększyła efektywność oczyszczania do 89,47% dla TPH i 75,52% dla WWA, natomiast w układzie z *Cirsium oleraceum* do 93,69% dla TPH i 79,55% dla WWA, co wskazuje na zasadność stosowania strategii łączonej w celu intensyfikacji procesu oczyszczania gleb z dołów urobkowych,
- analiza chromatograficzna wykonana z zastosowaniem autorskich procedur badawczych umożliwiła identyfikację i ilościowe oznaczenie poszczególnych węglowodorów ropopochodnych obecnych w glebie w trakcie jej oczyszczania, co pozwoliło na ocenę podatności analizowanych zanieczyszczeń na biodegradację oraz wyznaczenie wskaźników n-C_{17}/Pr i n-C_{18}/F jako miar efektywności zastosowanych strategii bioremediacyjnych. Wykazano, że węglowodory alifatyczne ulegały biodegradacji w kolejności n-C_{10}–n-C_{21} > n-C_6–n-C_9 > n-C_{22}–n-C_{30} > n-C_{31}–n-C_{36}, natomiast wielopierścieniowe węglowodory aromatyczne w kolejności WWA dwupierścieniowe > trójpierścieniowe > czteropierścieniowe > pięciopierścieniowe > sześciopierścieniowe, co wynikało z ich budowy chemicznej oraz właściwości fizykochemicznych.
- analiza morfologiczna roślin potwierdziła obserwacje z artykułów III i IV dotyczące zależności pomiędzy efektywnością oczyszczania gleby, a kondycją roślin fitoremediacyjnych,
- rozszerzenie zakresu badań w artykule V o analizę biomasy i tempa wzrostu roślin potwierdziło wpływ zastosowania strategii łączącej fitoremediację z bioaugmentacją na wzrost roślin. Wykazano o ok. 45% większą produkcję biomasy *Scirpus sylvaticus* i *Cirsium oleraceum* w układach inokulowanych niż w układach nieinokulowanych. Analogicznie biomasa korzeni i pędów w układach inokulowanych była wyższa o odpowiednio 51,5% i 41,7% dla *Scirpus sylvaticus* oraz 35,0% i 59,1% dla *Cirsium oleraceum,*
- analiza zawartości TPH i WWA w korzeniach i pędach wykazała aktywny udział testowanych roślin dziko rosnących w procesie oczyszczania, przy czym wyższe stężenia zanieczyszczeń oznaczono w korzeniach *Scirpus sylvaticus* i *Cirsium oleraceum*. Może to wskazywać na adsorpcję zanieczyszczeń ropopochodnych na

powierzchni korzeni, akumulację oraz możliwość ich częściowej degradacji i transformacji w tkankach korzeniowych,

- analizy chromatograficzne gleby i roślin pod kątem zawartości węglowodorów ropopochodnych w połączeniu z analizą biomasy wyraźnie wskazały, że rośliny w procesie fitoremediacji mogą pełnić podwójną funkcję. Po pierwsze mogą wspomagać biodegradację zanieczyszczeń poprzez poprawę warunków w strefie korzeniowej tj. zwiększenie biodostępności zanieczyszczeń, lepsze napowietrzenie gleby czy dostarczanie składników pokarmowych dla mikroorganizmów prowadzących degradację. Po drugie w ograniczonym stopniu mogą bezpośrednio uczestniczyć w usuwaniu zanieczyszczeń z gleby na drodze fitoekstrakcji,
- przeprowadzona analiza toksykologiczna gleb po zakończeniu eksperymentu potwierdziła skuteczność fitoremediacji wspomaganej bioaugmentacją z wykorzystaniem roślin dziko rosnących, czego potwierdzeniem były obniżone wartości współczynników toksyczności TU w testach Phytotoxkit™, Ostracodtoxkit F™, Microtox® SPT i MARA.
- Zastosowanie jako fitoremedianta roślin dziko rosnących należących do gatunków *Scirpus sylvaticus* i *Cirsium oleraceum* pochodzących z naturalnego wycieku ropy naftowej dowodzi, że są one odpowiednimi kandydatami do wykorzystania w procesie fitoremediacji gleb z dołów urobkowych, gdyż są w stanie zwiększyć potencjał biodegradacyjny oczyszczanej gleby poprzez wzbogacenie go o taksony wchodzące w skład mikrobiomów zamieszkujących ryzosferę obu gatunków roślin.
- Charakterystyka społeczności bakteryjnej po zakończeniu fitoremediacji wykazała zróżnicowanie struktury oraz różnorodności mikrobiologicznej gleby w zależności od zastosowanej strategii oczyszczania. Wykazano, że w procesie degradacji węglowodorów ropopochodnych w glebie z dołu urobkowego dominującą rolę odgrywały mikroorganizmy tlenowe. Poza tym w glebach poddanych fitoremediacji z wykorzystaniem roślin dziko rosnących stwierdzono obecność bakterii nieobecnych w glebie wyjściowej. Sugeruje to, że część z nich mogła stanowić element naturalnego mikrobiomu ryzosfery *Scirpus sylvaticus* oraz *Cirsium oleraceum*, w tym również bakterii o potencjale degradacyjnym w stosunku do węglowodorów ropopochodnych. Wyniki te wskazują, że wykorzystanie roślin dziko rosnących na terenach zanieczyszczonych ropą naftową w procesie fitoremediacji gleby z dołu urobkowego mogło sprzyjać wprowadzaniu do gleby mikroorganizmów wspierających biodegradację, co może przyczynić się do zwiększenia efektywności procesu oczyszczania.
- przeprowadzona analiza RDA wykazała, że głównymi czynnikami kształtującymi strukturę mikrobioty po zakończeniu fitoremediacji były rodzaj gleby oraz obecność biopreparatu MC,
- rozszerzenie zakresu badań o charakterystykę społeczności orzęsków, mogących pełnić rolę bioindykatorów, wykazało korzystny wpływ fitoremediacji na różnorodność i liczebność tych organizmów w glebie skażonej węglowodorami ropopochodnymi. W surowej glebie z dołu urobkowego stwierdzono obecność 9 gatunków orzęsek. Przeprowadzenie procesu fitoremediacji doprowadziło do wzrostu liczebności oraz różnorodności orzęsków, co było szczególnie widoczne w

układach bioaugmentowanych. W układach z *Scirpus sylvaticus* odnotowano 21 gatunków, natomiast w układach z *Cirsium oleraceum* stwierdzono 17 gatunków. Korzystny wpływ fitoremediacji na społeczność orzęsków wynikał prawdopodobnie ze zwiększonej dostępności i mobilności składników odżywczych w strefie korzeniowej, co sprzyjało rozwojowi mikroorganizmów i pośrednio wspierało regenerację grup troficznych orzęsków w glebie zanieczyszczonej ropą naftową. Zgodnie z dostępną wiedzą było to jedno z pierwszych badań analizujące strukturę społeczności orzęsków w procesach fitoremediacji gleb zanieczyszczonych węglowodorami ropopochodnymi.

Wyniki badań omówionych w artykule V przyczyniły się do realizacji założonych celów niniejszej rozprawy doktorskiej oraz stanowiły istotne źródło wiedzy na temat możliwości poprawy efektywności bioremediacji gleby z dołu urobkowego. Wykazano, że wykorzystanie w procesie fitoremediacji dziko rosnących roślin przystosowanych do wzrostu na terenach zanieczyszczonych ropą naftową, pozwala na istotną poprawę efektywności oczyszczania gleby z dołu urobkowego.

7. Wpływ nanomateriałów na efektywność bioremediacji gleby z dołu urobkowego – Artykuł VI

W artykule VI przedstawiono wyniki badań, w których zastosowano odmienną, zintegrowaną strategię bioremediacji gleb zanieczyszczonych węglowodorami ropopochodnymi, polegającą na połączeniu bioaugmentacji, wzbogacenia gleby w nanomateriały oraz fitoremediacji. Celem badań było nie tylko zwiększenie efektywności biodegradacji zanieczyszczeń, lecz również kompleksowa ocena skuteczności takiego podejścia w warunkach laboratoryjnych i półtechnicznych. W artykule wskazano, że zastosowanie nanomateriałów może stanowić skuteczne uzupełnienie strategii biologicznych, ze względu na ich właściwości fizykochemiczne, takie jak wysoka powierzchnia właściwa, reaktywność oraz zdolność do oddziaływania z zanieczyszczeniami organicznymi. Jednocześnie podkreślono, że połączenie inokulacji biopreparatem na bazie bakterii autochtonicznych ze wzbogaceniem gleby w nanomateriały nie było dotychczas szeroko opisywane w literaturze, co stanowiło istotny element nowości prowadzonych badań bioremediacyjnych.

W artykule VI przedstawiono wyniki badań biodegradacji zanieczyszczeń ropopochodnych w glebie pochodzącej z dołu urobkowego, poddanej inokulacji biopreparatem MC oraz wzbogaceniu nanomateriałami: tlenkiem grafenu (GO-NP) oraz tlenkiem cynku (ZnO-NP). Badania biodegradacji realizowano w skali laboratoryjnej z wykorzystaniem analiz respirometrycznych oraz chromatograficznych, na podstawie których dokonano oceny aktywności mikroorganizmów oraz stopnia degradacji zanieczyszczeń w skażonej glebie. Istotnym aspektem przeprowadzonych badań laboratoryjnych było zastosowanie różnych dawek nanomateriałów do wzbogacenia gleby, co umożliwiło określenie ich wpływu na przebieg procesów biodegradacyjnych oraz wyznaczenie dawki nanomateriałów zapewniającej największą efektywność biodegradacji zanieczyszczeń ropopochodnych. Uzyskane wyniki stanowiły podstawę do zaplanowania i realizacji kolejnego etapu badań z wykorzystaniem fitoremediacji.


Badania fitoremediacji realizowano w skali półtechnicznej z wykorzystaniem metody wazonowej z zastosowaniem *Lolium perenne*, jako rośliny fitoremediacyjnej. Celem badań była ocena skuteczności strategii łączącej bioaugmentację, wzbogacenie gleby w nanomateriały i fitoremediacje w procesie oczyszczania gleby zanieczyszczonej węglowodorami ropopochodnymi oraz weryfikacja wpływu wcześniej wyznaczonej optymalnej dawki nanomateriałów na efektywność tego procesu. Oceny efektywności zaproponowanej strategii bioremediacyjnej dokonano na podstawie analiz chromatograficznych gleby pod kątem zawartości węglowodorów alifatycznych oraz WWA, a także zawartości tych związków w korzeniach i pędach *Lolium perenne,* co dodatkowo umożliwiło analizę ich dystrybucji w układzie gleba–roślina. Ponadto w artykule VI przeprowadzono analizę biomasy roślin po zakończeniu fitoremediacji oraz ocenę ekotoksykologiczną gleby, co pozwoliło na kompleksową ocenę skuteczności zastosowanej strategii oczyszczania.

Wyniki badań zaprezentowane w artykule VI wskazują, że wzbogacenie gleby w nanomateriały (przy odpowiednio dobranej dawce nanomateriału) w połączeniu z

bioaugmentacją i fitoremediacją, może istotnie przyczynić się do zwiększenia efektywności biodegradacji zanieczyszczeń oraz poprawy jakości gleby, a zaproponowane rozwiązanie może stanowić podstawę do dalszych badań nad jego zastosowaniem w warunkach terenowych.

7.1. Omówienie wyników badań biodegradacji zaprezentowanych w artykule VI

Wyniki analiz respirometrycznych zaprezentowanych w artykule VI, wykazały wpływ zastosowanych nanomateriałów na aktywność mikrobiologiczną w układzie, odzwierciedlaną poprzez wzrost zużycia tlenu w układach reakcyjnych. W układzie kontrolnym, zaszczepionym biopreparatem MC, zużycie tlenu po 90 dniach testu wyniosło 1 944,8 mg O_2/dm^3, co wskazywało na gorszy efekt biodegradacji zanieczyszczeń niż w układach wzbogaconych w nanomateriały. W próbkach gleby inokulowanej i wzbogaconej w nanocząstki tlenku grafenu (GO-NP) zużycie tlenu po zakończeniu eksperymentu kształtowało się w granicach od 2 441 mg O_2/dm^3 do 5 147,8 mg O_2/dm^3, natomiast w próbkach inokulowanych i wzbogaconych w nanocząstki tlenku cynku (ZnO-NP) mieściło się w zakresie od 2 025,3 mg O_2/dm^3 do 4836,6 mg O_2/dm^3. Na podstawie analiz respirometrycznych stwierdzono, że efektywność biodegradacji zanieczyszczeń ropopochodnych w układach reakcyjnych była silnie zależna od dawki zastosowanego nanomateriału. W naszych badaniach optymalna ilość nanomateriałów wprowadzonych do gleby wynosiła 20 mg/kg s.m. gleby, podczas gdy najmniejszą skuteczność w poprawie aktywności mikrobiologicznej w układzie odnotowano przy 100 mg/kg s.m. gleby (Rys. 42).




Rysunek 42 Zależność ilości zużytego tlenu [mg/dm³] od czasu w glebie G6 inokulowanej biopreparatem oraz wzbogaconej w a) GO-NP, b)ZnO-NP.

Wyniki analiz chromatograficznych potwierdziły zależność pomiędzy efektywnością biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego, a ilością wprowadzonych nanomateriałów do gleby inokulowanej biopreparatem MC.


Efektywność biodegradacji TPH w glebie wyłącznie inokulowanej biopreparatem po 90 dniach badań wynosiła 44,23%. Wprowadzenie GO-NP do gleby przyczyniło się do wzrostu efektywności biodegradacji TPH, która po zakończeniu eksperymentu mieściła się w zakresie 50,41–70,47%, w zależności od ilości nanomateriału wprowadzonego do układu reakcyjnego. Warto zauważyć, że modyfikacja metody bioremediacji polegająca na wzbogaceniu gleby inokulowanej biopreparatem MC w GO-NP w ilości 20 mg/kg s.m. gleby przyczyniła się do poprawy efektywności biodegradacji TPH aż o 23,24%. Zbliżone, lecz nieco niższe wartości stopni biodegradacji odnotowano w przypadku wzbogacenia gleby w ZnO-NP. Po zakończeniu eksperymentu stopienie biodegradacji TPH w próbkach inokulowanych MC i wzbogaconych w ZnO-NP mieściły się w przedziale 49,69–67,68%, co świadczy o poprawie efektywności biodegradacji TPH w zakresie od 5,46-23,45%.

Podobnie zależności zaobserwowano w przypadku WWA, czego potwierdzeniem są obliczone wartości stopni biodegradacji tych związków po zakończeniu eksperymentu kształtujące się na poziomie 36,98% w glebie inokulowanej MC, 40,48–56,19% w glebie inokulowanej MC i wzbogaconej w GO-NP oraz 40,45–54,94% w glebie inokulowanej MC i wzbogaconej w ZnO-NP. Ponadto wykazano, że modyfikacja procesu biodegradacji prowadzonego metodą bioaugmentacji biopreparatem MC, poprzez wzbogacenie gleby w GO-NP lub ZnO-NP w ilości 20 mg/kg s. m. (dawka optymalna) przyczyniła się do poprawy efektywności biodegradacji WWA w glebie zdołu urobkowego o odpowiednio 19,21% i 17,96%.

Ponadto szczegółowa analiza chromatograficzna próbek gleby po zakończeniu eksperymentu wykazała, że zaproponowana strategia modyfikacji procesu bioaugmentacji, polegająca na dodatkowym wprowadzeniu do gleby nanomateriałów (GO-NP lub ZnO-NP.), istotnie wpłynęła na poprawę efektywności degradacji wszystkich oznaczonych w próbce węglowodorów ropopochodnych. Uzyskane wyniki wskazują, że stopień ich biodegradacji był zróżnicowany i zależał od struktury chemicznej cząsteczki, tj. od długości łańcucha węglowego w przypadku węglowodorów alifatycznych oraz liczby pierścieni aromatycznych w cząsteczce w przypadku WWA. (Rys. 43, Rys. 44). Obserwacje te są zgodne z typowymi zależnościami obserwowanymi w procesach biodegradacji węglowodorów ropopochodnych i mogą wskazywać, że zastosowane nanomateriały wpływały przede wszystkim na zwiększenie biodostępności zanieczyszczeń dla mikroorganizmów odpowiedzialnych za ich rozkład, co w konsekwencji mogło przyczyniać się do intensyfikacji procesu biodegradacji.



Rysunek 43 Średni stopień biodegradacji wybranych grup węglowodorów alifatycznych w glebie G6 inokulowanej biopreparatem oraz wzbogaconej w a) GO-NP, b)ZnO-NP. Różne małe litery oznaczają istotne statystycznie różnice między wariantami zgodnie z jednoczynnikową analizą wariancji (ANOVA) z testem post-hoc Tukeya HSD ($p < 0,05$).



Rysunek 44 Średni stopień biodegradacji wybranych grup WWA w glebie G6 inokulowanej biopreparatem oraz wzbogaconej w a) GO-NP, b)ZnO-NP. Różne małe litery oznaczają istotne statystycznie różnice między wariantami zgodnie z jednoczynnikową analizą wariancji (ANOVA) z testem post-hoc Tukeya HSD ($p < 0,05$).

Przeprowadzone badania laboratoryjne zaprezentowane w artykule VI wykazały, że wzbogacenie gleby w nanomateriały w połączeniu z inokulacją biopreparatem na bazie wyselekcjonowanych bakterii autochtonicznych może istotnie przyczynić się do poprawy efektywności oczyszczania gleb z terenów dołów urobkowych i stanowić ciekawą alternatywę dla tradycyjnych technik bioremediacyjnych.


Zadowalające wyniki przeprowadzonych badań biodegradacji, obejmujących analizy respirometryczne i chromatograficzne, stanowiły podstawę do podjęcia decyzji o rozszerzeniu badań z wykorzystaniem tych nanomateriałów oraz ich zastosowaniu w procesie fitoremediacji.

7.2. Omówienie wyników badań fitoremediacji wspomaganej bioaugmentacją i nanomateriałami zaprezentowanych w artykule VI

Badania fitoremediacji prowadzono w układzie kontrolnym (gleba obsadzona *Lolium perenne* i inokulowana biopreparatem MC) oraz w dwóch układach eksperymentalnych, w których gleba została dodatkowo wzbogacona w nanomateriały GO-NP i ZnO-NP. Ilość nanomateriałów wprowadzonych do gleby ustalono na podstawie wcześniejszych wyników biodegradacji i wynosiła 20 mg/kg s.m. gleby.

Analizy chromatograficzne gleby po zakończeniu badań fitoremediacji wykazały duży spadek stężenia TPH i WWA w wyniku zaproponowanych strategii bioremediacyjnych, jednak w układach wzbogaconych w nanomateriały był on zdecydowanie większy, aniżeli w układzie kontrolnym. Efektywność biodegradacji TPH i WWA w glebie obsadzonej *Lolium perenne* i inokulowanej biopreparatem wynosiła odpowiednio 63,22% i 61,86%, co wskazywało na na wysoką skuteczność zastosowanego układu fitoremediacyjnego oraz jego potencjał w intensyfikacji procesów biodegradacji zanieczyszczeń ropopochodnych.. Dodatkowe wzbogacenie gleby w GO-NP skutkowało dalszym wzrostem stopni biodegradacji TPH (81,85%) i WWA do (73,19%), co przekładało się na poprawę efektywności biodegradacji tych zanieczyszczeń o odpowiednio 18,63% i 11,33%. Nieznacznie niższe stopnie biodegradacji TPH i WWA odnotowano w glebie inokulowanej biopreparatem i wzbogaconej w ZnO-NP, które po zakończeniu eksperymentu kształtowały się na poziomie odpowiednio 80,39% i 70,66%. Uzyskane wyniki świadczyły o wzroście efektywności biodegradacji TPH i WWA w porównaniu z układem kontrolnym o 17,17% i 8,8%.

Szczegółowe analizy chromatograficzne węglowodorów alifatycznych i WWA potwierdziły poprawę biodegradacji wszystkich węglowodorów ropopochodnych w obecności GO-NP i ZnO-NP, co potwierdziło obserwacje poczynione na etapie badań laboratoryjnych, dotyczące wpływu nanomateriałów na poprawę biodostępności zanieczyszczeń dla mikroorganizmów rozkładających węglowodory (Rys. 45).




Rysunek 45 Średni stopień biodegradacji wybranych grup a) węglowodorów alifatycznych, b) WWA w glebie G6 poddanej fitoremediacji z użyciem Lolium perenne w różnych wariantach wzbogacenia gleby. Różne małe litery oznaczają istotne statystycznie różnice między wariantami zgodnie z jednoczynnikową analizą wariancji (ANOVA) z testem post-hoc Tukeya HSD ($p < 0,05$).

Analiza materiału roślinnego obejmująca analizę morfologiczną, analizę biomasy oraz analizę korzeni i pędów *Lolium perenne* pod kątem zawartości TPH i WWA, wykazały korelację pomiędzy efektywnością biodegradacji zanieczyszczeń w glebie, a ogólnym stanem roślin. W glebach wzbogaconych w nanomateriały zaobserwowano lepszy rozwój systemu korzeniowego oraz części nadziemnych (liści) *Lolium perenne*, co przejawiało się większą intensywnością wzrostu oraz wyższą biomasą w porównaniu z układem kontrolnym (Rys. 46).



Rysunek 46 Zjęcie Lolium perenne po miesiącu wzrostu na glebie G6 poddanej fitoremediacji w badanych wariantach: (1) gleba G6 + Lolium perenne + biopreparat; (2) gleba G6 + Lolium perenne + biopreparat + GO-NP (20 mg/kg s.m.); (3) gleba G6 + Lolium perenne + biopreparat + ZnO-NP (20 mg/kg s.m.).


Analiza suchej biomasy potwierdziła, że wzbogacenie gleby w nanomateriały pozytywnie wpłynęło na rozwój *Lolium perenne*. W porównaniu z układem kontrolnym (gleba poddana wyłącznie bioaugmentacji) sucha biomasa roślin wzrosła o 28,63% przy zastosowaniu GO-NP i o 22,21% przy zastosowaniu ZnO-NP. Analiza biomasy z rozdzieleniem na biomasę korzeni i pędów wykazała podobne zależności, mianowicie sucha biomasa korzeni *Lolium perenne* pobranych z gleb wzbogaconych w nanomateriały była o 29,70-30,84% większa niż z gleby kontrolnej, natomiast sucha biomasa pędów większa o 18,65-27,58% (Rys. 47).



Rysunek 47 Biomasa korzeni i pędów Lolium perenne pobranej z gleby poddanej fitoremediacji w różnych wariantach wzbogacenia gleby. Różne małe litery oznaczają istotne statystycznie różnice między wariantami zgodnie z jednoczynnikową analizą wariancji (ANOVA) z testem post-hoc Tukeya HSD ($p < 0,05$).

Przeprowadzona analiza materiału roślinnego pod kątem zawartości TPH i WWA w korzeniach oraz pędach *Lolium perenne* wykazała, że niewielkie ilości zanieczyszczeń ropopochodnych mogą być pobierane z gleby przez życicę trwałą, a następnie ulegać dalszym przemianom metabolicznym w jej tkankach. Ponadto badania prowadzone w różnych układach wzbogacenia gleby wskazały, że zastosowanie nanomateriałów w połączeniu z bioaugmentacją istotnie zwiększało akumulację analizowanych zanieczyszczeń w roślinach fitoremediacyjnych, co świadczy o intensyfikacji procesów ich pobierania z gleby w obecności GO-NP i ZnO-NP.

W układach wzbogaconych w nanomateriały zawartość TPH w korzeniach *Lolium perenne* była wyższa o 69,35–74,59%, natomiast w pędach o 82,19–90,67% w porównaniu z próbami kontrolnymi. Analogiczny trend zaobserwowano w tych samych układach dla WWA, których stężenie wzrosło odpowiednio o 55,31–59,84% w korzeniach oraz o 82,74–91,03% w pędach *Lolium perenne* (Rys. 48).




Rysunek 48 Stężenie a) TPH oraz b) WWA w korzeniach i pędach Lolium perenne pobranej z gleby poddanej fitoremediacji w różnych wariantach wzbogacenia gleby. Różne małe litery oznaczają istotne statystycznie różnice między wariantami zgodnie z jednoczynnikową analizą wariancji (ANOVA) z testem post-hoc Tukeya HSD ($p < 0,05$).

Dopełnieniem oceny efektywności zaproponowanych strategii bioremediacyjnych omówionych w artykule VI, była analiza ekotoksykologiczna badanych gleb po zakończeniu eksperymentu, której analogicznie do pozostałych artykułów omówionych w niniejszej rozprawie doktorskiej dokonano za pomocą pakietu testów toksykologicznych Phytotoxkit™, Ostracodtoxkit F™ oraz Microtox SPT. W przypadku wykorzystania w badaniach fitoremediacji nanomateriałów (GO-NP, ZnO-NP) ocena ta miała szczególne znaczenie ze względu na nadal niewystarczająco rozpoznany wpływ tych związków na środowisko glebowe oraz inne organizmy żywe.

Przeprowadzone analizy toksykologiczne wykazały że fitoremediacja gleby G6 charakteryzującej się podwyższonym poziomem toksyczności (wartości współczynników toksyczności TU w przeprowadzonych testach mieściły się w przedziale 9,02–13,05), z wykorzystaniem *Lolium perenne* jako fitoremedianta oraz inokulacji biopreparatem MC umożliwiła znaczne obniżenie jej toksyczności (wartości TU=5,14–6,81). Dodatkowe zbogacenie gleby w nanomateriały skutkowało uzyskaniem jeszcze lepszych wyników testów toksykologicznych, czego efektem było obniżenie wartości współczynników toksyczności TU w każdym z zastosowanych testów do wartości mieszczących się w przedziałach 2,5-2,82 dla gleby wzbogaconej w GO-NP i 2,70-3,30 dla gleby wzbogaconej w ZnO-NP. Uzyskane wyniki testów toksykologicznych potwierdziły, że modyfikacja fitoremediacji wspomaganej bioaugmentacją poprzez dodatkowe wzbogacenie gleby w GO-NP lub ZnO-NP (w odpowiednio dobranych dawkach) nie ogranicza się wyłącznie do redukcji stężenia zanieczyszczeń ropopochodnych, lecz prowadzi również do poprawy jakości środowiska glebowego. Ponadto wyniki przeprowadzonych testów toksykologicznych wskazują, że GO-NP i ZnO-NP wprowadzone do gleby w dawce 20 mg/kg s.m. nie wywołały efektu toksycznego u

zastosowanych w testach organizmów wskaźnikowych, co potwierdza bezpieczeństwo zastosowania proponowanej strategii bioremediacyjnej w warunkach terenowych. (Rys. 49).



Rysunek 49 Zestawienie wyników testów toksykologicznych gleby z dołu urobkowego poddanej fitoremediacji z wykorzystaniem Lolium perenne w różnych wariantach wzbogacenia gleby.

7.3 Podsumowanie artykułu VI

W artykule VI przedstawiono wyniki badań dotyczące możliwości zastosowania nanocząstek tlenku grafenu oraz tlenku cynku jako czynnika wspomagającego procesy bioremediacji gleb historycznie zanieczyszczonych węglowodorami ropopochodnymi. Kompleksowe badania biodegradacji prowadzone w warunkach laboratoryjnych w połączeniu z wynikami badań fitoremediacji prowadzonych metodą wazonową w warunkach półtechnicznych pozwoliły na sformułowanie najważniejszych wniosków wynikających z artykułu VI:

- badania biodegradacji i fitoremediacji wykazały, że wzbogacenie gleby w nanomateriały w połączeniu z bioaugmentacją stanowi skuteczne narzędzie intensyfikacji bioremediacji gleby z dołu urobkowego,
- analizy respirometryczne i chromatograficzne prowadzone w skali laboratoryjnej wykazały, że efektywność bioremediacji zanieczyszczeń ropopochodnych była uzależniona od dawki zastosowanych nanomateriałów. Stwierdzono, że przekroczenie optymalnego stężenia nanomateriałów w glebie prowadziło do ograniczenia aktywności mikroorganizmów, co podkreśla konieczność precyzyjnego projektowania tego typu strategii bioremediacyjnych. W przeprowadzonych badaniach najlepsze wyniki oczyszczania gleby z dołu

urobkowego uzyskano dla dawki nanomateriałów wynoszącej 20 mg/kg s.m. gleby,

- badania z wykorzystaniem dwóch rodzajów nanomateriałów potwierdziły ich skuteczność w intensyfikacji procesów biodegradacji, przy czym nieznacznie lepsze wyniki uzyskano dla nanotlenku grafenu (GO-NP),
- analizy chromatograficzne wykonane z zastosowaniem autorskich procedur badawczych umożliwiły identyfikację oraz ilościowe oznaczenie poszczególnych węglowodorów obecnych w glebie, co pozwoliło na ocenę ich podatności na biodegradację w układach wzbogaconych w nanomateriały i biopreparat,
- włączenie fitoremediacji z wykorzystaniem *Lolium perenne* do układów obejmujących bioaugmentację i nanomateriały pozwoliło na uzyskanie wyższej efektywności procesu oczyszczania gleby z dołu urobkowego. Badania fitoremediacyjne prowadzono przy zastosowaniu optymalnych dawek nanomateriałów wytypowanych na podstawie wcześniejszych badań laboratoryjnych,
- analizy chromatograficzne wykazały poprawę skuteczności biodegradacji wszystkich oznaczonych węglowodorów ropopochodnych w glebach obsadzonych *Lolium perenne* i wzbogaconych w biopreparat oraz nanomateriały. W układach tych, po zakończeniu eksperymentu, stopień biodegradacji TPH wynosił 81,85% dla GO-NP oraz 80,39% dla ZnO-NP, natomiast dla WWA odpowiednio 73,19% i 70,66%,
- analizy morfologiczne materiału roślinnego oraz analiza biomasy wykazały, że obecność nanomateriałów sprzyjała lepszemu rozwojowi *Lolium perenne* oraz zwiększeniu produkcji biomasy. Stwierdzono wzrost suchej masy roślin o 28,63% przy zastosowaniu GO-NP oraz o 22,21% w przypadku ZnO-NP w porównaniu do układów bez dodatku nanomateriałów. W układach wzbogaconych w nanomateriały odnotowano również większą biomasę korzeni o 29,70–30,84% oraz pędów o 18,65–27,58%,
- analiza zawartości TPH i WWA w korzeniach i pędach *Lolium perenne* potwierdziła zdolność roślin do pobierania węglowodorów ropopochodnych w trakcie fitoremediacji wspomaganej bioaugmentacją, przy czym większe ich ilości oznaczono w korzeniach, co było zgodne z wynikami przedstawionymi w artykułach IV i V. Wyższą akumulację zanieczyszczeń obserwowano w wariantach wzbogaconych w nanomateriały, co wskazuje na ich rolę w zwiększaniu biodostępności zanieczyszczeń oraz wspomaganiu procesów fitoekstrakcji,
- weryfikację zaproponowanej strategii bioremediacyjnej przeprowadzono na podstawie oceny ekotoksykologicznej gleby po zakończeniu eksperymentu, która wykazała poprawę jej jakości we wszystkich wariantach badawczych. Spadek wartości współczynników toksyczności TU w zastosowanych testach potwierdził, że intensyfikacja biodegradacji bezpośrednio przekłada się na ograniczenie negatywnego oddziaływania zanieczyszczeń na organizmy żywe. Wykazano jednocześnie, że wzbogacenie gleby w nanomateriały w optymalnej dawce nie

powodowało pogorszenia jakości gleby ani wzrostu jej toksyczności, lecz przyczyniało się do jej obniżenia, co ma istotne znaczenie w kontekście ograniczonej wiedzy dotyczącej wpływu nanomateriałów na środowisko,

Wyniki badań przedstawione w artykule VI wskazują na nową możliwość modyfikacji istniejących metod bioremediacji, polegającą na łącznym zastosowaniu fitoremediacji, bioaugmentacji oraz nanomateriałów, co stanowi innowacyjne podejście do intensyfikacji procesu oczyszczania gleby z dołu urobkowego i wskazuje na wysoki potencjał aplikacyjny opracowanej strategii i jej możliwe wykorzystanie w przyszłości w praktyce terenowej.

8. Efektywności biodegradacji węglowodorów ropopochodnych przy współwystępowaniu skażenia polichlorowanymi bifenylami (PCB) – Artykuł VII


Analizując dostępne źródła literaturowe dotyczące bioremediacji gleb historycznie zanieczyszczonych w wyniku działalności górniczej, można znaleźć informacje, że obecność w glebie polichlorowanych bifenyli, może ograniczać funkcje bakterii degradujących węglowodory ropopochodne ze względu na ich niską biodostępność i toksyczny wpływ na mikrobiotę glebową. W związku z powyższym w artykule VII postanowiono doświadczalnie sprawdzić skuteczność techniki bioaugmentacji w poprawie efektywności biodegradacji węglowodorów ropopochodnych przy współwystępowaniu w glebie zanieczyszczeń z grupy PCB. W badaniach zastosowano zarówno gleby skażone wyłącznie węglowodorami ropopochodnymi lub PCB, jak i gleby zanieczyszczone mieszaniną tych zanieczyszczeń. Nowością przedstawionych badań było porównanie skuteczności biodegradacji w układach zawierających PCB z układami wolnymi od PCB oraz porównanie działania pojedynczych szczepów bakteryjnych z autorskim biopreparatem M1. Do eksperymentów wykorzystano trzy szczepy bakteryjne: dwa szczepy autochtoniczne wyizolowane z obszaru dołu urobkowego (*Mycolicibacterium frederiksbergense* IN53 i *Rhodococcus erythropolis* IN129) oraz jeden szczep (*Rhodococcus* sp. IN306) pochodzący z gleby pustynnej skażonej ropą naftową o potwierdzonych właściwościach degradacyjnych względem PCB. Na bazie tych szczepów opracowano biopreparat M1, który zastosowano w badaniach laboratoryjnych i półtechnicznych prowadzonych metodą pryzmowania *ex-situ*.

Wyniki analiz respirometrycznych, chromatograficznych i toksykologicznych zaprezentowanych w artykule VII stanowią rozwinięcie wcześniejszych badań, dostarczając nowych informacji na temat wpływu współwystępowania PCB na biodegradację węglowodorów ropopochodnych. Jednocześnie potwierdzają, że biopreparaty złożone z kilku szczepów bakteryjnych, jak mieszanina M1, zwiększają efektywność procesu oraz pozwalają na skuteczniejsze oczyszczanie gleb skażonych różnymi rodzajami zanieczyszczeń.

8.1. Omówienie wyników zaprezentowanych w artykule VII


W badaniach respirometrycznych przeprowadzonych na glebach A, B i C oceniano intensywność procesów biodegradacyjnych w warunkach sterylnych po inokulacji pojedynczymi szczepami bakteryjnymi oraz biopreparatem M1. W glebie A (zanieczyszczonej PCB), największe zużycie tlenu zaobserwowano w układzie inokulowanym szczepem *Rhodococcus* sp. IN306 (3 274 mg O_2/dm^3), natomiast w glebie B (zanieczyszczonej węglowodorami ropopochodnymi), w układzie inokulowanym szczepem *Mycolicibacterium frederiksbergense* IN53 (4 235 mg O_2/dm^3), co wskazuje na wysoki potencjał tych szczepów pod kątem możliwości wykorzystania w procesach bioremediacji gruntów zanieczyszczonych tego rodzaju ksenobiotykami. Nieznacznie mniejszą intensywność procesów biodegradacyjnych zaobserwowano w układach zaszczepionych biopreparatem M1, dla których oznaczona ilość zużycia tlenu po

zakończeniu eksperymentu wynosiła 3 161 mg O_2/dm^3 (gleba A) i 3 975 mg O_2/dm^3 (gleba B). Warto zauważyć, że dla gleby pochodzącej z terenu dołu urobkowego (gleba B) zdecydowanie najniższą aktywność biologiczną odnotowano w układzie inokulowanym szczepem *Rhodococcus* sp. IN306 (3 193 mg O_2/dm^3), co podkreśla przewagę mikroorganizmów autochtonicznych nad obcym szczepem w rozkładzie węglowodorów ropopochodnych. Ponadto analizy respirometryczne, wykazały zdecydowanie wyższy poziom zużycia tlenu w układach zawierających glebę C (zanieczyszczoną węglowodorami ropopochodnymi i PCB) niż w glebach A i B, co sugeruje, że proces biodegradacji przebiegał w nich intensywniej, a obecność PCB w niskich stężeniach, nie stanowiła czynnika ograniczającego przebieg biodegradacji węglowodorów ropopochodnych w glebie, lecz mogła być dodatkowym źródłem węgla i energii dla mikrobioty glebowej. Poza tym przeprowadzone badania w warunkach laboratoryjnych wskazały, że w przypadku gleb zanieczyszczonych szerokim spektrum zanieczyszczeń, wskazane jest zastosowanie biopreparatu zawierającego co najmniej kilka szczepów bakteryjnych charakteryzujących się zróżnicowaną zdolnością do degradacji poszczególnych grup ksenobiotyków, co potwierdza najwyższa wartość zużycia tlenu odnotowana w układzie z glebą C poddaną inokulacji biopreparatem M1 (7 136 mg O_2/dm^3) (Rys. 50).



Rysunek 50 Zależność ilości zużytego tlenu [mg/dm^3] od czasu w układach zawierających glebę kontrolną (bez zanieczyszczeń) i gleby zanieczyszczone: a) PCB, b) TPH c) PCB oraz TPH poddane inokulacji szczepami: Mycolicibacterium frederiksbergense IN53, Rhodococcus erythropolis IN129, Rhodococcus sp. IN306 oraz biopreparatem M1.

Uzupełnieniem badań laboratoryjnych były analizy chromatograficzne, które wykazały spadek stężenia PCB w glebie A po inokulacji szczepem *Rhodococcus* sp. IN306 o 54,4%, natomiast po bioaugmentacji biopreparatem M1 o 46,2%. W glebie B najwyższą redukcję stężenia węglowodorów ropopochodnych uzyskano po inokulacji szczepem *Mycolicibacterium frederiksbergense* IN53, która wyniosła 37,2% oraz po bioaugmentacji biopreparatem M1, gdzie osiągnęła 33,6%. Wyniki te w połączeniu z wynikami analiz respirometrycznych wyraźnie wskazują, że spośród testowanych mieszanin inokulujących biopreparat M1 umożliwiał uzyskanie zadowalających stopni biodegradacji zarówno PCB jak i węglowodorów ropopochodnych. Analizy chromatograficzne przeprowadzone na próbkach gleby C poddanej inokulacji szczepami *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306 lub biopreparatem M1, potwierdziły te obserwacje. Po zakończeniu eksperymentu stopnie biodegradacji PCB i TPH w układzie inokulowanym biopreparatem M1 wynosiły odpowiednio 51,8% i 34,6%, natomiast w układach inokulowanych pojedynczymi szczepami bakteryjnymi były wyraźnie niższe (Rys. 51). Wynikało to z obecności w biopreparacie M1 szczepów bakterii o różnym potencjale biodegradacyjnym w stosunku do zanieczyszczeń obecnych w glebie, co umożliwiło skuteczną degradację zarówno PCB, jak i TPH.




Rysunek 51 Zestawienie stopni biodegradacji wybranych grup a) PCB, b) TPH po inokulacji gleby C szczepami Mycolicibacterium frederiksbergense IN53, Rhodococcus erythropolis IN129, Rhodococcus sp. IN306 oraz biopreparatem M1.

Potencjał biopreparatu M1 pod kątem możliwości wykorzystania w zabiegu inokulacji gleb zanieczyszczonych węglowodorami ropopochodnymi przy współwystępowaniu PCB zweryfikowano w warunkach półtechnicznych. Do tego celu wykorzystano niesterylną glebę D, charakteryzującą się wyższym stopniem skażenia niż gleby wykorzystane w badaniach laboratoryjnych (TPH=22 126 mg/kg s.m., PCB=8 752 µg/kg s.m.). Badania biodegradacji zanieczyszczeń prowadzono metodą pryzmowania


ex-situ przez okres sześciu miesięcy, co skutkowało obniżeniem stężenia zanieczyszczeń z grupy TPH o 70,8% i PCB o 84,5%.

W trakcie procesu bioremediacji monitorowano zmianę stężenia poszczególnych węglowodorów ropopochodnych oraz kongenerów PCB obecnych w glebie za pomocą analiz chromatograficznych. Wykazano, że w glebie D najłatwiej biodegradacji ulegały węglowodory alifatyczne o długości łańcucha węglowego n-C_{11}–n-C_{22} (87,2–93,2%), natomiast n-alkany zawierające powyżej 22 atomów węgla w cząsteczce (n-C_{23}–n-C_{40}) charakteryzowały się wyraźnie większą opornością na rozkład biologiczny (25,6-72,4%), co było zgodne z opisywanymi wcześniej zależnościami dotyczącymi wpływu długości łańcucha węglowego na biodegradację węglowodorów alifatycznych (Rys 52a). Ponadto wskazano, że spośród oznaczonych w glebie kongenerów PCB najłatwiej biodegradacji ulegały PCB28, PCB52 i PCB77, dla których stopień rozkładu po sześciu miesiącach oczyszczania wynosił od 87,7 do 94,6%, natomiast najtrudniej biodegradowalny okazał się kongener PCB180, dla którego uzyskano 58,6% biodegradacji (Rys. 52b). Uzyskane wyniki przeprowadzonych analiz chromatograficznych wskazują, że podatność na biodegradację zanieczyszczeń z grupy PCB zależała od liczby atomów chloru oraz obecności struktury *di-ortho* w cząsteczce.



Rysunek 52 Zestawienie stężeń a) kongenerów PCB, b) n-alkanów oznaczonych w niesterylnej glebie D poddanej inokulacji biopreparatem M1.

Na podstawie wyników analiz chromatograficznych obliczono stałe szybkości biodegradacji TPH oraz PCB w glebie D6, które wynosiły k = 0,0070 d^{-1} dla TPH i k = 0,0109 d^{-1} dla PCB a także dla poszczególnych frakcji: TriCBs (0,0170 d^{-1}), TetraCBs (0,0110 d^{-1}), PentaCBs (0,0095 d^{-1}), HexaCBs (0,0082 d^{-1}) i HeptaCBs (0,0050 d^{-1}) oraz n-alkanów $\Sigma nC_{11}–nC_{22}$ (0,0130 d^{-1}) i $\Sigma nC_{23}–nC_{40}$ (0,0049 d^{-1}). Uzyskane wartości stałych biodegradacji potwierdziły wysoką skuteczność biopreparatu M1 zarówno w zakresie biodegradacji węglowodorów ropopochodnych, jak i polichlorowanych bifenyli, a także umożliwiły porównanie tempa rozkładu poszczególnych grup zanieczyszczeń oznaczonych w oczyszczanej glebie (Rys. 53).




Rysunek 53 Porównanie obniżenia zawartości wybranych grup zanieczyszczeń a) PCB po normalizacji PCB209, b)TPH po normalizacji hopanem w kolejnych miesiącach oczyszczania gleby z dołu urobkowego techniką inokulacji biopreparatem M1.

W artykule VII standardowy pakiet testów toksykologicznych (Phytotoxkit™, Ostracodtoxkit F™ oraz Microtox SPT) wykorzystywanych do oceny ekotoksykologicznej gleby w trakcie oraz po zakończeniu oczyszczania rozszerzono o testy MARA oraz Ames. Było to spowodowane niewielką dostępnością danych literaturowych na temat możliwości powstawania toksycznych metabolitów pośrednich w trakcie oczyszczenia gleb zanieczyszczonych węglowodorami ropopochodnymi przy współwystępowaniu skażenia PCB. Szczególnie problematyczna pod tym względem mogła okazać się obecność w glebie PCB, które są zaliczane do związków wysoce toksycznych oraz wykazujących potencjał genotoksyczny, co w przypadku braku przeprowadzenia rozszerzonych testów toksykologicznych, mogło zwiększać niepewność oceny rzeczywistego poziomu ryzyka ekotoksykologicznego w glebie po bioremediacji.


Wyniki wszystkich testów toksykologicznych omówionych w artykule VII wykazały spadek toksyczności gleby D poddanej bioaugmentacji biopreparatem M1, czego potwierdzeniem były obniżone wartości współczynników toksyczności TU w stosunku do gleby wyjściowej. W teście Phytotoxkit™ uzyskano obniżenie wartości TU z

zakresu 8,5–10,5 do poziomu 3,0–3,1 w parametrze kiełkowania oraz z 10,1–15,0 do 3,6–4,6 w parametrze zahamowania wzrostu korzeni *Lepidium sativum, Sorghum saccharatum* oraz *Sinapis alba.* W pozostałych testach wartości współczynnika TU spadły z 16,9 na 4,9 (test Ostracodtoxkit F™), z 15,4 na 3,9 (test Microtox SPT) oraz z 19,6 na 5,2 (test MARA) (Rys.54). Wyniki przeprowadzonych testów Phytotoxkit™, Ostracodtoxkit F™, Microtox SPT oraz MARA wyraźnie wskazują, że inokulacja gleby D biopreparatem opracowanym na bazie szczepów *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129 oraz *Rhodococcus* sp. IN306 umożliwiła nie tylko redukcję zawartości zanieczyszczeń w oczyszczanej glebie, ale także wpłynęła korzystnie na poprawę jej właściwości ekotoksykologicznych..



Rysunek 54 Zestawienie wyników testów toksykologicznych gleby z dołu urobkowego zanieczyszczonej TPH i PCB poddanej inokulacji biopreparatem M1.

Wyniki testu Amesa (test mutagenności Amesa z użyciem bakterii *Salmonella Typhimurium*), przeprowadzonego w celu wykrycia w badanej glebie potencjalnych związków mutagennych oraz kancerogennych, wykazały że w wyniku inokulacji gleby D biopreparatem M1 po 6 miesiącach bioremediacji doszło do obniżenia wartości współczynnika mutagenności z 7,85–9,85 do poziomu 0,68–1,86. Wyniki te wskazują na istotne zmniejszenie potencjału mutagennego badanej gleby w wyniku przeprowadzonych zabiegów bioremediacyjnych oraz brak powstawania metabolitów pośrednich o charakterze mutagennym (Rys. 55).



Rysunek 55 Wpływ stężenia PCB i TPH na liczbę rewertantów w glebie a) wyjściowej (Soil D), b) po 3 miesiącach biodegradacji (Soil D3), c) po 6 miesiącach biodegradacji (Soil D6).

8.2. Podsumowanie artykułu VII

W artykule VII zaprezentowano wyniki badań biodegradacji dostarczające istotnych informacji na temat wpływu zanieczyszczeń współwystępujących, takich jak PCB, na przebieg biodegradacji węglowodorów ropopochodnych w glebie. Na podstawie przeprowadzonych analizy respirometrycznych, chromatograficznych oraz toksykologicznych sformułowano najważniejsze wnioski wynikające z artykułu VII:

- wyniki analiz respirometrycznych i chromatograficznych wykazały, że zastosowane w badaniach szczepy bakteryjne charakteryzowały się zróżnicowaną skutecznością w biodegradacji badanych zanieczyszczeń organicznych. W przypadku PCB najwyższą efektywność biodegradacji uzyskano dla inokulacji gleby szczepem *Rhodococcus* sp. IN306, natomiast w przypadku węglowodorów ropopochodnych najskuteczniejszy okazał się szczep *Mycolicibacterium frederiksbergense* IN53. W warunkach współwystępowania w glebie węglowodorów ropopochodnych i PCB najwyższą efektywność biodegradacji obu grup zanieczyszczeń uzyskano dla biopreparatu M1,
- wyniki analiz laboratoryjnych potwierdziły, że w celu skutecznego oczyszczania gleb zanieczyszczonych substancjami należącymi do różnych grup związków chemicznych (węglowodory alifatyczne, PCB) z wykorzystaniem techniki bioaugmentacji, wskazane jest stosowanie biopreparatów zawierających kilka szczepów bakteryjnych o różnym spektrum działania,
- spośród analizowanych mieszanin inokulujących największym potencjałem w zakresie wspomagania biodegradacji węglowodorów ropopochodnych w glebie współzanieczyszczonej PCB charakteryzował się biopreparat B1, który wytypowano do dalszych badań porównawczych prowadzonych w skali półtechnicznej,
- badania biodegradacji prowadzone metodą pryzmowania *ex-situ* w skali półtechnicznej potwierdziły skuteczność przyjętej strategii oczyszczania opartej na inokulacji biopreparatem M1, opracowanym na bazie szczepów bakterii autochtonicznych z dodatkiem szczepu nieautochtonicznego,

- monitoring procesu oczyszczania z wykorzystaniem analiz chromatograficznych umożliwił kompleksową ocenę przebiegu biodegradacji, w tym określenie sumarycznego stopnia rozkładu zanieczyszczeń, identyfikację poszczególnych n-alkanów oraz wybranych kongenerów PCB wraz z oceną stopnia ich degradacji, wyznaczenie wskaźników biodegradacji, takich jak stosunki n-C_{17}/Pr oraz n-C_{18}/F stanowiących miarę efektywności procesu, a także obliczenie stałych szybkości biodegradacji wybranych grup zanieczyszczeń,
- przeprowadzenie monitoringu toksykologicznego z wykorzystaniem zestawu testów obejmujących Phytotoxkit™, Ostracodtoxkit F™, Microtox SPT, MARA oraz Ames umożliwiło ocenę zmian toksyczności gleby w trakcie procesu oczyszczania i potwierdziło skuteczność zastosowanej inokulacji biopreparatem M1 w poprawie jej własności użytkowych. Przeprowadzenie badań toksykologicznych było szczególnie istotne z uwagi na wciąż niewystarczającą ilość danych dotyczących możliwości powstawania produktów toksycznych podczas degradacji węglowodorów ropopochodnych przy współwystępowaniu PCB.

Wyniki badań przedstawione w artykule VII przyczyniły się do realizacji założonych celów niniejszej rozprawy doktorskiej oraz stanowiły istotne uzupełnienie wiedzy dotyczącej biodegradacji węglowodorów ropopochodnych w warunkach współwystępowania PCB. Problematyka ta jest stosunkowo rzadko podejmowana, co wynika z faktu, że związki te nie są obecnie produkowane, a ich obecność w środowisku związana jest głównie z historycznym skażeniem gleb powstałym m.in. w wyniku wycieków zużytych olejów transformatorowych. Przeprowadzone badania, zaprezentowane w artykule VII pozwoliły na wypełnienie tej luki badawczej w zakresie bioremediacji gleb zanieczyszczonych wieloskładnikowo.

Kompleksowy charakter badań zaprezentowanych w artykule VII, obejmujący analizy respirometryczne, chromatograficzne oraz ocenę toksykologiczną, umożliwił wieloaspektową ocenę skuteczności zaproponowanej strategii oczyszczania gleby z dołu urobkowego zanieczyszczonej węglowodorami ropopochodnymi przy współwystępowaniu PCB, co zwiększa możliwości zastosowania opracowanej metody w warunkach rzeczywistych

9. Podsumowanie i wnioski

Historyczne zanieczyszczenie gruntów substancjami ropopochodnymi, w szczególności na terenach związanych z dawną działalnością górnictwa nafty i gazu, stanowi istotny problem ekologiczny. Obecność tych zanieczyszczeń w glebach stwarza realne zagrożenie dla zdrowia ludzi i zwierząt, co uzasadnia konieczność prowadzenia skutecznych działań rekultywacyjnych ukierunkowanych na ich usunięcie oraz przywrócenie wartości użytkowych zdegradowanych terenów. W Instytucie Nafty i Gazu – Państwowym Instytucie badawczym od lat prowadzone są badania na temat możliwości wykorzystania procesów bioremediacyjnych do oczyszczania gruntów na obszarach dawnych składowisk odpadów wiertniczych tzw. dołów urobkowych. Pomimo wysokiej skuteczności opracowanej w INiG-PIB (w Zakładzie Technologii Eksploatacji Płynów Złożowych i w Zakładzie Mikrobiologii) technologii etapowego oczyszczania odpadów wiertniczych zanieczyszczonych zestarzałymi związkami ropopochodnymi zdeponowanych w starych dołach urobkowych, w przypadku gleb z terenów kopalni Grabownica charakteryzujących się bardzo wysokim stężeniem zanieczyszczeń ropopochodnych, po trzech latach badań nie udało się osiągniąć zakładanego stopnia oczyszczania. Zasadne zatem było opracowanie bardziej zaawansowanych technik bioremediacji umożliwiających dalsze doczyszczanie gleb z terenów dołów urobkowych.

Postawiono hipotezę, że *głębsze doczyszczenie gruntów z dołów urobkowych można będzie osiągnąć poprzez modyfikację istniejących technik bioremediacyjnych i opracowanie zintegrowanych strategii zwiększających efektywność biodegradacji węglowodorów ropopochodnych*.

Realizacja niniejszej pracy doktorskiej umożliwiła weryfikację postawionej hipotezy badawczej i osiągnięcie wszystkich założonych celów przedstawionych w rozdziale 1.6.

W pracy zaprezentowano i poddano analizie różnorodne podejścia ukierunkowane na zwiększenie efektywności procesu biodegradacji zanieczyszczeń ropopochodnych w glebach pochodzących z terenów dołów urobkowych. Szczególną wartość naukową stanowiło interdyscyplinarne podejście do badanego zagadnienia, integrujące problematykę z zakresu wiertnictwa, inżynierii środowiska, analityki chemicznej, mikrobiologii oraz ekotoksykologii. Tak szeroka perspektywa badawcza umożliwiła kompleksowe ujęcie problemu oraz wieloaspektową ocenę skuteczności zaproponowanych strategii bioremediacji, a także sformułowanie najważniejszych osiągnięć badawczych:

- Opracowanie skutecznych i przyjaznych dla środowiska metod doczyszczania gleb zanieczyszczonych węglowodorami ropopochodnymi, wymagało przeprowadzenia kompleksowych badań obejmujących: analizy fizyczno-chemiczne, respirometryczne, chromatograficzne, mikrobiologiczne i ekotoksykologiczne. Należy podkreślić, że badania prowadzono na rzeczywistych próbkach gleb pochodzących z terenów dołów urobkowych po trzyletnim procesie oczyszczania, jako najbardziej reprezentatywnych dla realizacji założonych celów.

- Analizy chromatograficzne prowadzone z wykorzystaniem opracowanych i zmodyfikowanych procedur analitycznych umożliwiły:
    - ✓ *identyfikację oraz ilościowe oznaczenie poszczególnych węglowodorów wchodzących w skład TPH, BTEX, WWA i PCB;.*
    - ✓ *określenie zmian stężeń analizowanych zanieczyszczeń w wyniku zastosowanych strategii bioremediacyjnych;*
    - ✓ *identyfikację węglowodorów z grupy izoprenoidów, takich jak pristan i fitan, należących do związków trudno biodegradowalnych, co umożliwiło określenie stopnia biodegradacji n-alkanów na podstawie zmian wartości wskaźników n-C_{17}/Pr oraz n-C_{18}/F;*
    - ✓ *obliczenie stopni biodegradacji węglowodorów alifatycznych i aromatycznych, co było podstawą oceny skuteczności zaproponowanych strategii bioremediacyjnych*.

- Przeprowadzone badania toksykologiczne z wykorzystaniem biotestów o zróżnicowanej wrażliwości Phytotoxkit™, Ostracodtoxkit F™, Microtox®SPT, MARA i Ames wskazały, że w wyniku zaproponowanych strategii bioremediacyjnych uzyskano zmniejszenie toksyczności gleby, które korelowało ze spadkiem zawartości zanieczyszczeń ropopochodnych w oczyszczonej glebie.

- Badania prowadzone w warunkach laboratoryjnych i półtechnicznych z wykorzystaniem analiz respirometrycznych, chromatograficznych i toksykologicznych wykazały, że modyfikacja procesu inokulacji gleby biopreparatem opracowanym na bazie bakterii autochtonicznych, poprzez dodatek γ-PGA, spowodowała poprawę efektywności biodegradacji zanieczyszczeń ropopochodnych w badanej glebie o 13,18% dla TPH, 10,11% dla BTEX i 12,32% dla WWA w stosunku do gleby wyłącznie inokulowanej biopreparatem. Uzyskane wyniki wykazały, że γ-PGA wpłynął na poprawę biodostępności zanieczyszczeń ropopochodnych, co skutkowało większą efektywnością oczyszczania badanej gleby.

- Modyfikacja procesu inokulacji gleby biopreparatem opracowanym na bazie bakterii autochtonicznych poprzez zastosowanie biosurfaktantów pozwoliła na poprawę efektywności biodegradacji zarówno węglowodorów alifatycznych jak i wielopierścieniowych węglowodorów aromatycznych w glebie z terenu dołu urobkowego. Spośród przebadanych biosurfaktantów największym potencjałem pod kątem poprawy efektywności biodegradacji TPH i WWA wspomaganej bioaugmentacją wykazała się autorska mieszanina PSR opracowana na bazie trzech różnych związków powierzchniowo-czynnych tj. γ-PGA, surfaktyny i ramnolipidów. Pozostałe biosurfaktanty wykorzystane w badaniach również istotnie wpłynęły na poprawę skuteczności oczyszczania gleby z dołu urobkowego, a ich skuteczność malała w kolejności JBR 425 > JBR 320 > γ-PGA >ramnolipidy>surfaktyna, co wskazuje, że biosurfaktanty mogą zwiększać biodostępność zanieczyszczeń w trakcie procesów bioremediacyjnych. Zastosowanie najbardziej optymalnego wariantu wzbogacenia gleby tj. inokulacji

biopreparatem w połączeniu z mieszaniną PSR wpłynęło na poprawę biodegradacji TPH i WWA w stosunku do gleby wyłącznie bioaugmentowanej o odpowiednio 32,97% i 33,35%.

- Wzbogacenie biopreparatu na bazie bakterii autochtonicznych w wybrane szczepy niepatogennych grzybów autochtonicznych spowodowała poprawę efektywności biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego o 9,82% dla TPH i 7,32% dla WWA. Obecność grzybów w biopreparacie przyczyniła się do zwiększenia efektywności degradacji wszystkich analizowanych węglowodorów ropopochodnych obecnych w badanej glebie, przy czym szczególnie wyraźną poprawę odnotowano w przypadku węglowodorów alifatycznych o dłuższych łańcuchach węglowych oraz WWA, należących do związków trudno biodegradowalnych.

- W celu głębszego doczyszczenia gleb zanieczyszczonych węglowodorami ropopochodnymi, pochodzących z terenów dołów urobkowych przeprowadzono badania fitoremediacji wykorzystaniem jako fitoremedianta roślin uprawnych (*Zea mays)* oraz ozdobnych (*Echinacea purpurea*) wspomaganej inokulacją biopreparatem na bazie autochtonicznych bakterii i grzybów. Proces fitoremadiacji gleby z terenu dołu urobkowego z wykorzystaniem jako fitoremedianta *Zea mays* umożliwił obniżenie zawartości zanieczyszczeń ropopochodnych (TPH i WWA) odpowiednio 22,81% i 14,48% natomiast jej połączenie z inokulacją przyczyniło się do wzrostu efektywności oczyszczania gleby (TPH o 26,99% i WWA o 22,26%). Zbieżne wyniki uzyskano podczas fitoremediacji z wykorzystaniem *Echinacea purpurea* wspomaganej inokulacją na bazie autochtonicznych bakterii i grzybów. Zastosowanie w procesach fitoremediacyjnych zmodyfikowanej inokulacji biopreparatem na bazie bakterii i grzybów poprzez dodatek γ-PGA skutkowało dalszym wzrostem stopnia biodegradacji TPH i WWA, co potwierdziło skuteczność γ-PGA w poprawie biodostępności zanieczyszczeń ropopochodnych. Uzyskane wyniki badań pozwalają na stwierdzenie, że zastosowanie strategii łączącej dwie techniki tj. fitoremediację z inokulacją pozwala na uzyskanie większej efektywności oczyszczania gleb z terenów dołów urobkowych.

- Wykorzystanie jako fitoremedianta dziko rosnących roślin (*Scirpus sylvaticus* oraz *Cirsium oleraceum*) pobranych z obszaru naturalnego wycieku ropy naftowej znajdującego się w pobliżu dołu urobkowego podczas sześciomiesięcznego procesu fitoremediacji wspomaganej inokulacją konsorcjum bakteryjnym na bazie bakterii autochtonicznych uzyskano w glebie z dołu urobkowego obniżenie stężenia zastarzałych zanieczyszczeń ropopochodnych: TPH z 4,320 do 455 – 261 mg kg^{-1} m. s. (89,1 – 93.69%) oraz WWA z 7,19 mg kg^{-1} m. s do 1,14 – 1,53 mg kg^{-1} m. s. (75,2 – 79,5%). Porównując uzyskane wyniki procesu fitoremediacji z wykorzystaniem dzikich roślin wspomaganego inokulacją konsorcjum bakteryjnym z wcześniejszymi badaniami procesów fitoremediacyjnych z wykorzystaniem roślin uprawnych i ozdobnych takich jak *Zea mays,* czy

Echinacea purpurea, można stwierdzić, że wprowadzenie dziko rosnących roślin z wycieku naturalnego ropy naftowej przyczyniło się osiągnięcia wyższego stopnia redukcji zanieczyszczeń ropopochodnych (TPH i WWA), który korelował z niższym stopniem toksyczności oczyszczanej gleby. Uzyskane wyniki wskazują, że zastosowanie jako fitoremedianta dziko rosnących roślin należących do gatunków *(Scirpus sylvaticus Cirsium oleraceum)* pochodzących z naturalnego wycieku ropy naftowej są one odpowiednimi kandydatami, gdyż są w stanie wytrzymać skażenie i uczestniczyć w naturalnym procesie bioatenuacji i mogą zwiększyć potencjał biodegradacyjny oczyszczanej gleby poprzez wzbogacenie go o taksony wchodzące w skład mikrobiomów zamieszkujących ryzosferę obu gatunków roślin.

- Przeprowadzone badania obejmujące analizę biomasy oraz tempa wzrostu roślin wykazały, że wzbogacenie procesu fitoremediacji prowadzonej z wykorzystaniem dziko rosnacych roślin (*Scirpus sylvaticus* oraz *Cirsium oleraceum*) poprzez inokulację biopreparatem opracowanym na bazie bakterii autochtonicznych wpłynęła na lepszy wzrost roślin, co w konsekwencji skutkowało zwiększeniem efektywności bioremediacji. Ponadto przeprowadzone analizy zawartości zanieczyszczeń ropopochodnych (TPH i WWA) w korzeniach i pędach roślin fitoremediujących potwierdziły, że inokulacja może istotnie wpływać na poprawę fitoekstrakcji zanieczyszczeń z gleby do tkanek roślinnych, w których mogą ulegać akumulacji, transformacji lub degradacji. Ponadto, oznaczono nieznacznie wyższe zawartości węglowodorów ropopochodnych w materiale roślinnym *Cirsium oleraceum,* co wynika prawdopodobnie ze zdecydowanie lepiej rozwiniętego systemu korzeniowego, pozwalającego na usprawnienie mechanizmów i oddziaływań zachodzących w układzie roślina-mikrobiota glebowa.

- Spadek toksyczności w próbkach po przeprowadzonym procesie fitoremediacji w wykorzystaniem dzikich roślin wspomaganych inokulacją konsorcjum bakteryjnym koreluje ze spadkiem zawartości zanieczyszczeń ropopochodnych w glebie i wzrostem biomasy roślin korzeni i pędów roślin. W badaniach fitotoksyczności z wykorzystaniem testu Phytotoxkit™ odnotowano zmienną wrażliwość badanych roślin na zanieczyszczenia występujące w badanych glebach, w kolejności: *Lepidium sativum* < *Sorghum saccharatum* < *Sinapis alba*. Natomiast czułość pozostałych biotestów była porównywalna i zwiększana w kolejności: Ostracodtoxkit F™< Microtox SPT< MARA .

- Ocena struktury społeczności bakteryjnej przy zastosowania jsko fitoremedianta roślin dziko rosnacych, wykazała wyraźny spadek (wyrażony względną liczebnością) typowych mikroorganizmów obligatoryjnie beztlenowych, takich jak archeony metanogenne i przedstawiciele *Desulfobacterota* we wszystkich remediowanych glebach. Zmniejszona liczebność niektórych taksonów bakterii tlenowych i fakultatywnie beztlenowych może sugerować, że zostały one wyparte przez inne mikroorganizmy lub narażone na wzmożone żerowanie pierwotniaków w zrekultywowanych glebach. Zrekultywowane gleby zawierały przedstawicieli

społeczności mikrobiologicznej, których nie było w glebie zanieczyszczonej. Zatem wszystkie te taksony były prawdopodobnie częścią mikrobiomów zasiedlających ryzosferę obu gatunków roślin. Wykazano również, że bioaugmentacja gleby zanieczyszczonej węglowodorami skutkowała wzrostem liczebności *Mycobacterium/Mycolibacterium*, co sugeruje, że jest ono w stanie przetrwać w zremediowanych glebach i ich obecność w ryzosferze przyspiesza degradację zanieczyszczeń organicznych i zwiększa odporność roślin na zanieczyszczenia obecne w glebie, zwiększając tym samym tolerancję roślin na niekorzystne warunki środowiskowe.

- W badaniach wprowadzono nowy element badawczy dotyczący oceny zmian w zbiorowisku pierwotniaków glebowych (orzęsków), które przyczyniają się one do obiegu składników odżywczych w ekosystemie poprzez zjadanie bakterii i przyspieszanie obrotu biomasą bakteryjną i materią organiczną gleby, uwalniając w ten sposób zamaskowane składniki odżywcze z biomasy drobnoustrojów niedostępne dla producentów pierwotnych, co czyni je bioindykatorami stresu środowiskowego. Przeprowadzenie procesu fitoremediacji z wykorzystaniem dziko rosnących roślin wspomaganej inokulacją biopreparatem na bazie autochtonicznych bakterii przyczyniło się do wzrostu liczebności orzęsków. Obliczone wskaźniki ekologicznej różnorodności biologicznej są cennym elementem charakterystyki procesów fitoremediacyjnych z wykorzystaniem dzikich roślin należących do gatunku (*Scirpus sylvaticus* i *Cirsium oleraceum).*

- Przeprowadzone badania skuteczności biodegradacji zastarzałych zanieczyszczeń ropopochodnych w glebie z terenu dołu urobkowego pod kątem wykorzystania nanocząstek tlenku grafenu (GO-NP) i tlenku cynku (ZnO-NP) w procesie fitoremediacji z wykorzystaniem jako fitoremedianta *Lolium perenne* w połączeniu z inokulacją konsorcjum bakteryjnym na bazie bakterii autochtonicznych oraz wykazały, że ich obecność może istotny wpływ na poprawę efektywności biodegradacji zarówno węglowodorów alifatycznych jak i wielopierścieniowych węglowodorów aromatycznych w oczyszczanej glebie. Spośród testowanych nanomateriałów najwyższym potencjałem pod kątem poprawy efektywności biodegradacji zanieczyszczeń ropopochodnych w glebie inokulowanej biopreparatem i obsadzonej *Lolium perenne* charakteryzował się tlenek grafenu, którego użycie skutkowało wzrostem redukcji TPH do 81,85% i WWA do 73,19%, co przekładało się na poprawę efektywności biodegradacji tych zanieczyszczeń o odpowiednio 18,63% i 11,33%. Nieznacznie niższe efekty uzyskano przy wykorzystaniu nanocząsteczek tlenku cynku.

- Przeprowadzone badania w skali laboratoryjnej i półtechnicznej (metoda pryzmowania *ex-situ*) pozwoliły na ocenę możliwości zastosowania do biodegradacji zanieczyszczeń ropopochodnych (TPH i WWA) przy współwystępowaniu skażenia PCB w glebie z dołów urobkowych szczepów bakteryjnych: *Mycolicibacterium frederiksbergense IN53, Rhodococcus erythropolis IN129, Rhodococcus sp. IN306* oraz biopreparatu stworzonego na

bazie tych szczepów. Wyniki zarówno testów respirometrycznych, jak i analiz chromatograficznych wykazały, że wybrane szczepy bakteryjne o potencjalnej zdolności rozkładu PCB są rzeczywiście zdolne do usuwania tego typu zanieczyszczeń z gleby, a także są zdolne do rozkładu węglowodorów ropopochodnych (w szczególności alifatycznych). Natomiast, biopreparat opracowany na bazie testowanych szczepów bakteryjnych, charakteryzował się zadowalającą zdolnością biodegradacji zarówno dla poszczególnych kongenerów PCB, jak i TPH co rozszerza jego spektrum działania wpływając na efektywność biodegradacji zarówno węglowodorów ropopochodnych jak i polichlorowanych bifenyli.

- Kompleksowa ocena zaproponowanych strategii poprawy efektywności biodegradacji zanieczyszczeń ropopochodnych w glebach pobranych z terenów dołów urobkowych stanowi podstawę do możliwości wdrożenia ich w przyszłości w warunkach terenowych.

Bibliografia

1. Abbasian F., Lockington R., Mallavarapu M., Naidu R.,. A Comprehensive Review of Aliphatic Hydrocarbon Biodegradation by Bacteria. *Applied Biochemistry and Biotechnology* **2015**, 176, 670–699. https://doi.org/10.1007/s12010-015-1603-5
2. Abdel-Shafy H.I., Mansour M.S.M. A review on polycyclic aromatic hydrocarbons: Source, environmental impact, effect on human health and remediation. *Egyptian Journal of Petroleum* **2016**, 25, 107–123. https://doi.org/10.1016/j.ejpe.2015.03.011
3. Alazaiza M.Y.D., Alzghoul T.M., Ramu M.B., Nassani D.E. Catalytic Nanomaterials for Soil and Groundwater Remediation: Global Research Trends (2010–2024). *Catalysts* **2025**, 15, 981. https://doi.org/10.3390/catal15100981
4. Al-Dossary M.A., Abood S.A., Al-Saad H.T. Factors affecting polycyclic aromatic hydrocarbon biodegradation by *Aspergillus flavus*. *Remediation Journal* **2020**, 30, 17–25. https://doi.org/10.1002/rem.21658
5. Al-Khayri J.M., Rashmi R., Surya Ulhas R., Sudheer W.N., Banadka A., Nagella P., Aldaej M.I., Rezk A.A.-S., Shehata W.F., Almaghasla M.I. The Role of Nanoparticles in Response of Plants to Abiotic Stress at Physiological, Biochemical, and Molecular Levels. *Plants* **2023**, 12, 292. https://doi.org/10.3390/plants12020292
6. Arghiropol D., Rusu T., Moldovan M., Paltinean G.-A., Silaghi-Dumitrescu L., Sarosi C., Petean I. Petroleum Hydrocarbon Pollution and Sustainable Uses of Indigene Absorbents for Spill Removal from the Environment—A Review. *Sustainability* **2025**, 17, 8018. https://doi.org/10.3390/su17178018
7. Atlas R.M. Microbial degradation of petroleum hydrocarbons: an environmental perspective. *Microbiological Reviews* **1981**, 45, 180–209. https://doi.org/10.1128/mr.45.1.180-209.1981
8. Atlas R.M., Bartha R. Microbial ecology: fundamentals and applications, 4th ed, Benjamin Cummings series in the life sciences. Benjamin-Cummings Publishing Company, Subs of Addison Wesley Longman, Inc, New York **1998**.
9. Azeez N.A., Dash S.S., Gummadi S.N., Deepa V.S. Nano-remediation of toxic heavy metal contamination: Hexavalent chromium [Cr(VI)]. *Chemosphere* **2021** 266, 129204. https://doi.org/10.1016/j.chemosphere.2020.129204
10. Badmus S.O., Amusa H.K., Oyehan T.A., Saleh T.A. Environmental risks and toxicity of surfactants: overview of analysis, assessment, and remediation techniques. *Environmental Science and Pollution Research* **2021**, 28, 62085–62104. https://doi.org/10.1007/s11356-021-16483-w
11. Baker A.J.M., Brooks R. Terrestrial higher plants which hyperaccumulate metallic elements. A review of their distribution, ecology and phytochemistry. *Biorecovery* **1989**, 1(2), 81-126.
12. Baltacı N.G., Bayındırlı K.N., Baltacı M.Ö. *Alcanivorax*: Unique Genus of Hydrocarbon-Degrading Bacteria. *Biotech and App Biochem* **2026**, 73, 862–870. https://doi.org/10.1002/bab.70048

13. Banks M.K., Kulakow P., Schwab A.P., Chen Z., Rathbone, K. Degradation of Crude Oil in the Rhizosphere of *Sorghum bicolor*. *International Journal of Phytoremediation* **2003**, 5, 225–234. https://doi.org/10.1080/713779222
14. Baoune H., Aparicio J.D., Acuña A., El Hadj-khelil A.O., Sanchez L., Polti M.A., Alvarez A.,. Effectiveness of the *Zea mays*-*Streptomyces* association for the phytoremediation of petroleum hydrocarbons impacted soils. *Ecotoxicology and Environmental Safety* **2019a**, 184, 109591. https://doi.org/10.1016/j.ecoenv.2019.109591
15. Baoune H., Aparicio J.D., Pucci G., Ould El Hadj-Khelil A., Polti M.A. Bioremediation of petroleum-contaminated soils using *Streptomyces* sp. Hlh1. *Journal of Soils and Sediments* **2019b**, 19, 2222–2230. https://doi.org/10.1007/s11368-019-02259-w
16. Baragaño D., Alonso J., Gallego J.R., Lobo M.C., Gil-Díaz M. Magnetite nanoparticles for the remediation of soils co-contaminated with As and PAHs. *Chemical Engineering Journal* **2020**, 399, 125809. https://doi.org/10.1016/j.cej.2020.125809
17. Baran A., Tarnawski M. Phytotoxkit/Phytotestkit and Microtox® as tools for toxicity assessment of sediments. *Ecotoxicology and Environmental Safety* **2013**, 98, 19–27. https://doi.org/10.1016/j.ecoenv.2013.10.010
18. Barathi S., Vasudevan N. Utilization of petroleum hydrocarbons by Pseudomonas fluorescens isolated from a petroleum-contaminated soil. *Environment International* **2001**, 26, 413–416. https://doi.org/10.1016/S0160-4120(01)00021-6
19. Bhanu Revathi K., Meghana G., Anuradha S., Shinomol George K. Understanding mechanisms underlying genes regulating the production of biosurfactant, in: Green Sustainable Process for Chemical and Environmental Engineering and Science. Elsevier **2022**, 649–663. https://doi.org/10.1016/B978-0-323-85146-6.00035-8
20. Bjerk T.R., Severino P., Jain S., Marques C., Silva A.M., Pashirova T., Souto E.B. Biosurfactants: Properties and Applications in Drug Delivery, Biotechnology and Ecotoxicology. *Bioengineering* **2021**, 8, 115. https://doi.org/10.3390/bioengineering8080115
21. Blinova I., Niskanen J., Kajankari P., Kanarbik L., Käkinen A., Tenhu H., Penttinen O.-P., Kahru A. Toxicity of two types of silver nanoparticles to aquatic crustaceans *Daphnia magna* and *Thamnocephalus platyurus*. *Environmental Science and Pollution Research* **2013**, 20, 3456–3463. https://doi.org/10.1007/s11356-012-1290-5
22. Borowik A., Wyszkowska J., Gałązka A., Kucharski J. Role of *Festuca rubra* and *Festuca arundinacea* in determinig the functional and genetic diversity of microorganisms and of the enzymatic activity in the soil polluted with diesel oil. *Environmental Science and Pollution Research* **2019**, 26, 27738–27751. https://doi.org/10.1007/s11356-019-05888-3
23. Brakstad O.G., Bonaunet K. Biodegradation of Petroleum Hydrocarbons in Seawater at Low Temperatures (0–5 °C) and Bacterial Communities Associated with Degradation. *Biodegradation* **2006**, 17, 71–82.

https://doi.org/10.1007/s10532-005-3342-8

24. Brakstad O.G., Throne-Holst M., Netzer R., Stoeckel D.M., Atlas R.M. Microbial communities related to biodegradation of dispersed M acondo oil at low seawater temperature with Norwegian coastal seawater. *Microbial Biotechnology* **2015**, 8, 989–998. https://doi.org/10.1111/1751-7915.12303
25. Brown L.M., Gunasekera T.S., Striebich R.C., Ruiz O.N. Draft Genome Sequence of Gordonia sihwensis Strain 9, a Branched Alkane-Degrading Bacterium. *Genome Announcements* **2016**, 4, e00622-16. https://doi.org/10.1128/genomeA.00622-16
26. Brzeszcz J. Drobnoustroje środowiskowe, zdolne do jednoczesnego rozkładu węglowodorów alifatycznych i aromatycznych- perspektywy wykorzystania w praktyce bioremediacji gleb zanieczyszczonych substancjami ropopochodnymi **2017** (Praca doktorska).
27. Brzeszcz J., Kaszycki P. Aerobic bacteria degrading both n-alkanes and aromatic hydrocarbons: an undervalued strategy for metabolic diversity and flexibility. *Biodegradation* **2018**, 29, 359–407. https://doi.org/10.1007/s10532-018-9837-x
28. Brzeszcz J., Kapusta P., Turkiewicz A.,. Zastosowanie metod molekularnych w badaniach bioremediacji substancji ropopochodnych. *Nafta-Gaz* **2013**, 11, 829–842.
29. Brzeszcz J., Steliga T., Kapusta P., Turkiewicz A., Kaszycki P. r-strategist versus K-strategist for the application in bioremediation of hydrocarbon-contaminated soils. *International Biodeterioration & Biodegradation* **2016**, 106, 41–52. https://doi.org/10.1016/j.ibiod.2015.10.001
30. Brzeszcz J., Steliga T., Ryszka P., Kaszycki P., Kapusta P. Bacteria degrading both n-alkanes and aromatic hydrocarbons are prevalent in soils. Environmental *Science and Pollution Research* **2023**, 31, 5668–5683. https://doi.org/10.1007/s11356-023-31405-8
31. Callahan B.J., McMurdie P.J., Rosen M.J., Han A.W., Johnson A.J.A., Holmes S.P. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods* **2016**, 13, 581–583. https://doi.org/10.1038/nmeth.3869
32. Chandra S., Sharma R., Singh K., Sharma A. Application of bioremediation technology in the environment contaminated with petroleum hydrocarbon. *Annals of Microbiology* **2013**, 63, 417–431. https://doi.org/10.1007/s13213-012-0543-3
33. Chaudhary D.K., Kim J. New insights into bioremediation strategies for oil-contaminated soil in cold environments. *International Biodeterioration & Biodegradation* **2019**, 142, 58–72. https://doi.org/10.1016/j.ibiod.2019.05.001
34. Chen X., Wang S., Lai H., Deng L., Zhong Q., Okoye C.O., Niu Q., Jing Y., Wang J., Jiang J. Nano-Biotechnology in Soil Remediation: Use of Nanomaterials to Promote Plant Growth and Stress Tolerance. *Nanomaterials* **2025**, 15, 1743. https://doi.org/10.3390/nano15221743
35. Cheng J., Sun Z., Yu Y., Li X., Li T. Effects of modified carbon black nanoparticles on plant-microbe remediation of petroleum and heavy metal co-contaminated soils. *International Journal of Phytoremediation* **2019**, 21, 634–642. https://doi.org/10.1080/15226514.2018.1556581

36. Chicca I., Becarelli S., Dartiahl C., La China S., De Kievit T., Petroni G., Di Gregorio S., Levin D.B. Degradation of BTEX mixture by a new *Pseudomonas putida* strain: role of the quorum sensing in the modulation of the upper BTEX oxidative pathway. *Environmental Science and Pollution Research* **2020**, 27, 36203–36214. https://doi.org/10.1007/s11356-020-09650-y
37. Cooney J.J., Silver S.A., Beck E.A. Factors influencing hydrocarbon degradation in three freshwater lakes. *Microbial Ecology* **1985**, 11, 127–137. https://doi.org/10.1007/BF02010485
38. Cui J.-Q., H, Q.-S., Li, M.-H., Che H., Sun, M.-B., Wen J.-P. Comparative Study on Different Remediation Strategies Applied in Petroleum-Contaminated Soils. *International Journal of Environmental Research and Public Health* **2020**, 17, 1606. https://doi.org/10.3390/ijerph17051606
39. Czarny J., Staninska-Pięta J., Piotrowska-Cyplik A., Juzwa W., Wolniewicz A., Marecik R., Ławniczak Ł., Chrzanowski Ł. *Acinetobacter* sp. as the key player in diesel oil degrading community exposed to PAHs and heavy metals. *Journal of Hazardous Materials* **2020**, 383, 121168. https://doi.org/10.1016/j.jhazmat.2019.121168
40. Daâssi D., Qabil Almaghribi F. Petroleum-contaminated soil: environmental occurrence and remediation strategies. *3 Biotech* **2022**, 12, 139. https://doi.org/10.1007/s13205-022-03198-z
41. Dandie C.E., Weber J., Aleer S., Adetutu E.M., Ball A.S., Juhasz A.L. Assessment of five bioaccessibility assays for predicting the efficacy of petroleum hydrocarbon biodegradation in aged contaminated soils. *Chemosphere* **2010**, 81, 1061–1068. https://doi.org/10.1016/j.chemosphere.2010.09.059
42. Das N., Chandran P. Microbial Degradation of Petroleum Hydrocarbon Contaminants: An Overview. *Biotechnology Research International* **2011**, 1–13. https://doi.org/10.4061/2011/941810
43. Das A., Das N., Rajkumari J., Pandey Prisha, Pandey Piyush, Exploring the bioremediation potential of *Bacillus* spp. for sustainable mitigation of hydrocarbon contaminants. *Environmental Sustainability* **2024**, 7, 135–156. https://doi.org/10.1007/s42398-024-00309-9
44. Dhanya M.S. Biosurfactant-enhanced bioremediation of petroleum hydrocarbons: potential issues, challenges, and future prospects, in: Bioremediation for Environmental Sustainability. Elsevier, **2021**, 215–250. https://doi.org/10.1016/B978-0-12-820524-2.00010-9
45. Dini S., Bekhit A.E.-D.A., Roohinejad S., Vale J.M., Agyei D. The Physicochemical and Functional Properties of Biosurfactants: A Review. *Molecules* **2024**, 29, 2544. https://doi.org/10.3390/molecules29112544
46. Dos Santos J.J., Maranho L.T. Rhizospheric microorganisms as a solution for the recovery of soils contaminated by petroleum: A review. *Journal of Environmental Management* **2018**, 210, 104–113. https://doi.org/10.1016/j.jenvman.2018.01.015
47. Doty S.L. Enhancing phytoremediation through the use of transgenics and endophytes. *New Phytologist* **2008**, 179, 318–333. https://doi.org/10.1111/j.1469-8137.2008.02446.x

48. Dudhagara D.R., Dave B.P. Mycobacterium as Polycyclic Aromatic Hydrocarbons (PAHs) Degrader, in: Ribón, W. (Ed.), *Mycobacterium* - Research and Development. *InTech*. **2018**.https://doi.org/10.5772/intechopen.73546
49. Elijah A.A. A Review of the Petroleum Hydrocarbons Contamination of Soil, Water and Air and the Available Remediation Techniques, Taking into Consideration the Sustainable Development Goals. *Earthline Journal of Chemical Sciences* **2022**, 97–113. https://doi.org/10.34198/ejcs.7122.97113
50. Eras-Muñoz E., Farré A., Sánchez A., Font X., Gea T. Microbial biosurfactants: a review of recent environmental applications. *Bioengineered* **2022**, 13, 12365–12391. https://doi.org/10.1080/21655979.2022.2074621
51. Eskandari S., Hoodaji M., Tahmourespour A., Abdollahi A., Baghi T., Eslamian S., Ostad-Ali-Askari K. Bioremediation of Polycyclic Aromatic Hydrocarbons by *Bacillus Licheniformis* ATHE9 and *Bacillus Mojavensis* ATHE13 as Newly Strains Isolated from Oil-Contaminated Soil. *Journal of Geography, Environment and Earth Science International* **2017**, 11, 1–11. https://doi.org/10.9734/JGEESI/2017/35447
52. Fan Y., Wang J., Gao C., Zhang Y., Du, W. A novel exopolysaccharide-producing and long-chain n-alkane degrading bacterium *Bacillus licheniformis* strain DM-1 with potential application for in-situ enhanced oil recovery. *Scientific Reports* **2020**, 10, 8519. https://doi.org/10.1038/s41598-020-65432-z
53. Feitkenhauer H., Müller R., MAuml;rkl H. Degradation of polycyclic aromatic hydrocarbons and long chain alkanes at 6070 °C by *Thermus* and *Bacillus* spp. *Biodegradation* **2003**, 14, 367–372. https://doi.org/10.1023/A:1027357615649
54. Feng L., Jiang X., Huang Y., Wen D., Fu T., Fu R. Petroleum hydrocarbon-contaminated soil bioremediation assisted by isolated bacterial consortium and sophorolipid. *Environmental Pollution* **2021**, 273, 116476. https://doi.org/10.1016/j.envpol.2021.116476
55. Finlay B.J., Black H.I.J., Brown S., Clarke K.J., Esteban G.F., Hindle R.M., Olmo J.L., Rollett A., Vickerman K. Estimating the Growth Potential of the Soil Protozoan Community. *Protist* **2000**, 151, 69–80. https://doi.org/10.1078/1434-4610-00008
56. Firmin S.L., Laurette N.N. Biodegradation of high-molecular-weight PAHs in polluted mangrove sediments using indigenous microflora and exogenous strains *Rhodococcus erythropolis* T902.1 and *Bacillus subtilis*. *Biodegradation* **2026**, 37, 41. https://doi.org/10.1007/s10532-026-10264-3
57. Foght J.M., Westlake D.W.S., Johnson W.M., Ridgway H.F. Environmental gasoline-utilizing isolates and clinical isolates of *Pseudomonas aeruginosa* are taxonomically indistinguishable by chemotaxonomic and molecular techniques. *Microbiology* **1996,** 142, 2333–2340. https://doi.org/10.1099/00221287-142-9-2333
58. Foissner W., Berger H. A user-friendly guide to the ciliates (Protozoa, Ciliophora) commonly used by hydrobiologists as bioindicators in rivers, lakes, and waste waters, with notes on their ecology. *Freshwater Biology* **1996**, 35, 375–482. https://doi.org/10.1111/j.1365-2427.1996.tb01775.x

59. Foissner W., Berger H., Kohmann F. Taxonomische Und ¨ Okologische Revision Der Ciliaten Des Saprobien Systems - Band II: Peritrichia, Heterotrichida, Odontostomatida. BayerischesLandesamt für Wasserwirtschaft, München, Germany, **1992**
60. Foissner W., Berger H., Schaumburg J. Identification and Ecology of Limnetic Plankton Ciliates. BayerischesLandesamt für Wasserwirtschaft, München, Germany, **1999**
61. Foissner W., Blatterer H., Berger H., Kohmann F. Taxonomische und ¨ okologischerevision der Ciliaten des saprobiensystems. Band I: Cyrtophorida, Oligotrichida, Hypotrichia, Colpodea. BayerischesLandesamt für Wasserwirtschaft, München, Germany, **1991**
62. Foucault Y., Durand M.-J., Tack K., Schreck E., Geret F., Leveque T., Pradere P., Goix S., Dumat C. Use of ecotoxicity test and ecoscores to improve the management of polluted soils: case of a secondary lead smelter plant. *Journal of Hazardous Materials* **2013**, 246–247, 291–299. https://doi.org/10.1016/j.jhazmat.2012.12.042
63. Fudryn G., Kowala Z. Odnowa zanieczyszczonych gruntów metodami in-situ. *Ochrona Środowiska* **1996**, 2, 27–34.
64. Furtak K., Gawryjołek K., Gałązka A., Grządziel J. The Response of Red Clover (Trifolium pratense L.) to Separate and Mixed Inoculations with *Rhizobium leguminosarum* and *Azospirillum brasilense* in Presence of Polycyclic Aromatic Hydrocarbons. *International Journal of Environmental Research and Public Health* **2020**, 17, 5751. https://doi.org/10.3390/ijerph17165751
65. Gałązka A. Przegląd biologicznych metod oczyszczania gleb skażonych wielopierścieniowymi węglowodorami aromatycznymi. *Zeszyty problemowe postępów nauk rolniczych* **2009**, 535, 103–110.
66. Gałązka A., Gałązka R. Phytoremediation of Polycyclic Aromatic Hydrocarbons in Soils Artificially Polluted Using Plant-Associated-Endophytic Bacteria and *Dactylis glomerata* as the Bioremediation Plant. *Polish Journal of Microbiology* **2015**, 64, 241–252. https://doi.org/10.5604/01.3001.0009.2119
67. Gałązka A., Król M., Pierzyński A. Bioremediation of crude oil derivatives in soils naturally and artificially polluted with the use of maize as the test plant part I. pahs degradation. *Acta Scientiarum Polonorum. Agricultura* **2010**, 9, 13–24.
68. Gan S., Lau E.V., Ng H.K. Remediation of soils contaminated with polycyclic aromatic hydrocarbons (PAHs). *Journal of Hazardous Materials* **2009**, 172, 532–549. https://doi.org/10.1016/j.jhazmat.2009.07.118
69. Gao H., Wu M., Liu H., Zhang T., Zhang X. Cell toxic damages during polycyclic aromatic hydrocarbons biodegradation by Pseudomonas aeruginosa G24. *Journal of Water Process Engineering* **2023**, 54, 103992. https://doi.org/10.1016/j.jwpe.2023.103992
70. García-Díaz C., Nebbioso A., Piccolo A., Barrera-Cortés J., Martínez-Palou R. Remediation of Hydrocarbon-Contaminated Soil by Washing with Novel Chemically Modified Humic Substances. *Journal of Environmental Quality* **2015**, 44, 1764–1771. https://doi.org/10.2134/jeq2014.09.0399

71. Ghorbannezhad H., Moghimi H., Dastgheib S.M.M. Biodegradation of high molecular weight hydrocarbons under saline condition by halotolerant *Bacillus subtilis* and its mixed cultures with Pseudomonas species. *Scientific Reports* **2022**, 12, 13227. https://doi.org/10.1038/s41598-022-17001-9
72. Ghosal D., Ghosh S., Dutta T.K., Ahn Y. Current State of Knowledge in Microbial Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *Frontiers in Microbiology* **2016**, 7. https://doi.org/10.3389/fmicb.2016.01369
73. Gomes M.P. Nanophytoremediation: advancing phytoremediation efficiency through nanotechnology integration. *Discover Plants* **2025**, 2, 8. https://doi.org/10.1007/s44372-025-00090-x
74. Greco Lucchina P., Catania V., Di Trapani D., Petta E.M., Sciré Calabrisotto L., Vinti G., Quatrini P., Viviani G. Sequential biostimulation and bioaugmentation treatments of a diesel-contaminated soil: effect on hydrocarbon degradation and soil bacterial communities. *Applied Soil Ecology* **2026**, 219, 106802. https://doi.org/10.1016/j.apsoil.2026.106802
75. Guirado M., García-Delgado C., Pindado O., De La Torre B.O., Escolano O., Eymar E., Millán R. Bioremediation study of a hydrocarbon-contaminated soil by profiling aromatic and aliphatic chains. *Applied Soil Ecology* **2023**, 190, 104983. https://doi.org/10.1016/j.apsoil.2023.104983
76. Gupta N., Banerjee S., Koley A., Bharali P., GhoshThakur R., Hoque R.R., Balachandran S. Strategies for remediation of polycyclic aromatic hydrocarbons in contaminated soil: A systematic review and bibliometric analysis. *Applied Soil Ecology* **2024**, 204, 105688. https://doi.org/10.1016/j.apsoil.2024.105688
77. Hall J., Soole K., Bentham R. Hydrocarbon Phytoremediation in the Family *Fabacea* —A Review. *International Journal of Phytoremediation* **2011**, 13, 317–332. https://doi.org/10.1080/15226514.2010.495143
78. Heidari S., Fotouhi Ghazvini R., Zavareh M., Kafi M. Physiological responses and phytoremediation ability of Eastern Coneflower (*Echinacea purpurea*) for crude oil contaminated soil. *Caspian Journal of Environmental Sciences* **2018**, 16. https://doi.org/10.22124/cjes.2018.2957
79. Heidarzadeh N., Jebeli M., Gitipour S. Cement-Based Solidification/Stabilization of Soil Contaminated by Polycyclic Aromatic Hydrocarbons Using Organophilic Clay*, Research Square* **2021**, 1-17. https://doi.org/10.21203/rs.3.rs-974704/v1
80. Hesham A.E.-L., Mohamed E.A., Mawad A.M.M., Elfarash A., Abd El-Fattah B.S., El-Rawy M.A. Molecular Characterization of *Fusarium Solani* Degrades a Mixture of Low and High Molecular Weight Polycyclic Aromatic Hydrocarbons. *The Open Biotechnology Journal* **2017**, 11, 27–35. https://doi.org/10.2174/1874070701711010027
81. Hong Y., Liao D., Chen J., Khan S., Su J., Li H. A comprehensive study of the impact of polycyclic aromatic hydrocarbons (PAHs) contamination on salt marsh plants *Spartina alterniflora*: implication for plant-microbe interactions in phytoremediation. *Environmental Science and Pollution Research* **2015**, 22, 7071–7081. https://doi.org/10.1007/s11356-014-3912-6
82. Horel A., Schiewer S. Microbial Degradation of Different Hydrocarbon Fuels with

Mycoremediation of Volatiles. *Microorganisms* **2020**, 8, 163. https://doi.org/10.3390/microorganisms8020163

83. Hossain Md.F., Akter Mst.A., Sohan Md.S.R., Sultana Dr.N., Reza M.A., Hoque K.Md.F. Bioremediation potential of hydrocarbon degrading bacteria: isolation, characterization, and assessment. *Saudi Journal of Biological Sciences* **2022**, 29, 211–216. https://doi.org/10.1016/j.sjbs.2021.08.069
84. http://arep.med.harvard.edu/labgc/adnan/projects/Utilities/revcomp.html (dostęp 21.03.2024)
85. https://tigret.eu/kategoria/toksycznosc/ (dostęp 20.09.2023)
86. https://www.ncbi.nlm.nih.gov/bioproject?term=PRJNA1224491&cmd=DetailsSearch (dostęp 24.01.2026)
87. http://www.ncbi.nlm.nih.gov/ BLAST (dostęp 14.03.2024)
88. Hu B., Wang M., Geng S., Wen L., Wu M., Nie Y., Tang Y.-Q., Wu X.-L. Metabolic Exchange with Non-Alkane-Consuming Pseudomonas stutzeri SLG510A3-8 Improves *n* -Alkane Biodegradation by the Alkane Degrader *Dietzia* sp. Strain DQ12-45-1b. *Applied and Environmental Microbiology* **2020**, 86, e02931-19. https://doi.org/10.1128/AEM.02931-19
89. Huang S., Dai C., Zhou Y., Peng H., Yi K., Qin P., Luo S., Zhang X. Comparisons of three plant species in accumulating polycyclic aromatic hydrocarbons (PAHs) from the atmosphere: a review. *Environmental Science and Pollution Research* **2018**, 25, 16548–16566. https://doi.org/10.1007/s11356-018-2167-z
90. Huizenga J.M., Semprini L. Influence of growth substrate and contaminant mixtures on the degradation of BTEX and MTBE by *Rhodococcus rhodochrous* ATCC strain 21198. *Biodegradation* **2023**, 34, 461–475. https://doi.org/10.1007/s10532-023-10037-2
91. Ijaz A., Imran A., Anwar Ul Haq M., Khan Q.M., Afzal M. Phytoremediation: recent advances in plant-endophytic synergistic interactions. *Plant Soil* **2016**, 405, 179–195. https://doi.org/10.1007/s11104-015-2606-2
92. Innes C. Pleurotus spp. as Agents of Mycoremediation: A Review. University Honors Theses, Portland State University **2023**, 1358 https://doi.org/10.15760/honors.1387
93. Ivshina I.B., Kuyukina M.S., Krivoruchko A.V., Tyumina E.A. The *Rhodococcus ruber Actinomycetes*, Key and Universal Biooxidizers of C2−C4 Gaseous Alkanes. *Microbiology* **2025**, 94, 145–171. https://doi.org/10.1134/S0026261724607620
94. Jaekel U., Musat N., Adam B., Kuyper M., Grundmann, O., Musat F. Anaerobic degradation of propane and butane by sulfate-reducing bacteria enriched from marine hydrocarbon cold seeps. *The ISME Journal* **2013**, 7, 885–895. https://doi.org/10.1038/ismej.2012.159
95. Janbandhu A., Fulekar M.H. Biodegradation of phenanthrene using adapted microbial consortium isolated from petrochemical contaminated environment. *Journal of Hazardous Materials* **2011**, 187, 333–340. https://doi.org/10.1016/j.jhazmat.2011.01.034
96. Jiang C., Liu S., Phenrat T., Sui Q. Nanomaterial-Enabled Environmental

Remediation and Removal of Emerging Pollutants. *Toxics* **2025**, 13, 810. https://doi.org/10.3390/toxics13100810

97. Kadri T., Magdouli S., Rouissi T., Brar S.K. Ex-situ biodegradation of petroleum hydrocarbons using *Alcanivorax borkumensis* enzymes. *Biochemical Engineering Journal* **2018**, 132, 279–287. https://doi.org/10.1016/j.bej.2018.01.014
98. Kalia A., Sharma S., Semor N., Babele P.K., Sagar S., Bhatia R.K., Walia A. Recent advancements in hydrocarbon bioremediation and future challenges: a review. *3 Biotech* **2022**, 12*,* 135. https://doi.org/10.1007/s13205-022-03199-y
99. Kamath R., Rentz J.A., Schnoor J.L., Alvarez P.J.J. Chapter 16 Phytoremediation of hydrocarbon-contaminated soils: principles and applications, in: Studies in Surface Science and Catalysis. Elsevier, **2004**, 447–478. https://doi.org/10.1016/S0167-2991(04)80157-5
100. Kaplan C.W., Kitt, C.L. Bacterial Succession in a Petroleum Land Treatment Unit. *Applied and Environmental Microbiology* **2004**, 70, 1777–1786. https://doi.org/10.1128/AEM.70.3.1777-1786.2004
101. Karlapudi A.P., Venkateswarulu T.C., Tammineedi J., Kanumuri L., Ravuru B.K., Dirisala V.R., Kodali V.P. Role of biosurfactants in bioremediation of oil pollution-a review. *Petroleum* **2018**, 4, 241–249. https://doi.org/10.1016/j.petlm.2018.03.007
102. Kessler A., Hedberg J., Blomberg E., Odnevall I. Reactive Oxygen Species Formed by Metal and Metal Oxide Nanoparticles in Physiological Media—A Review of Reactions of Importance to Nanotoxicity and Proposal for Categorization. *Nanomaterials* **2022**, 12, 1922. https://doi.org/10.3390/nano12111922
103. Khan M.A.I., Biswas B., Smith E., Naidu R., Megharaj M. Toxicity assessment of fresh and weathered petroleum hydrocarbons in contaminated soil- a review. *Chemosphere* **2018**, 212, 755–767. https://doi.org/10.1016/j.chemosphere.2018.08.094
104. Khan S., Afzal M., Iqbal S., Khan Q.M. Plant–bacteria partnerships for the remediation of hydrocarbon contaminated soils. *Chemosphere* **2013**, 90, 1317–1332. https://doi.org/10.1016/j.chemosphere.2012.09.045
105. Kim D., Chae J.-C., Zylstra G.J., Kim Y.-S., Kim S.-K., Nam M.H., Kim Y.M., Kim E. Identification of a Novel Dioxygenase Involved in Metabolism of *o* -Xylene, Toluene, and Ethylbenzene by *Rhodococcus* sp. Strain DK17. *Applied and Environmental Microbiology* **2004**, 70, 7086–7092. https://doi.org/10.1128/AEM.70.12.7086-7092.2004
106. Klindworth A., Pruesse E., Schweer T., Peplies J., Quast C., Horn M., Glöckner F.O. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research* **2013**, 41, e1–e1. https://doi.org/10.1093/nar/gks808
107. Kluk D., Steliga T. Efektywna metoda identyfikacji zanieczyszczeń ropopochodnych (TPH) i wielopierścieniowych węglowodorów aromatycznych (WWA) w glebach. *Nafta-Gaz* **2017**, 73, 488–495. https://doi.org/10.18668/NG.2017.07.06

108. Koshlaf E., Ball A.S. Soil bioremediation approaches for petroleum hydrocarbon polluted environments. *AIMS Microbiology* **2017**, 3, 25–49. https://doi.org/10.3934/microbiol.2017.1.25
109. Kotani T., Kawashima Y., Yurimoto H., Kato N., Sakai Y. Gene structure and regulation of alkane monooxygenases in propane-utilizing *Mycobacterium* sp. TY-6 and Pseudonocardia sp. TY-7. *Journal of Bioscience and Bioengineering* **2006,** 102, 184–192. https://doi.org/10.1263/jbb.102.184
110. Kowalska E., Więch M. Procesy i systemy termicznej degradacji odpadów niebezpiecznych. *Elektronika* **2004**, 36–40.
111. Kuppusamy S., Thavamani P., Venkateswarlu K., Lee Y.B., Naidu R., Megharaj M. Remediation approaches for polycyclic aromatic hydrocarbons (PAHs) contaminated soils: Technological constraints, emerging trends and future directions. *Chemosphere* 2017, 168, 944–968. https://doi.org/10.1016/j.chemosphere.2016.10.115
112. Kurth E.G., Doughty D.M., Bottomley P.J., Arp D.J., Sayavedra-Soto L.A. Involvement of BmoR and BmoG in n-alkane metabolism in '*Pseudomonas butanovora*.' *Microbiology* **2008**, 154, 139–147. https://doi.org/10.1099/mic.0.2007/012724-0
113. Laczi K., Erdeiné Kis Á., Szilágyi Á., Bounedjoum N., Bodor A., Vincze G.E., Kovács T., Rákhely G., Perei K. New Frontiers of Anaerobic Hydrocarbon Biodegradation in the Multi-Omics Era. *Frontiers in Microbiology* **2020**, 11, 590049. https://doi.org/10.3389/fmicb.2020.590049
114. Lalucat J., Bennasar A., Bosch R., García-Valdés E., Palleroni N.J. Biology of *Pseudomonas stutzeri*. *Microbiology and Molecular Biology Reviews* **2006**, 70, 510–547. https://doi.org/10.1128/MMBR.00047-05
115. Leahy J.G., Colwell R.R. Microbial degradation of hydrocarbons in the environment. *Microbiological Reviews* **1990**, 54, 305–315. https://doi.org/10.1128/mr.54.3.305-315.1990
116. Li F., Guo S., Wu B., Wang S. Pilot-scale electro-bioremediation of heavily PAH-contaminated soil from an abandoned coking plant site. *Chemosphere* **2020**, 244, 125467. https://doi.org/10.1016/j.chemosphere.2019.125467
117. Li L., Zhu P., Wang X., Zhang Z. Phytoremediation effect of *Medicago sativa* colonized by Piriformospora indica in the phenanthrene and cadmium co-contaminated soil. *BMC Biotechnology* **2020**, 20. https://doi.org/10.1186/s12896-020-00613-2
118. Li Q. Mining secondary metabolism of achromobacter and analysis of key genes of petroleum degradation. IOP Conference Series: Earth and Environmental Science **2021**, 692, 042032. https://doi.org/10.1088/1755-1315/692/4/042032
119. Liao C., Xu W., Lu G., Liang X., Guo C., Yang C., Dang Z. Accumulation of Hydrocarbons by Maize (*Zea mays* L.) in Remediation of Soils Contaminated with Crude Oil. *International Journal of Phytoremediation* **2015**, 17, 693–700. https://doi.org/10.1080/15226514.2014.964840
120. Lima T.M.S., Procópio L.C., Brandão F.D., Leão B.A., Tótola M.R., Borges A.C. Evaluation of bacterial surfactant toxicity towards petroleum degrading

microorganisms. *Bioresource Technology* **2011**, 102, 2957–2964. https://doi.org/10.1016/j.biortech.2010.09.109

121. Liu M., Chen G., Xu L., He Z., Ye Y. Environmental remediation approaches by nanoscale zero valent iron (nZVI) based on its reductivity: a review. *RSC Advances* **2024**, 14, 21118–21138. https://doi.org/10.1039/D4RA02789B
122. Liu R., Jadeja R.N., Zhou Q., Liu Z. Treatment and Remediation of Petroleum-Contaminated Soils Using Selective Ornamental Plants. *Environmental Engineering Science* **2012**, 29, 494–501. https://doi.org/10.1089/ees.2010.0490
123. Liu X., Yang W., He L., Zhang X., Cao T., Wan Y., Yuan L., Mi X. Unlocking metabolic suppression in *Acinetobacter baumannii* with synergistic N-P-surfactant to enable petroleum degradation in saline soils. *Journal of Environmental Sciences* **2026,** S100107422600046X. https://doi.org/10.1016/j.jes.2026.01.044
124. Lopez E.S., Elufisan T.O., Bustos P., Charles C.P.M., Mendoza-Herrera A., Guo X. Complete Genome Report of a Hydrocarbon-Degrading *Sphingobium yanoikuyae* S72. *Applied Sciences* **2022**, 12, 6201. https://doi.org/10.3390/app12126201
125. Lü H., Tang G.-X., Huang Y.-H., Mo C.-H., Zhao H.-M., Xiang L., Li Y.-W., Li H., Cai Q.-Y., Li Q.X. Response and adaptation of rhizosphere microbiome to organic pollutants with enriching pollutant-degraders and genes for bioremediation: A critical review. *Science of The Total Environment* **2024**, 912, 169425. https://doi.org/10.1016/j.scitotenv.2023.169425
126. Ławniczak Ł., Woźniak-Karczewska M., Loibner A.P., Heipieper H.J., Chrzanowski Ł. Microbial Degradation of Hydrocarbons—Basic Principles for Bioremediation: A Review. *Molecules* **2020**, 25, 856. https://doi.org/10.3390/molecules25040856
127. Ma C., Han L., Shang H., Hao Y., Xu X., White J.C., Wang Z., Xing B. Nanomaterials in agricultural soils: Ecotoxicity and application. *Current Opinion in Environmental Science & Health* **2023**, 31, 100432. https://doi.org/10.1016/j.coesh.2022.100432
128. Ma J., Zhuang Y., Wang Y., Zhu N., Wang T., Xiao H., Chen J. Update on new trend and progress of the mechanism of polycyclic aromatic hydrocarbon biodegradation by *Rhodococcus*, based on the new understanding of relevant theories: a review. *Environmental Science and Pollution Research* **2023**, 30, 93345–93362. https://doi.org/10.1007/s11356-023-28894-y
129. Ma Y., Wang J., Liu Y., Wang X., Zhang B., Zhang W., Chen T., Liu G., Xue L., Cui X. Nocardioides: "Specialists" for Hard-to-Degrade Pollutants in the Environment. *Molecules* **2023**, 28, 7433. https://doi.org/10.3390/molecules28217433
130. Ma Y.-L., Lu W., Wan L.-L., Luo N. Elucidation of Fluoranthene Degradative Characteristics in a Newly Isolated *Achromobacter xylosoxidans* DN002. *Applied Biochemistry and Biotechnology* **2015**, 175, 1294–1305. https://doi.org/10.1007/s12010-014-1347-7
131. Małachowska-Jutsz A., Janosz W., Rudek J. Toksyczność gleby zanieczyszczonej olejem silnikowym poddanej samooczyszczaniu oraz fitoremediacji. *Ochrona*

Środowiska **2012**, 34, 15–22.
132. Maletic S., Dalmacija B., Roncevic S. Petroleum Hydrocarbon Biodegradability in Soil – Implications for Bioremediation, in: Kutcherov, V. (Ed.), Hydrocarbon. *InTech* **2013**. https://doi.org/10.5772/50108
133. Malina G., Szczepański A. Likwidacja zanieczyszczeń substancjami ropopochodnymi w środowisku wodno-gruntowym. , Państwowa Inspekcja Ochrony Środowiska, Biblioteka Monitoringu Środowiska **1994**.
134. Malina G. Likwidacja zagrożenia środowiska gruntowo-wodnego na terenach zanieczyszczonych. . Wydawnictwo Politechniki Częstochowskiej **2007**.
135. Malińska K. Application of a modified OxiTop® respirometer for laboratory composting studies. *Archives of Environmental Protection* **2016**, 42, 56–62. https://doi.org/10.1515/aep-2016-0007
136. Mamindy-Pajany Y., Hamer B., Roméo M., Géret F., Galgani F., Durmiši E., Hurel C., Marmier N. The toxicity of composted sediments from Mediterranean ports evaluated by several bioassays. *Chemosphere* **2011**, 82, 362–369. https://doi.org/10.1016/j.chemosphere.2010.10.005
137. Manjunatha B.K., Vamshee V.K., Poojitha N.G., Kiran R.J., Mythili S.T., Divakara R., Sreenivasa R.A., Vidya S.M. Bioremediation of polycyclic aromatic hydrocarbons (acenaphthene, anthracene, fluoranthene, fluorene) by *Aspergillus niger* HQ170509.1 fungus. *International Journal of Environmental Science and Technology* **2025**, 22, 12567–12582. https://doi.org/10.1007/s13762-025-06370-z
138. Mekonnen B.A., Aragaw T.A., Genet M.B. Bioremediation of petroleum hydrocarbon contaminated soil: a review on principles, degradation mechanisms, and advancements. *Frontiers in Environmental Science* **2024,** 12, 1354422. https://doi.org/10.3389/fenvs.2024.1354422
139. Mnif S., Sayadi S., Chamkha M. Biodegradative potential and characterization of a novel aromatic-degrading bacterium isolated from a geothermal oil field under saline and thermophilic conditions. *International Biodeterioration & Biodegradation* **2014**, 86, 258–264. https://doi.org/10.1016/j.ibiod.2013.09.015
140. Mohanapragash A. G., Kaleeswari R. K., Madhupriyaa D., Salma Santhosh S., Meena S., Baskar M., Umamaheswari T., Selvamurugan M., Ramesh T. Nanotechnology in sustainable soil remediation: exploring a review on carbon-based, clay-based, dendrimer, polymer, and nanocomposite based nanomaterials for environmental cleanup. *International Journal of Environmental Analytical Chemistry* **2025**,105, 8533–8558. https://doi.org/10.1080/03067319.2025.2478610
141. Moore F.P., Barac T., Borremans B., Oeyen L., Vangronsveld J., Van Der Lelie D., Campbell C.D., Moore E.R.B. Endophytic bacterial diversity in poplar trees growing on a BTEX-contaminated site: The characterisation of isolates with potential to enhance phytoremediation. *Systematic and Applied Microbiology* **2006**, 29, 539–556. https://doi.org/10.1016/j.syapm.2005.11.012
142. Morillo E., Madrid F., Lara-Moreno A., Villaverde J. Soil bioremediation by cyclodextrins. A review. *International Journal of Pharmaceutics* **2020**, 591, 119943. https://doi.org/10.1016/j.ijpharm.2020.119943
143. Moses H., Odigure J.O., Otaru A.J., Oduoza C.F. A potential remediation of total

petroleum hydrocarbon from contaminated soils using surfactant suspended multiwall carbon nanotubes. *Results in Engineering* **2023**, 19, 101371. https://doi.org/10.1016/j.rineng.2023.101371

144. Mrozik A., Piotrowska-Seget Z., Łabuzek Z. Bacteria in bioremediation of hydrocarbon-contaminated environments . *Postępy Mikrobiologii* **2005**, 44, 227–238.
145. Mukherjee A.K., Bhagowati P., Biswa B.B., Chanda A., Kalita B. A comparative intracellular proteomic profiling of *Pseudomonas aeruginosa* strain ASP-53 grown on pyrene or glucose as sole source of carbon and identification of some key enzymes of pyrene biodegradation pathway. *Journal of Proteomics* **2017**, 167, 25–35. https://doi.org/10.1016/j.jprot.2017.07.020
146. Mullaeva S.A., Delegan Ya.A., Streletskii R.A., Sazonov O.I., Petrikov K.V., Ivanova A.A., Dyatlov I.A., Shemyakin I.G., Bogun A.G., Vetrova A.A. Pseudomonas veronii strain 7–41 degrading medium-chain n-alkanes and polycyclic aromatic hydrocarbons. *Scientific Reports* **2022**, 12, 20527. https://doi.org/10.1038/s41598-022-25191-5
147. Naderi Peikam E., Jalali M. Application of three nanoparticles (Al_2O_3, SiO_2 and TiO_2) for metal-contaminated soil remediation (measuring and modeling). *International Journal of Environmental Science and Technology* **2019**, 16, 7207–7220. https://doi.org/10.1007/s13762-018-2134-8
148. Naloka K., Jaroonrunganan J., Woratecha N., Khondee N., Nojiri H., Pinyakong O. Physiological changes in *Rhodococcus ruber* S103 immobilized on biobooms using low-cost media enhance stress tolerance and crude oil-degrading activity. *Scientific Reports* **2022**, 12, 10474. https://doi.org/10.1038/s41598-022-14488-0
149. Niyommaneerat W., Nakajima F., Tobino T., Yamamoto K. Development of a chronic sediment toxicity test using the benthic ostracod *Heterocypris incongruens* and their application to toxicity assessments of urban road dust. *Ecotoxicology and Environmental Safety* **2017**, 143, 266–274. https://doi.org/10.1016/j.ecoenv.2017.05.011
150. Nwadinigwe A.O., Ugwu E.C. Overview of Nano-phytoremediation Applications, in: Ansari, A.A., Gill, S.S., Gill, R., R. Lanza, G., Newman, L. (Eds.), Phytoremediation. Springer International Publishing, Cham, **2018**, 377–382. https://doi.org/10.1007/978-3-319-99651-6_15
151. Nwankwegu A.S., Orji M.U., Onwosi C.O. Studies on organic and in-organic biostimulants in bioremediation of diesel-contaminated arable soil. *Chemosphere* **2016**, 162, 148–156. https://doi.org/10.1016/j.chemosphere.2016.07.074
152. Oleszczuk P., Jośko I., Kuśmierz M., Futa B., Wielgosz E., Ligęza S., Pranagal J. Microbiological, biochemical and ecotoxicological evaluation of soils in the area of biochar production in relation to polycyclic aromatic hydrocarbon content. *Geoderma* **2014**, 213, 502–511. https://doi.org/10.1016/j.geoderma.2013.08.027
153. Ortega-Calvo J.J., Lahlou M., Saiz-Jimenez C. Effect of organic matter and clays on the biodegradation of Phenanthrene in soils. *International Biodeterioration & Biodegradation* **1997**, 40, 101–106. https://doi.org/10.1016/S0964-8305(97)00045-0

154. Ossai I.C., Ahmed A., Hassan A., Hamid F.S. Remediation of soil and water contaminated with petroleum hydrocarbon: A review. *Environmental Technology & Innovation* **2020**, 17, 100526. https://doi.org/10.1016/j.eti.2019.100526
155. Özlü V., Elif Fırat M. Fly ash and nano-graphene enhanced stabilization of engine oil-contaminated soils. *Nanotechnology Reviews* **2025**, 14, 20250210. https://doi.org/10.1515/ntrev-2025-0210
156. Pacwa-Płociniczak M., Płaza G.A., Piotrowska-Seget Z., Cameotra S.S. Environmental Applications of Biosurfactants: Recent Advances. *International Journal of Molecular Sciences* **2011**, 12, 633–654. https://doi.org/10.3390/ijms12010633
157. Pandolfo E., Barra Caracciolo A., Rolando L. Recent Advances in Bacterial Degradation of Hydrocarbons. *Water* **2023**, 15, 375. https://doi.org/10.3390/w15020375
158. Parthipan P., Preetham E., Machuca L.L., Rahman P.K.S.M., Murugan K., Rajasekar A. Biosurfactant and Degradative Enzymes Mediated Crude Oil Degradation by *Bacterium Bacillus subtilis* A1. *Frontiers in Microbiology* **2017**, 8. https://doi.org/10.3389/fmicb.2017.00193
159. Patel A.B., Shaikh S., Jain K.R., Desai C., Madamwar D. Polycyclic Aromatic Hydrocarbons: Sources, Toxicity, and Remediation Approaches. *Frontiers in Microbiology* **2020**, 11, 562813. https://doi.org/10.3389/fmicb.2020.562813
160. Patowary K., Patowary R., Kalita M.C., Deka S. Characterization of Biosurfactant Produced during Degradation of Hydrocarbons Using Crude Oil As Sole Source of Carbon. *Frontiers in Microbiology* **2017**, 8. https://doi.org/10.3389/fmicb.2017.00279
161. Pawlik M., Cania B., Thijs S., Vangronsveld J., Piotrowska-Seget Z. Hydrocarbon degradation potential and plant growth-promoting activity of culturable endophytic bacteria of *Lotus corniculatus* and *Oenothera biennis* from a long-term polluted site. *Environmental Science and Pollution Research* **2017**, 24, 19640–19652. https://doi.org/10.1007/s11356-017-9496-1
162. Pawlik M., Piotrowska-Seget Z. Endophytic Bacteria Associated with *Hieracium piloselloides* : Their Potential for Hydrocarbon-Utilizing and Plant Growth-Promotion. *Journal of Toxicology and Environmental Health* **2015**, 78, 860–870. https://doi.org/10.1080/15287394.2015.1051200
163. Pawluśkiewic, B., Małuszyńska I., Małuszyński M., Dąbrowski P., Gnatowski T. Biometric indicators of growth and development of *Lolium perenne* and *Trifolium repens* in terms of recultivation of soil contaminated with petroleum. *Journal of Water and Land Development* **2022**, 70–79. https://doi.org/10.24425/jwld.2022.143722
164. Peng S., Zhou Q., Cai Z., Zhang Z. Phytoremediation of petroleum contaminated soils by *Mirabilis Jalapa* L. in a greenhouse plot experiment. *Journal of Hazardous Materials* **2009**, 168, 1490–1496. https://doi.org/10.1016/j.jhazmat.2009.03.036
165. Pérez-Hernández H., Pérez-Moreno A.Y., Méndez-López A., Fernández-Luqueño F. Effect of ZnO Nanoparticles During the Process of Phytoremediation of Soil

Contaminated with As and Pb Cultivated with Sunflower (*Helianthus annuus* L.). *International Journal of Environmental Research* **2023**, 18, 7. https://doi.org/10.1007/s41742-023-00556-4

166. Petrović O., Knežević P., Marković J., Rončević S. Screening method for detection of hydrocarbon-oxidizing bacteria in oil-contaminated water and soil specimens. *Journal of Microbiological Methods* **2008**, 74, 110–113. https://doi.org/10.1016/j.mimet.2008.03.012
167. Podsiadło Ł., Krzyśko-Łupicka T. Techniki bioremediacji substancji ropopochodnych i metody oceny ich efektywności. *Inżynieria i Ochrona Środowiska* **2013**, 16, 456–476.
168. Polyak Y.M., Bakin L.G., Chugunova M.V., Mayachkina N.V., Gerasimov A.O., Bure V.M. Effect of remediation strategies on biological activity of oil-contaminated soil - A field study. *International Biodeterioration & Biodegradation* **2018**, 126, 57–68. https://doi.org/10.1016/j.ibiod.2017.10.004
169. Qi Y.-B., Wang C.-Y., Lv C.-Y., Lun Z.-M., Zheng C.-G. Removal Capacities of Polycyclic Aromatic Hydrocarbons (PAHs) by a Newly Isolated Strain from Oilfield Produced Water. *International Journal of Environmental Research and Public Health* **2017**, 14, 215. https://doi.org/10.3390/ijerph14020215
170. Quast C., Pruesse E., Yilmaz P., Gerken J., Schweer T., Yarza P., Peplies J., Glöckner F.O. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research* **2012**, 41, D590–D596. https://doi.org/10.1093/nar/gks1219
171. Rabêlo F.H.S., Vangronsveld J., Baker A.J.M., Van Der Ent A., Alleoni L.R.F. Are Grasses Really Useful for the Phytoremediation of Potentially Toxic Trace Elements? A Review. *Frontiers in Plant Science* **2021**, 12, 778275. https://doi.org/10.3389/fpls.2021.778275
172. Radhakrishnan A., Balaganesh P., Vasudevan M., Natarajan N., Chauhan A., Arora J., Ranjan A., Rajput V.D., Sushkova S., Minkina T., Basniwal R.K., Kapardar R., Srivastav R. Bioremediation of Hydrocarbon Pollutants: Recent Promising Sustainable Approaches, Scope, and Challenges. *Sustainability* **2023**, 15, 5847. https://doi.org/10.3390/su15075847
173. Radwan K., Ślosorz Z., Rakowska J. Efekty środowiskowe usuwania zanieczyszczeń ropopochodnych. *Technika i Technologia* **2012**, 107–114.
174. Rai P.K., Kim K.-H., Lee S.S., Lee J.-H. Molecular mechanisms in phytoremediation of environmental contaminants and prospects of engineered transgenic plants/microbes. *Science of The Total Environment* **2020**, 705, 135858. https://doi.org/10.1016/j.scitotenv.2019.135858
175. Rakowska J. Usuwanie substancji ropopochodnych z dróg i gruntów, Monografie CNBOP-PIB 2012. Wydawnictwo Centrum Naukowo-Badawczego Ochrony Przeciwpożarowej im. Józefa Tuliszkowskiego, Józefów **2012**.
176. Ramakrishna T.R.B., Nalder T.D., Yang W., Marshall S.N., Barrow C.J. Controlling enzyme function through immobilisation on graphene, graphene derivatives and other two dimensional nanomaterials. *Journal of Materials Chemistry B* **2018**, 6, 3200–3218. https://doi.org/10.1039/C8TB00313K

177. Reeves R.D. Tropical hyperaccumulators of metals and their potential for phytoextraction. *Plant and Soil* **2003**, 249, 57–65. https://doi.org/10.1023/A:1022572517197
178. Reeves R.D., Baker A.J.M., Jaffré T., Erskine P.D., Echevarria G., Van Der Ent A. A global database for plants that hyperaccumulate metal and metalloid trace elements. *New Phytologist* **2018**, 218, 407–411. https://doi.org/10.1111/nph.14907
179. Ren W., Liu H., Mao T., Teng Y., Zhao R., Luo Y. Enhanced remediation of PAHs-contaminated site soil by bioaugmentation with graphene oxide immobilized bacterial pellets. *Journal of Hazardous Materials* **2022,** 433, 128793. https://doi.org/10.1016/j.jhazmat.2022.128793
180. Revathy T., Jayasri M.A., Suthindhiran K. Biodegradation of PAHs by *Burkholderia* sp. VITRSB1 Isolated from Marine Sediments. *Scientifica* **2015**, 1–9. https://doi.org/10.1155/2015/867586
181. Rohrbacher F., St-Arnaud M. Root Exudation: The Ecological Driver of Hydrocarbon Rhizoremediation. *Agronomy* **2016**, 6, 19. https://doi.org/10.3390/agronomy6010019
182. Rosa A.P., Triguis J.A. Bioremediation process on Brazil shoreline: Laboratory experiments. *Environmental Science and Pollution Research - International* **2007**, 14, 470–476. https://doi.org/10.1065/espr2007.02.377
183. Rouvalis A., Theodoropoulos C., Georgudaki J.I. Assessment of toxicity of the untreated and *Pleurotus ostreatus* treated olive mill wastewater by using microbiotests. *International Journal of Environmental Engineering* **2013**, 5, 373. https://doi.org/10.1504/IJEE.2013.056623
184. Sakshi Singh S.K., Haritash A.K. Polycyclic aromatic hydrocarbons: soil pollution and remediation. *International Journal of Environmental Science and Technology* **2019**, 16, 6489–6512. https://doi.org/10.1007/s13762-019-02414-3
185. Salari M., Rahmanian V., Hashemi S.A., Chiang W.-H., Lai C.W., Mousavi S.M., Gholami A. Bioremediation Treatment of Polyaromatic Hydrocarbons for Environmental Sustainability. *Water* **2022**, 14, 3980. https://doi.org/10.3390/w14233980
186. Sarfaraz A., Sumbal S., Qin Y., Faqir Y., Zveushe O.K., Zhou L., Zhang W., Li, J., Lv Z., Han Y., Dong F. Bioaugmentation-assisted phytoremediation of petroleum hydrocarbon-contaminated soils. *Journal of Environmental Chemical Engineering* **2025**, 13, 115895. https://doi.org/10.1016/j.jece.2025.115895
187. Sarkar J., Kazy S.K., Gupta A., Dutta A., Mohapatra B., Roy A., Bera P., Mitra A., Sar P. Biostimulation of Indigenous Microbial Community for Bioremediation of Petroleum Refinery Sludge. *Frontiers in Microbiology* **2016**, 7, 1407. https://doi.org/10.3389/fmicb.2016.01407
188. Sattar S., Siddiqui S., Shahzad A., Bano A., Naeem M., Hussain R., Khan N., Jan B.L., Yasmin H. Comparative Analysis of Microbial Consortiums and Nanoparticles for Rehabilitating Petroleum Waste Contaminated Soils. *Molecules* **2022**, 27, 1945. https://doi.org/10.3390/molecules27061945
189. Sharma J., Sundar D., Srivastava P. Biosurfactants: Potential Agents for Controlling Cellular Communication, Motility, and Antagonism. *Frontiers in*

Molecular Biosciences **2021**, 8, 727070. https://doi.org/10.3389/fmolb.2021.727070

190. Silva N.M., De Oliveira A.M.S.A., Pegorin S., Giusti C.E., Ferrari V.B., Barbosa D., Martins L.F., Morais C., Setubal J.C., Vasconcellos S.P., Da Silva A.M., De Oliveira J.C.F., Pascon R.C., Viana-Niero C. Characterization of novel hydrocarbon-degrading *Gordonia paraffinivorans* and *Gordonia sihwensis* strains isolated from composting. *PLoS ONE* **2019**, 14, e0215396. https://doi.org/10.1371/journal.pone.0215396
191. Singh P., Tiwary B.N. Optimization of conditions for polycyclic aromatic hydrocarbons (PAHs) degradation by *Pseudomonas stutzeri* P2 isolated from Chirimiri coal mines. *Biocatalysis and Agricultural Biotechnology* **2017**, 10, 20–29. https://doi.org/10.1016/j.bcab.2017.02.001
192. Singha L.P., Pandey P. Rhizosphere assisted bioengineering approaches for the mitigation of petroleum hydrocarbons contamination in soil. *Critical Reviews in Biotechnology* **2021**, 41, 749–766. https://doi.org/10.1080/07388551.2021.1888066
193. Siuta J. Podstawy biodegradacji ropopochodnych składników w glebach i odpadach. *Inżynieria Ekologiczna* **2001**, 2, 23–35.
194. Sivasamy S., Rajangam S., Kanagasabai T., Bisht D., Prabhakaran R., Dhandayuthapani S. Biocatalytic Potential of *Pseudomonas* Species in the Degradation of Polycyclic Aromatic Hydrocarbons. *Journal of Basic Microbiology* **2025**, 65, e2400448. https://doi.org/10.1002/jobm.202400448
195. Smułek W., Sydow M., Zabielska-Matejuk J., Kaczorek E. Bacteria involved in biodegradation of creosote PAH – A case study of long-term contaminated industrial area. *Ecotoxicology and Environmental Safety* **2020**, 187, 109843. https://doi.org/10.1016/j.ecoenv.2019.109843
196. Song L., Niu X., Zhang N., Li T. Effect of biochar-immobilized *Sphingomonas* sp. PJ2 on bioremediation of PAHs and bacterial community composition in saline soil. *Chemosphere* **2021**, 279, 130427. https://doi.org/10.1016/j.chemosphere.2021.130427
197. Souza E.C., Vessoni-Penna T.C., De Souza Oliveira R.P. Biosurfactant-enhanced hydrocarbon bioremediation: An overview. *International Biodeterioration & Biodegradation* **2014**, 89, 88–94. https://doi.org/10.1016/j.ibiod.2014.01.007
198. Souza L.R.R., Pomarolli L.C., Da Veiga M.A.M.S. From classic methodologies to application of nanomaterials for soil remediation: an integrated view of methods for decontamination of toxic metal(oid)s. *Environmental Science and Pollution Research* **2020**, 27, 10205–10227. https://doi.org/10.1007/s11356-020-08032-8
199. Stancu M.M. Characterization of new diesel-degrading bacteria isolated from freshwater sediments. *International Microbiology* **2022**, 26, 109–122. https://doi.org/10.1007/s10123-022-00277-2
200. Steliga T. Bioremediacja odpadów wiertniczych zanieczyszczonych substancjami ropopochodnymi ze starych dołów urobkowych. Instytut Nafty i Gazu, Kraków **2009**
201. Steliga T., Jakubowicz P., Kapusta P. Changes in toxicity during in situ

bioremediation of weathered drill wastes contaminated with petroleum hydrocarbons. *Bioresource Technology* **2012**, 125, 1–10. https://doi.org/10.1016/j.biortech.2012.08.092

202. Steliga T., Jakubowicz P., Kapusta P., Wojtowicz K. Efektywność biodegradacji zanieczyszczeń ropopochodnych w zestarzałym odpadzie z dołu urobkowego. *Nafta-Gaz* **2018a**, 74, 752–758. https://doi.org/10.18668/NG.2018.10.07
203. Steliga T., Jakubowicz P., Wojtowicz K., Kluk D. Zastosowanie testów toksykologicznych w przemyśle naftowym. *Nafta-Gaz* **2018b**, 74, 684–689. https://doi.org/10.18668/NG.2018.09.07
204. Steliga T., Kluk D. Assessment of the Suitability of *Melilotus officinalis* for Phytoremediation of Soil Contaminated with Petroleum Hydrocarbons (TPH and PAH), Zn, Pb and Cd Based on Toxicological Tests. *Toxics* **2021**, 9, 148. https://doi.org/10.3390/toxics9070148
205. Steliga T., Kluk D. Application of Festuca arundinacea in phytoremediation of soils contaminated with Pb, Ni, Cd and petroleum hydrocarbons. *Ecotoxicology and Environmental Safety* **2020,** 194, 110409. https://doi.org/10.1016/j.ecoenv.2020.110409
206. Steliga T., Uliasz M. Spent drilling muds management and natural environment protection. *Gospodarka Surowcami Mineralnymi* **2014**, 30, 135–155. https://doi.org/10.2478/gospo-2014-0011
207. Steliga T., Wojtowicz K. Wykorzystanie testów respirometrycznych do oceny efektywności biodegradacji osadów z instalacji kopalnianych. *Nafta-Gaz* **2019**, 75, 29–37. https://doi.org/10.18668/NG.2019.01.05
208. Steliga T.; Wojtowicz K.; Kapusta P.; Brzeszcz J. Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture. *Molecules* **2020**, *25*, 709, doi:10.3390/molecules25030709.
209. Surygała J. Kryteria doboru metod oczyszczania gruntów. Oficyna Wydawnicza Politechniki Wrocławskiej pt. „Zanieczyszczenia naftowe w gruncie", pod redakcją J. Surygały **2000**, 189-205.
210. Tang H., Yu H., Li Q., Wang X., Gai Z., Yin G., Su F., Tao F., Ma C., Xu P. Genome Sequence of *Pseudomonas putida* Strain B6-2, a Superdegrader of Polycyclic Aromatic Hydrocarbons and Dioxin-Like Compounds. *Journal of Bacteriology* **2011**, 193, 6789–6790. https://doi.org/10.1128/JB.06201-11
211. Tiwari B., Manickam N., Kumari S., Tiwari A. Biodegradation and dissolution of polyaromatic hydrocarbons by *Stenotrophomonas* sp. *Bioresource Technology* **2016**, 216, 1102–1105. https://doi.org/10.1016/j.biortech.2016.06.047
212. Tonelli F.C.P., Tonelli F.M.P., Lemos M.S., Nunes N.A.D.M. Mechanisms of phytoremediation, in: Phytoremediation. Elsevier, **2022**, 37–64. https://doi.org/10.1016/B978-0-323-89874-4.00023-6
213. Towell M.G., Paton G.I., Semple K.T. The biodegradation of cable oil components: Impact of oil concentration, nutrient addition and bioaugmentation. *Environmental Pollution* **2011,** 159, 3777–3783. https://doi.org/10.1016/j.envpol.2011.06.043

214. Ulhassan Z., Khan I., Hussain M., Khan A.R., Hamid Y., Hussain S., Allakhverdiev S.I., Zhou W. Efficacy of metallic nanoparticles in attenuating the accumulation and toxicity of chromium in plants: Current knowledge and future perspectives. *Environmental Pollution* **2022**, 315, 120390. https://doi.org/10.1016/j.envpol.2022.120390
215. Ulrici W. Contaminated Soil Areas, Different Countries and Contaminants, Monitoring of Contaminants, in: Rehm, H. -J., Reed, G. (Eds.), Biotechnology Set. Wiley **2001**, 5–41. https://doi.org/10.1002/9783527620999.ch1m
216. Varjani S., Upasani V.N. Influence of abiotic factors, natural attenuation, bioaugmentation and nutrient supplementation on bioremediation of petroleum crude contaminated agricultural soil. *Journal of Environmental Management* **2019**, 245, 358–366. https://doi.org/10.1016/j.jenvman.2019.05.070
217. Vogel T.M. Bioaugmentation as a soil bioremediation approach. *Current Opinion in Biotechnology* **1996**, 7, 311–316. https://doi.org/10.1016/S0958-1669(96)80036-X
218. Wang D., Xie Y., Jaisi D.P., Jin Y. Effects of low-molecular-weight organic acids on the dissolution of hydroxyapatite nanoparticles. *Environmental Science: Nano* **2016**, 3, 768–779. https://doi.org/10.1039/C6EN00085A
219. Wang M., Zheng X., Oba B.T., Lin Y., Shen C., Huang X., Yang F., Xiao Q., Ding Y. Innovations in nanomaterials for remediation of heavy metal−polluted soil: Advances, mechanistic insights, and future prospects. *Nano Materials Science* **2026**, 8, 11–35. https://doi.org/10.1016/j.nanoms.2025.06.008
220. Wang T., Jing J., Huang P., Guo X., Li C., Qu Y. Bioremediation of alkane–containing saline soils using the long–chain alkane–degrading bacterium *Pseudomonas aeruginosa* DL: Effects, communities, and networks. *Journal of Hazardous Materials* **2025,** 488, 137401. https://doi.org/10.1016/j.jhazmat.2025.137401
221. Wang Y., Pan C., Chu W., Vipin A.K., Sun L. Environmental Remediation Applications of Carbon Nanotubes and Graphene Oxide: Adsorption and Catalysis. *Nanomaterials* **2019**, 9, 439. https://doi.org/10.3390/nano9030439
222. Weyens N., Taghavi S., Barac T., Van Der Lelie D., Boulet J., Artoi T., Carleer R., Vangronsveld J. Bacteria associated with oak and ash on a TCE-contaminated site: characterization of isolates with potential to avoid evapotranspiration of TCE. *Environmental Science and Pollution Research* **2009a**, 16, 830–843. https://doi.org/10.1007/s11356-009-0154-0
223. Weyens N., Van Der Lelie D., Artois T., Smeets K., Taghavi S., Newman L., Carleer R., Vangronsveld J. Bioaugmentation with Engineered Endophytic Bacteria Improves Contaminant Fate in Phytoremediation. *Environmental Science & Technology* **2009b**, 43, 9413–9418. https://doi.org/10.1021/es901997z
224. Weyens N., Van Der Lelie D., Taghavi S., Newman L., Vangronsveld J. Exploiting plant–microbe partnerships to improve biomass production and remediation. *Trends in Biotechnology* **2009c**, 27, 591–598. https://doi.org/10.1016/j.tibtech.2009.07.006
225. Weyens N., Van Der Lelie D., Taghavi S., Vangronsveld J. Phytoremediation:

plant–endophyte partnerships take the challenge. *Current Opinion in Biotechnology* **2009d**, 20, 248–254. https://doi.org/10.1016/j.copbio.2009.02.012

226. Wilkes H., Buckel W., Golding B.T., Rabus R. Metabolism of Hydrocarbons in ***n***-Alkane-Utilizing Anaerobic Bacteria. *Microbial Physiology* **2016**, 26, 138–151. https://doi.org/10.1159/000442160
227. Willumsen P., Karlson U., Stackebrandt E., Kroppenstedt R.M. *Mycobacterium frederiksbergense* sp. nov., a novel polycyclic aromatic hydrocarbon-degrading Mycobacterium species. *International Journal of Systematic and Evolutionary Microbiology* **2001**, 51, 1715–1722. https://doi.org/10.1099/00207713-51-5-1715
228. Wojtowicz K. Opracowanie metodyki oznaczania WWA w próbkach gleb z wykorzystaniem chromatografii cieczowej HPLC. *Nafta-Gaz* **2022**, 78, 141–153. https://doi.org/10.18668/NG.2022.02.06
229. Wojtowicz K. Opracowanie metodyki oznaczania BTEX w próbkach gleb z wykorzystaniem chromatografii gazowej z przystawką headspace. *Nafta-Gaz* **2018**, 74, 201–207. https://doi.org/10.18668/NG.2018.03.03
230. Wojtowicz K., Jakubowicz P. Opracowanie metodyki oznaczania polichlorowanych bifenyli w próbkach gleb. *Nafta-Gaz* **2019**, 75, 420–429. https://doi.org/10.18668/NG.2019.07.06
231. Wojtowicz K., Steliga T. Wpływ mieszaniny biosurfaktantów na efektywność bioremediacji gruntu zanieczyszczonego węglowodorami ropopochodnymi. *Nafta-Gaz* **2024**, 80, 269–278. https://doi.org/10.18668/NG.2024.05.02
232. Wojtowicz K., Steliga T. Zastosowanie *Lotus corniculatus* w badaniach biodegradacji TPH i WWA wspomaganej fitoremediacją. *Nafta-Gaz* **2023**, 79, 443–454. https://doi.org/10.18668/NG.2023.07.01
233. Wojtowicz K.; Steliga T.; Brzeszcz J.; Fyda J.; Skalski T.; Kapusta P. Phytoremediation of Soil Contaminated with Petroleum Hydrocarbons Using the Wild Plants *Scirpus Sylvaticus* and *Cirsium Oleraceum*, Supported by Bioaugmentation. *International Biodeterioration & Biodegradation* **2026a**, *206*, 106217, doi:10.1016/j.ibiod.2025.106217.
234. Wojtowicz K., Steliga T., Kapusta P. Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using *Echinacea purpurea*. *Applied Sciences* **2023a**, 13, 13077. https://doi.org/10.3390/app132413077
235. Wojtowicz K., Steliga T., Kapusta P., Brzeszcz J. The Role of Graphene Oxide and Zinc Oxide Nanoparticles in Enhancing the Effectiveness of Phytoremediation of Petroleum Hydrocarbon-Contaminated Soils Using *Lolium perenne*. *Molecules* **2026b**, 31, 890. https://doi.org/10.3390/molecules31050890
236. Wojtowicz K., Steliga T., Kapusta P., Brzeszcz J.. Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (*Zea mays*). *Molecules* **2023b**, 28, 6104. https://doi.org/10.3390/molecules28166104
237. Wojtowicz K.; Steliga T.; Kapusta P.; Brzeszcz J.; Skalski T. Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil. *Materials* **2022**, *15*, 400,

doi:10.3390/ma15020400

238. Wojtowicz K.; Steliga T.; Skalski T.; Kapusta P. Influence of Biosurfactants on the Efficiency of Petroleum Hydrocarbons Biodegradation in Soil. *Sustainability* **2025**, *17*, 6520, doi:10.3390/su17146520.
239. Wu M., Li W., Dick W.A., Ye X., Chen K., Kost D., Chen L. Bioremediation of hydrocarbon degradation in a petroleum-contaminated soil and microbial population and activity determination. *Chemosphere* **2017**, 169, 124–130. https://doi.org/10.1016/j.chemosphere.2016.11.059
240. Xi Y., Song Y., Johnson D.M., Li M., Liu H., Huang Y. Se enhanced phytoremediation of diesel in soil by *Trifolium repens*. *Ecotoxicology and Environmental Safety* **2018**, 154, 137–144. https://doi.org/10.1016/j.ecoenv.2018.01.061
241. Xiang W., Liang Y., Hong S., Wang G., You J., Xue Y., Ma Y. Degradation of long-chain n-alkanes by a novel thermal-tolerant *Rhodococcus* strain. *Archives of Microbiology* **2022**, 204, 259. https://doi.org/10.1007/s00203-022-02872-3
242. Xiao T., Li P., Fei W., Wang J. Effects of vegetation roots on the structure and hydraulic properties of soils: A perspective review. *Science of The Total Environment* **2024**, 906, 167524. https://doi.org/10.1016/j.scitotenv.2023.167524
243. Xiao Y., Jiang R., Wu X., Zhong Q., Li Y., Wang H. Comparative Genomic Analysis of *Stenotrophomonas maltophilia* Strain W18 Reveals Its Adaptative Genomic Features for Degrading Polycyclic Aromatic Hydrocarbons. *Microbiology Spectrum* **2021**, 9, e01420-21. https://doi.org/10.1128/Spectrum.01420-21
244. Xu R., Lau N.L.A., Ng K.L., Obbard J.P. Application of a Slow-Release Fertilizer for Oil Bioremediation in Beach Sediment. *Journal of Environmental Quality* **2004**, 33, 1210–1216. https://doi.org/10.2134/jeq2004.1210
245. Yang K.M., Poolpak T., Pokethitiyook P. Rhizodegradation: The Plant Root Exudate and Microbial Community Relationship, in: Newman, L., Ansari, A.A., Gill, S.S., Naeem, M., Gill, R. (Eds.), Phytoremediation. Springer International Publishing, Cham, **2023**, 209–229. https://doi.org/10.1007/978-3-031-17988-4_11
246. Yemele O.M., Zhao Z., Nkoh J.N., Ymele E., Usman M. A systematic review of polycyclic aromatic hydrocarbon pollution: A combined bibliometric and mechanistic analysis of research trend toward an environmentally friendly solution. *Science of The Total Environment* **2024**, 926, 171577. https://doi.org/10.1016/j.scitotenv.2024.171577
247. Yousaf S., Andria V., Reichenauer T.G., Smalla K., Sessitsch A. Phylogenetic and functional diversity of alkane degrading bacteria associated with Italian ryegrass (*Lolium multiflorum*) and Birdsfoot trefoil (*Lotus corniculatus*) in a petroleum oil-contaminated environment. *Journal of Hazardous Materials* **2010**, 184, 523–532. https://doi.org/10.1016/j.jhazmat.2010.08.067
248. Yu K.S.H., Wong A.H.Y., Yau K.W.Y., Wong Y.S., Tam N.F.Y. Natural attenuation, biostimulation and bioaugmentation on biodegradation of polycyclic aromatic hydrocarbons (PAHs) in mangrove sediments. *Marine Pollution Bulletin* **2005**, 51, 1071–1077. https://doi.org/10.1016/j.marpolbul.2005.06.006

249. Zajic J.E., Seffens W., Panchal C. Biosurfactants. *Critical Reviews in Biotechnology* **1983,** 1, 87–107. https://doi.org/10.3109/07388558309082580
250. Zhang L., Zhang C., Cheng Z., Yao Y., Chen J. Biodegradation of benzene, toluene, ethylbenzene, and o-xylene by the bacterium *Mycobacterium cosmeticum* byf-4. *Chemosphere* **2013**, 90, 1340–1347. https://doi.org/10.1016/j.chemosphere.2012.06.043
251. Zhang X., Li L., Shi X., Chen S., Liang W., Zhu,Y., Li H. Influence of Thermal Desorption Technology on Removal Effects and Properties of PAH-Contaminated Soil Based on Engineering Experiments. *Agronomy* **2024**, 14, 1117. https://doi.org/10.3390/agronomy14061117
252. Zhang Y., Liu H., Yu S., Pei R., Hu H., Wang W., Xu P., Tang H. Characterization of C16–C36 alkane degradation and oily sludge bioremediation by *Rhodococcus erythropolis* XP. *Applied and Environmental Microbiology* **2025**, 91, e02124-25. https://doi.org/10.1128/aem.02124-25
253. Zhou E., Crawford R.L. Effects of oxygen, nitrogen, and temperature on gasoline biodegradation in soil. *Biodegradation* **1995**, 6, 127–140. https://doi.org/10.1007/BF00695343
254. Zhou X., Zhang B., Meng Q., Li L. Effects of Graphene Oxide on Endophytic Bacteria Population Characteristics in Plants from Soils Contaminated by Polycyclic Aromatic Hydrocarbons. *Molecules* **2024**, 29, 2342. https://doi.org/10.3390/molecules29102342

Streszczenie

Gospodarka odpadami wiertniczymi od kilku lat stanowi jeden z kluczowych problemów środowiskowych, z jakimi mierzy się obecnie branża górnictwa nafty i gazu. Odpady te przez wiele lat były sukcesywnie deponowane w tzw. dołach urobkowych, co doprowadziło do trwałego skażenia gruntów na tych terenach oraz konieczność przeprowadzenia działań rekultywacyjnych. Bioremediacja jest obecnie jedną z najczęściej wykorzystywanych metod oczyszczania terenów historycznie zanieczyszczonych węglowodorami ropopochodnymi, co wynika z jej wysokiej skuteczności, relatywnie niskiego oddziaływania środowiskowego oraz stosunkowo niskich kosztów prowadzenia procesu. Jednocześnie obserwuje się, że w przypadku gruntów silnie zdegradowanych, klasyczne techniki bioremediacyjne mogą wykazywać ograniczoną efektywność procesu, dlatego w niniejszej pracy doktorskiej podjęto próbę modyfikacji i opracowania zintegrowanych strategii bioremediacycjnych umożliwiających głębsze doczyszczenie gleb z terenów dołów urobkowych.

Niniejsza rozprawa doktorska została opracowana na podstawie siedmiu spójnych tematycznie publikacji naukowych, w których zaproponowano różne możliwości modyfikacji istniejących technik bioremediacyjnych obejmujących zarówno modyfikację techniki inokulacji gleby biopreparatem opracowanym na bazie bakterii autochtonicznych, jak i zastosowanie tzw. strategii zintegrowanych.

W części literaturowej omówiono zagadnienia związane z problemem skażenia środowiska substancjami ropopochodnymi, charakterystykę węglowodorów ropopochodnych oraz metodami oczyszczania gleb zanieczyszczonych tego typu substancjami, ze szczególnym uwzględnieniem biodegradacji oraz metodami wspomagającymi biodegradację zanieczyszczeń ropopochodnych. Ważnym aspektem poruszonym w tej części pracy było zwrócenie uwagi na szczególny charakter gleb pochodzących z terenów dołów urobkowych, które stanowiły materiał badawczy w artykułach omówionych w niniejszej rozprawie doktorskiej.

W części doświadczalnej omówiono materiał badawczy oraz metodyki badawcze wykorzystywane do oceny efektywności zaproponowanych modyfikacji procesu bioremediacji gleb z terenów dołów urobkowych, a także szczegółowo omówiono wyniki badań zaprezentowanych w artykułach I-VII, stanowiących podstawę niniejszej rozprawy doktorskiej. Ważnym aspektem niniejszej pracy była kompleksowość badań zaprezentowanych w załączonych publikacjach naukowych, obejmujących wykorzystanie analiz respirometrycznych, chromatograficznych, mikrobiologicznych oraz toksykologicznych, co umożliwiło pełną ocenę efektywności zaproponowanych modyfikacji procesów bioremediacyjnych, a także możliwość wykorzystania uzyskanych wyników w przyszłości, w warunkach przemysłowych.

W poszczególnych artykułach omówionych w pracy zaprezentowano nowe podejścia prowadzenia procesu bioremediacji wspomaganej bioaugmentację ukierunkowane na poprawę efektywności biodegradacji zanieczyszczeń ropopochodnych w glebach z dołów urobkowych. W artykule I zaproponowano możliwość modyfikacji techniki inokulacji biopreparatem opracowanym na bazie bakterii autochtonicznych poprzez zastosowanie γ-PGA, a efekty tych badań zostały opublikowane w czasopiśmie

Materials (Wojtowicz et al., 2022). Wyniki uzyskane w I artykule stanowiły podstawę do rozwijania koncepcji modyfikacji techniki inokulacji biopreparatem poprzez wzbogacenie jej w inne substancje wpływające na poprawę biodostępności zanieczyszczeń takie jak biosurfaktanty. Kompleksowe laboratoryjne badania biodegradacji z wykorzystaniem zmodyfikowanej techniki inokulacji biopreparatem poprzez wprowadzenie do układu reakcyjnego biosurfaktantów umożliwiły ocenę ich wpływu na biodegradację węglowodorów alifatycznych oraz WWA w glebie z dołu urobkowego i zostały opublikowane w czasopiśmie *Sustainability* (Wojtowicz et al., 2025).

Kolejnym etapem badań była modyfikacja biopreparatu opracowanego na bazie bakterii autochtonicznych w wybrane szczepy grzybów autochtonicznych oraz wprowadzenie fitoremediacji z wykorzystaniem roślin uprawnych *Zea mays* i ozdobnych *Echinacea purpurea*, w celu głębszego doczyszczenia gleb z terenów dołów urobkowych. Bazując na wynikach uzyskanych w artykułach I i II, zbadano również możliwość połączenia bioaugmentacji, fitoremediacji i γ-PGA, wykazując przewagę tzw. metod zintegrowanych nad metodami klasycznymi w poprawie efektywności oczyszczania gleb historycznie zanieczyszczonych węglowodorami ropopochodnymi. Wyniki tych badań zostały opublikowane w czasopiśmie *Molecules* (Wojtowicz et al., 2023b) i *Applied Science* (Wojtowicz et al., 2023a).

W artykule V rozwijano możliwość wykorzystania techniki fitoremediacji wspomaganej bioaugmentacją, jednakże w badaniach postanowiono wykorzystać dziką roślinność z gatunków *Scirpus sylvaticus* oraz *Cirsium oleraceum*, pobraną z terenów skażonych ropą naftową zlokalizowanych w sąsiedztwie dołu urobkowego, z którego pobrano glebę do badań. Wykorzystanie dzikiej roślinności wykazującej wysoka tolerancję na wysokie stężenie zanieczyszczeń ropopochodnych w glebie skutkowało istotna poprawą efektywności oczyszczania gleby z dołu urobkowego. Ponadto w artykule przeprowadzono charakterystykę społeczności bakteryjnej oraz orzęskowej w glebie po zakończeniu eksperymentu, co stanowiło rozszerzenie zakresu badań omawianych w pozostałych artykułach. Wyniki tych prac zostały opublikowane w czasopiśmie *International Biodeterioration & Biodegradation* (Wojtowicz et al., 2026a).

W artykule VI zaproponowano nowe podejście do poprawy efektywności biodegradacji zanieczyszczeń ropopochodnych, polegające na połączenie bioaugmentacji z wzbogaceniem gleby w wybrane nanomateriały. Badania biodegradacji prowadzone w skali laboratoryjnej wykazały wysoką skuteczność zaproponowanej modyfikacji procesu bioremediacji gleby z dołu urobkowego, w związku z czym rozwijano ten koncept poprzez wprowadzenie fitoremediacji z wykorzystaniem trawy *Lollium perenne,* a wyniki tych badań zostały zaprezentowane w czasopiśmie *Molecules* (Wojtowicz et al., 2026b).

W ostatnim z omówionych w pracy doktorskiej artykule, zbadano wpływ zanieczyszczeń współwystępujących takich jak PCB na efektywność biodegradacji zanieczyszczeń ropopochodnych w glebie z dołu urobkowego. Wykazano skuteczność biopreparatów opracowanych na bazie kilku szczepów bakteryjnych o różnym spektrum działania, w poprawie efektywności oczyszczania gleb skażonych różnymi rodzajami zanieczyszczeń. Wyniki tych badań zostały opublikowane w czasopiśmie *Molecules* (Steliga et al., 2020).

Kompleksowe badania prowadzone w skali laboratoryjnej i półtechniczne zaproponowanych technik modyfikacji procesów bioremediacyjnych omówionych w artykułach I-VII pozwoliły na zrealizowanie wszystkich założonych celów niniejszej

pracy doktorskiej oraz sformułowanie najważniejszych wniosków. Wykazano, że zaproponowane modyfikacje istniejących technik bioremediacyjnych oraz zastosowanie technik zintegrowanych istotnie przyczyniło się do poprawy efektywności bioremediacji gleb z terenów dołów urobkowych i wykazują potencjał do wykorzystania ich w przyszłości w warunkach terenowych.

Abstract

The management of drilling waste has become one of the major environmental challenges facing the oil and gas industry. For many years, drilling waste was systematically deposited in so-called waste pits, resulting in the long-term contamination of soils in these areas and creating the need for remediation measures. Bioremediation is currently one of the most widely applied methods for the remediation of sites historically contaminated with petroleum hydrocarbons due to its high efficiency, relatively low environmental impact, and comparatively low operating costs. However, it has been observed that conventional bioremediation techniques may exhibit limited effectiveness in heavily degraded soils. Therefore, this doctoral thesis aimed to modify existing bioremediation techniques and develop integrated bioremediation strategies to achieve more thorough remediation of soils from waste pit areas.

This doctoral thesis is based on seven thematically coherent scientific publications presenting various approaches to the modification of existing bioremediation techniques, including both the modification of soil inoculation using a biopreparation developed on the basis of autochthonous bacteria and the application of integrated remediation strategies.

The literature review discusses issues related to environmental contamination with petroleum-derived compounds, the characteristics of petroleum hydrocarbons, and methods for the remediation of soils contaminated with these substances, with particular emphasis on biodegradation and techniques supporting the biodegradation of petroleum contaminants. Particular attention was paid to the specific characteristics of soils originating from waste pit areas, which constituted the research material used in the studies included in this doctoral thesis.

The experimental section presents the research material and methodologies used to evaluate the effectiveness of the proposed modifications to the bioremediation of waste pit soils and provides a detailed discussion of the results presented in Articles I–VII, which constitute the basis of this doctoral thesis. An important aspect of this work is the comprehensive nature of the research presented in the included scientific publications, encompassing respirometric, chromatographic, microbiological, and toxicological analyses. This multidisciplinary approach enabled a comprehensive assessment of the effectiveness of the proposed modifications to bioremediation processes and demonstrated the potential applicability of the obtained results under future industrial conditions.

The individual articles included in this doctoral thesis present novel bioaugmentation-assisted bioremediation approaches aimed at improving the biodegradation efficiency of petroleum-derived contaminants in waste pit soils. Article I presents a modification of the biopreparation inoculation technique based on

autochthonous bacteria through the application of γ-PGA. The results of this study were published in the journal *Materials* (Wojtowicz et al., 2022).

The findings reported in Article I provided the basis for the further development of the biopreparation inoculation strategy by enriching it with additional compounds enhancing contaminant bioavailability, such as biosurfactants. Comprehensive laboratory biodegradation studies employing the modified inoculation technique supplemented with biosurfactants enabled the assessment of their effects on the biodegradation of aliphatic hydrocarbons and polycyclic aromatic hydrocarbons (PAHs) in waste pit soil. The results of these studies were published in the journal *Sustainability* (Wojtowicz et al., 2025).

The next stage of the research involved modifying the biopreparation based on autochthonous bacteria by incorporating selected strains of autochthonous fungi. In addition, phytoremediation using the crop plant *Zea mays* and the ornamental plant *Echinacea purpurea* was introduced to achieve more thorough remediation of waste pit soils. Based on the results obtained in Articles I and II, the possibility of combining bioaugmentation, phytoremediation, and γ-PGA was also investigated, demonstrating the superiority of integrated approaches over conventional methods in enhancing the remediation efficiency of soils historically contaminated with petroleum hydrocarbons. The results of these studies were published in the journals *Molecules* (Wojtowicz et al., 2023b) and *Applied Sciences* (Wojtowicz et al., 2023a).

Article V further developed the concept of bioaugmentation-assisted phytoremediation by employing wild plant species, namely *Scirpus sylvaticus* and *Cirsium oleraceum*, collected from crude oil-contaminated areas located in the vicinity of the waste pit from which the soil samples were obtained. The use of wild vegetation exhibiting high tolerance to elevated concentrations of petroleum-derived contaminants significantly improved the efficiency of waste pit soil remediation. Furthermore, the bacterial and ciliate communities present in the soil after completion of the experiment were characterised, thereby extending the scope of the investigations beyond those presented in the remaining articles. The results of these studies were published in the journal *International Biodeterioration & Biodegradation* (Wojtowicz et al., 2026a).

Article VI proposed a novel approach to enhancing the biodegradation efficiency of petroleum-derived contaminants by combining bioaugmentation with soil amendment using selected nanomaterials. Laboratory-scale biodegradation experiments demonstrated the high effectiveness of this modification of the bioremediation process applied to waste pit soil. Consequently, the concept was further developed by incorporating phytoremediation using *Lolium perenne*, and the results of these studies were published in the journal *Molecules* (Wojtowicz et al., 2026b).

The final article included in this doctoral thesis investigated the influence of co-occurring contaminants, such as polychlorinated biphenyls (PCBs), on the biodegradation efficiency of petroleum-derived contaminants in waste pit soil. The study demonstrated the effectiveness of biopreparations developed on the basis of several bacterial strains with different metabolic capabilities in improving the remediation efficiency of soils contaminated with multiple classes of pollutants. The results of this research were published in the journal *Molecules* (Steliga et al., 2020).

The comprehensive laboratory-scale and pilot-scale investigations of the proposed

modifications to bioremediation processes presented in Articles I–VII enabled the successful achievement of all objectives of this doctoral thesis and the formulation of its principal conclusions. The results demonstrated that the proposed modifications of existing bioremediation techniques, together with the application of integrated remediation strategies, significantly enhanced the efficiency of the bioremediation of waste pit soils and exhibit considerable potential for future application under field conditions.

Załączniki

W tym rozdziale zaprezentowane zostały pełne wersje artykułów naukowych opisanych skrótowo w rozdziałach 3-8. Artykuły zostały wymienione w następującej kolejności:

Artykuł I: **Katarzyna Wojtowicz**, T. Steliga, P., Kapusta, J. Brzeszcz, T. Skalski, *Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil*, Materials, Volume 15(2), 400, 2022. DOI: 10.3390/ma15020400

Artykuł II: **Katarzyna Wojtowicz**, T. Steliga, T. Skalski, P., Kapusta, *Influence of Biosurfactants on the Efficiency of Petroleum Hydrocarbons Biodegradation in Soil*, Sustainability, Volume 17(14), 6520, 2025. DOI: 10.3390/su17146520

Artykuł III: **Katarzyna Wojtowicz**, T. Steliga, P., Kapusta, J. Brzeszcz, *Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (Zea mays)*, Molecules, Volume 28(16), 6104, 2023. DOI: 10.3390/molecules28166104

Artykuł IV: **Katarzyna Wojtowicz**, T. Steliga, P., Kapusta, *Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using Echinacea purpurea*, Applied Sciences (Basel), Volume 13(24), 13077, 2023. DOI: 10.3390/app132413077

Artykuł V: **Katarzyna Wojtowicz**, T. Steliga, J. Brzeszcz, J. Fyda, T. Skalski, P., Kapusta, *Phytoremediation of soil contaminated with petroleum hydrocarbons using the wild plants Scirpus sylvaticus and Cirsium oleraceum, supported by bioaugmentation*, International Biodeterioration & Biodegradation, Volume 206, 106217, 2026. DOI: 10.1016/j.ibiod.2025.106217

Artykuł VI: **Katarzyna Wojtowicz**, T. Steliga, P., Kapusta, J. Brzeszcz, *The Role of Graphene Oxide and Zinc Oxide Nanoparticles in Enhancing the Effectiveness of Phytoremediation of Petroleum Hydrocarbon-Contaminated Soils Using Lolium perenne*, Molecules, Volume 31(5), 890, 2026. DOI: 10.3390/molecules31050890

Artykuł VII I: T. Steliga, **Katarzyna Wojtowicz**, P., Kapusta, J. Brzeszcz, *Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture*, Molecules, Volume 25(3), 709, 2020. DOI: 10.3390/molecules25030709

materials

MDPI

Article

Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil

Katarzyna Wojtowicz [1,*], Teresa Steliga [1], Piotr Kapusta [1], Joanna Brzeszcz [1] and Tomasz Skalski [2]

[1] Oil and Gas Institute—National Research Institute, Lubicz 25 A, 31-503 Krakow, Poland; teresa.steliga@inig.pl (T.S.); piotr.kapusta@inig.pl (P.K.); joanna.brzeszcz@inig.pl (J.B.)
[2] Biotechnology Centre, Silesian University of Technology, Krzywoustego 8, 44-100 Gliwice, Poland; tomasz.skalski@polsl.pl
* Correspondence: katarzyna.wojtowicz@inig.pl

Abstract: Biodegradation is a method of effectively removing petroleum hydrocarbons from the natural environment. This research focuses on the biodegradation of aliphatic hydrocarbons, monoaromatic hydrocarbons such as benzene, toluene, ethylbenzene, and all three xylene isomers (BTEX) and polycyclic aromatic hydrocarbons (PAHs) as a result of soil inoculation with a biopreparation A1 based on autochthonous microorganisms and a biopreparation A1 with the addition of γ-PGA. The research used biopreparation A1 made of the following strains: *Dietzia* sp. *IN133*, *Gordonia* sp. *IN138 Mycolicibacterium frederiksbergense IN53*, *Rhodococcus erythropolis IN119*, *Rhodococcus* sp. *IN136* and *Pseudomonas* sp. *IN132*. The experiments were carried out in laboratory conditions (microbiological tests, respirometric tests, and in semi-technical conditions (*ex-situ* prism method). The biodegradation efficiency was assessed on the basis of respirometric tests, chromatographic analyses and toxicological tests. As a result of inoculation of AB soil with the biopreparation A1 within 6 months, a reduction of total petroleum hydrocarbons (TPH) (66.03%), BTEX (80.08%) and PAHs (38.86%) was achieved and its toxicity was reduced. Inoculation of AB soil with the biopreparation A1 with the addition of γ-PGA reduced the concentration of TPH, BTEX and PAHs by 79.21%, 90.19%, and 51.18%, respectively, and reduced its toxicity. The conducted research has shown that the addition of γ-PGA affects the efficiency of the biodegradation process of petroleum pollutants, increasing the degree of TPH biodegradation by 13.18%, BTEX by 10.11% and PAHs by 12.32% compared to pure biopreparation A1.

Keywords: biodegradation; petroleum contaminants; biopreparation; inoculation; polyglutamic acid; ecotoxicology

check for updates

Citation: Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J.; Skalski, T. Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil. *Materials* **2022**, *15*, 400. https://doi.org/10.3390/ma15020400

Academic Editor: Camelia Ungureanu

Received: 17 December 2021
Accepted: 4 January 2022
Published: 6 January 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

Oil mining exploits valuable, non-renewable resources of natural gas and crude oil, and also contributes to the degradation of the biologically active surface of the earth. The works carried out in the oil and gas mining are accompanied by the necessity to collect and manage large amounts of waste materials, which were stored in waste pits in the years 1920–1950 [1]. In terms of chemicals, these wastes are diverse, multi-component systems that changed during deposition and initial treatment processes. An important parameter determining the changes taking place in the deposited waste is the presence of petroleum pollutants, which age as a result of the course of many processes, including evaporation, dissolution, transport of water-soluble substances, and biodegradation. The neutralization of old petroleum pollutants in the areas of liquidated mines and in the areas of old waste pits is one of the key ecological problems faced by the oil and gas mining industry.

One of the main pollutants causing the inhibition of the development and growth of flora in these areas are petroleum hydrocarbons (TPH, BTEX, PAHs), the concentration

of which in old soil varies widely and varies within wide ranges. Individual terrestrial ecosystems, constituting an ecological unit which includes groups of living organisms, form a certain whole adapted to the conditions in the biotope. Introducing petroleum pollutants to ecosystems disturbs the course of the natural cycles of matter and energy circulation. The stimulation of the development of some groups of microorganisms and the inhibition of the activity of the remaining ones are observed. In addition, petroleum pollution causes soil distortions, leading to the reduction of water capacity due to partial or complete filling of soil pores, impeding or preventing the exchange of air between the soil and the atmosphere. The excess of organic carbon in relation to nitrogen and phosphorus content causes an acute shortage of these components for microorganisms and plants, degradation of the physicochemical properties of soil colloids (sorption capacity, ion exchange), and causes visible and unfavorable changes in the pH of the environment [2,3].

The elimination of hazardous, toxic and mutagenic substances from the environment (soil) is one of the main goals of environmental engineering. There are many techniques for remediating soil contaminated with petroleum substances, including thermal (thermal combustion), physical-chemical (soil washing, solidification, vacuum desorption, photolysis, chemical degradation, electrokinetic method) and biological (biodegradation, phytoremediation) methods. One of the most frequently used methods of treatment of soil contaminated with petroleum substances is biodegradation. Biodegradation is a much more economical method (no need to use complicated apparatus), safe and environmentally friendly than conventional thermal and physico-chemical methods. It is characterized by high efficiency and a wide spectrum of activity thanks to the use of agricultural microorganisms to decompose various types of pollutants. On the other hand, not all chemicals can be removed from the environment by biodegradation, and the time of land remediation by biological methods may be longer than by thermal or physico-chemical methods. Biodegradation can be carried out *in-situ*, i.e., in the place of contamination, or *ex-situ*, i.e., outside the place where contamination occurs. The advantage of conducting the biodegradation process with the *in-situ* method is the possibility of cleaning land in urbanized areas, while the *ex-situ* method enables the intensification of the contaminated soil remediation processes by controlling the process conditions and removing leachate containing harmful substances.

The biodegradation of petroleum hydrocarbons is a complex process depending on many factors that can be divided into 3 main groups: factors related to pollution (xenobiotic concentration [4,5], age of pollution [6,7], physico-chemical properties [6,8–10], symmetry and particle size [11,12]), environmental factors [4] (temperature [13–15], pH of the environment [4], the presence of oxygen [4,6,9,16–18], humidity [4,6], the presence of nutrients [4,18–20], properties of the soil matrix [6,17,21]) and biological factors (quantitative and qualitative composition of microorganisms, bioavailability of the pollutant for organisms supporting decomposition, spatial distribution, the adaptability of microorganisms, interactions between species and within the same species, enzymatic activity [4,22–27]).

The ability of microorganisms to adapt to pollutants plays a particularly important role in the decomposition of xenobiotics. At the molecular level, adaptation is explained by the genetic variation of microorganisms (resulting from the presence of the relevant genes), and then the selection of those cells that have acquired the ability to use xenobiotics as a source of carbon and energy. Petroleum pollutants constitute a complex, multi-component system and differ in their susceptibility to microbial attack. The susceptibility of hydrocarbons to microbial degradation can be broadly classified as follows: linear alkanes > branched alkanes > monoaromatic hydrocarbons (BTEX) > cyclic alkanes > polycyclic aromatic hydrocarbons [28]. There are several hundred species of microorganisms capable of biodegradation of petroleum pollutants, but most of them can metabolize a limited range of hydrocarbons [4]. For this reason, for their biodegradation, it is advisable to use a mixture of microorganism cultures with an extensive enzyme apparatus [29,30]. It is preferable to prepare bacterial consortia—biopreparations based on autochthonous microorganisms previously isolated from the soil in order to avoid the antagonistic influence of the autochthonous soil microbiota on foreign cultures, which creates a high probability of survival

in a contaminated biocenosis, and thus guaran—tees a sufficiently high biodegradation potential. The most commonly reported bacteria in the literature capable of degrading both aliphatic and aromatic hydrocarbons the following types: *Achromobacter* [31], *Acinetobacter* [32], *Bacillus* [33–36], *Dietzia* [37], *Gordonia* [38–40], *Micrococcus* [41,42], *Mycobacterium/Mycolicibacterium* [43–51], *Nocardia* [52,53], *Pseudomonas* [54–58], *Rhizobium* [59,60], *Rhodococcus* [44,61–66], *Sphingomonas* [17,67–70], *Stenotrophomonas* [71–73] and *Sphingobacterium* [34,74,75]. A combination of bioaugmentation treatments (inoculation of the contaminated soil-water environment with microorganisms characterized by high biodegradation efficiency of petroleum pollutants, including TPH, BTEX and PAHs) and biostimulation (introduction of oxygen, water, biogenic substances and surface-active substances to the soil-water environment in order to increase the activity of microorganisms autochthonous species) allows for a significant improvement in the efficiency of the biodegradation process of petroleum hydrocarbons.

Currently, more and more research is being carried out on accelerating the biodegradation process of petroleum pollutants through biotechnological processes with the use of a bacterial consortium (biopreparations) prepared on the basis of autochthonous bacteria and additives accelerating the decomposition of hydrocarbons. An interesting solution seems to be the use of a biopreparation in the inoculation process in combination with biopolimers, e.g., γ-PGA.

Biopolymers—polymers produced by living organisms, fulfilling specific functions in these organisms. A number of biopolymers have an industrial application; in the oil and gas industry, polysaccharide biopolymers such as starch, cellulose and guar gum and their derivatives are used, and to a lesser extent xanthan gum, scleroglucan and chitin/chitosan [76]. Biopolymers that are not polysaccharide compounds have not aroused interest so far. However, in recent years, the attention of a number of researchers has focused on a compound of a polyamide nature—γ-polyglutamic acid (γ-PGA) [77]. A characteristic feature of γ-PGA is that it is made of one type of monomer—glutamic acid. The monomers in γ-PGA are linked by an amide bond between the α-amino group of one monomer and the α-carboxyl group of the other. γ-PGA is a product of bacteria and archaea, and only a few species have the ability to produce it [78,79]. Particularly preferred is the genus Bacillus, because most of the γ-PGA-synthesizing organisms belong to this genus (*Bacillus amyloliquefaciens*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus subtilis*, as well as the highly pathogenic organism—*Bacillus anthracis*) [80]. Particular species and even strains produce γ-PGA with different molecular weights; from 100 kD to 5000 kD [81,82], which is reflected in its physical properties. The higher the molecular weight, the greater the density of the aqueous γ-PGA solution. Depending on the environmental conditions, γ-PGA can assume various conformational states, e.g., in lower pH it will be in the form of an α-helix, and in higher pH in the form of a random coil [83]. On the other hand, increased temperature causes the aggregation of γ-PGA molecules and the formation of fibrous structures [84]. PGA can be produced from a variety of precursors. In general, γ-PGA-synthesizing microorganisms fall into 2 groups. The first one is strictly dependent on the presence of L-glutamic acid in the medium, the second one can use various precursors, such as citric acid, glucose, α-ketoglutaric acid, glycerol and various amino acids, e.g., alanine, arginine, ornithine, proline and aspartic acid, in particular glutamine. By growing the bacteria from the above-mentioned compounds in various combinations, γ-PGAs of various molecular weights can be obtained and the efficiency of the process can be controlled [77].

The changing properties of γ-PGA (depending on the environment in which it is located) have attracted the attention of many researchers in terms of the industrial application of this biopolymer. Moreover, the compound has many advantages: it is highly soluble in water, non-toxic to living organisms even in very high concentrations, biodegradable (although mainly by organisms capable of synthesizing it), and even edible [85]. Therefore, it is currently used as a flocculant [86,87], an additive to fertilizers [88] and food [89]. Its presence in human food has been known for a relatively long time—traditional Japanese natto (based on fermented soybeans) contains a lot of γ-PGA, and now it is suggested that

γ-PGA has a positive effect on lowering sugar levels, and therefore has a health-promoting effect [90]. Other features—biocompatibility and lack of immunogenicity—allow it to be used as a medical and cosmetic agent [91]. The use of γ-PGA in environmental engineering has also begun to be considered. Favorable results have been obtained in the application of γ-PGA in the remediation of soils contaminated with heavy metals [92–94]. γ-PGA has also been reported to have a positive effect on the removal of crude oil from marine sediments [95] and trichlorethylene (TCE) from the water bearing zone [96]. Therefore, we decided to experimentally check whether the presence of γ-PGA would support the decomposition of petroleum substances (TPH, BTEX, PAHs) in soils inoculated with a biopreparation developed on the basis of bacterial microorganisms.

We hypothesized that it was possible to increase the efficiency of biodegradation of petroleum pollutants by using a biopreparation based on autochthonous bacteria with the addition of γ-PGA in the inoculation process. The aim of the research was: (a) to determine the efficiency of the biodegradation process of petroleum pollutants (TPH, BTEX, PAHs) during inoculation with a biopreparation prepared on the basis of autochthonous bacteria isolated, multiplied and selected from the treated soil, carried out for a period of 6 months on a semi-technical scale using the *ex-situ* prism method, (b) determining the degree of increasing the efficiency of biodegradation of petroleum pollutants during the application of the PGA additive in the inoculation process as a function of the duration of biodegradation processes, (c) toxicological monitoring of soil as a function of time during soil biodegradation processes together with the sensitivity assessment of the biotests applied. The efficiency of biodegradation of petroleum substances was assessed on the basis of respirometric tests, chromatographic analyses and toxicological tests with the use of a new generation toxicological test set, the bioindicators of which belonged to different taxonomic groups (bacteria, crustaceans and plants).

2. Materials and Methods

2.1. Soil and Microorganisms

In the study of the biodegradation of petroleum substances, the AB soil (initial soil) collected from the waste pit G11 (N: 49°40′31.23″, E: 22°04′30.86″) in the area of the crude oil mine located in south-eastern Poland was used. This soil is characterized by a high concentration of petroleum pollutants, i.e., aliphatic hydrocarbons (19,774.23 mg/kg DM), monoaromatic hydrocarbons (17.35 mg/kg DM) and polycyclic aromatic hydrocarbons (27.03 mg/kg DM). The content (%) of individual components of petroleum pollutants in the soil is provided in the Supplementary Materials on Figure S1.

For soil inoculation, biopreparation A1 was used, developed on the basis of bacterial strains (came from the hydrocarbon-degrading microbial collection of the Department of Microbiology at the Oil and Gas Institute–National Research Institute, Krakow, Poland), isolated formerly from the AB soil: *Dietzia* sp. *IN133*, *Gordonia* sp. *IN138 Mycolicibacterium frederiksbergense IN53*, *Rhodococcus erythropolis IN119*, *Rhodococcus* sp. *IN136* and *Pseudomonas* sp. *IN132*. A commercially available preparation, Ambiogel® (Ambioteco, Staszów, Poland), containing 10% (*w/w*) pure gamma-polyglutamic acid, was used as a γ-PGA additive. Table 1 presents the species affiliation of bacterial strains occurring in the biopreparation A1.

All reagents used for the cultivation of microorganisms came from Avantor Performance Materials Poland S.A. (former POCH Gliwice) (Gliwice, Poland) and were of analytical purity with the exception of nutrient broth which were purchased from Biomaxima S.A. (Lublin, Poland).

Table 1. Genes involve in petroleum hydrocarbon degradation, found in genomes for the closest relatives of all strains.

Stain	The Closest Relative Based on 16S rRNA Accession Number, (% of Identity) *	The Closest Relative for Which the Genome Sequence Is Available in NCBI GenBank, Accession Number, (% of Identity) *	Putative Gene Encoding for the Enzymes Degrading Hydrocarbons
Dietzia sp. *IN133*	*Dietzia* sp. JS16-p6b CP024869 99.93%	*Dietzia* sp. JS16-p6b CP024869 99.93%	Alkane 1-monooxygenase (4 copies), biphenyl 2,3-dioxygenase, (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Gordonia sp. *IN138*	*Gordonia terrae* RL-JC02 CP049836 99.65%	*Gordonia terrae* RL-JC02 CP049836 99.65%	Alkane 1-monooxygenase, pentachlorophenol monooxygenase, naphthalene 1,2-dioxygenase subunit alpha (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Mycolicibacterium frederiksbergense IN53	*Mycolicibacterium frederiksbergense* DSM 44346 (typical strain) NR_025393.1 99.58%	*Mycolicibacterium frederiksbergense* LB 501T 99.8%	Alkane 1-monooxygenase (2 copies), pentachlorophenol monooxygenase, naphthalene 1,2-dioxygenase subunit alpha (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Rhodococcus erythropolis IN119	*Rhodococcus erythropolis* KD-1 CP050124 99.42%	*Rhodococcus erythropolis* KD-1 CP050124 99.42%	Alkane 1-monooxygenase (5 copies), pentachlorophenol monooxygenase, cyclohexanone monooxygenase (2 copies), biphenyl 2,3-dioxygenase (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Rhodococcus sp. *IN136*	*Rhodococcus fascians* KHO60 MN099376 99.71%	*Rhodococcus fascians* D188 CP015235 99.71%	Alkane 1-monooxygenase (3 copies), methane monooxygenase 2,4-dichlorophenol 6-monooxygenase, 1 element of 1,2-benzene dioxygenase system, 2,3-dihydroxybiphenyl 1,2-dioxygenase
Pseudomonas sp. *IN132*	*Pseudomonas citronellolis* PY-1 MH685460 99.49%	*Pseudomonas citronellolis* SJTE-3 CP015878 99.49%	Alkane 1-monooxygenase, phenol 2-monooxygenase, 2,4-dichlorophenol 6-monooxygenase, naphthalene 1,2-dioxygenase, biphenyl 2,3-dioxygenase, 2,3-dihydroxybiphenyl 1,2-dioxygenase

* the presented data concerns the data gathered in the NCBI GenBank database (November 2021).

2.2. Biopreparation Development

The stains were grown separately in nutrient broth supplemented with the addition of sodium acetate (0.2%, *w/v*) and incubated at 25 °C at 150 rpm for 72 h to obtain a cell number of 10^8–10^9 cfu/cm^3. The biopreparation used for further research was prepared by mixing equal volumes of all strains. The ability to use crude oil and various hydrocarbons as the sole source of carbon and energy was tested by incubating each of the strains in a mineral medium with the addition of particular compounds. The test medium was described in the study of Brzeszcz et al., 2020 [97,98]. Simultaneously, tests were carried out according to the method described by Wrenn and Venos (1986) [99]. The characteristics

of the individual strains were established on the basis of microscopic observations, colony morphology, growth on the selective agar media and biochemical profiles (API Coryne and 32GN bioMerieux tests). The strains were identified phylogenetically by sequencing the 16S rRNA gene as previously described [100]. The obtained sequences were deposited in the NCBI database, and their accession numbers together with the properties of the strains are presented in Table 2.

Table 2. Hydrocarbon-degrading capabilities of bacterial strains comprising the biopreparation A1.

Stain	NCBI Accession Number	$nC_{18}H_{38}$	iso $C_{19}H_{40}$	TOL + XYL	NAP	ANTH	PHEN	FLU	FLTH	PYR
Dietzia sp. *IN133*	KT923300	+	+	+	+	−	−	−	−	−
Gordonia sp. *IN138*	KT923297	+	+	+	+	+/−	−	−	−	−
Mycolicibacterium frederiksbergense IN53	JN572675	+	+	−	+	+	+	−	−	−
Rhodococcus erythropolis IN119	KT923331	+	+	+	+	−	−	−	−	−
Rhodococcus sp. *IN136*	KT923330	+	+	−	+	−	−	−	+	−
Pseudomonas sp. *IN132*	KT923299	+	+	+	+	+/−	+	+/−	+/−	+/−

$nC_{18}H_{38}$: n-octane iso-$C_{19}H_{40}$: pristane, TOL: toluene, XYL: mixture of xylenes NAP: naphtalene, ANTH: anthracene, PHEN: phenanthrene FLU: fluorene, FLTH: fluoranthene, PYR: pyrene, +: growth, −: no growth +/−: ambiguous observation.

2.3. Biodegradation Experiments

A workstation for tests carried out on a semi-technical scale in the conditions of *ex-situ* treatment of soil contaminated with petroleum substances was created in the Oil and Gas Institute—National Research Institute, Krosno Division. The structure of the test workstation was as follows: a foil was placed directly on the floor to protect the surface against the spread of contamination of the room, for which a gravel bed of appropriate granulation was used as the first layer. Perforated pipes were laid on the gravel layer, through which compressed air was supplied from the compressor unit. Then the pipes were covered with another layer of gravel, secured with a technical fabric to prevent clogging. On the base prepared in this way, two identical prisms of AB soil were placed in the amount of 30 kg (per 1 prism) (Figure 1).

Before the soil inoculation process, the proper C:N:P ratio was ensured, which should be at the level of 100:10:1 [36,44,100,101] in order to ensure optimal conditions for the proper activity of microorganisms contained in the biopreparation. This was carried out by supplementing nitrogen and phosphorus with the Azofoska mineral fertilizer with the following composition: 13.6% total N, 8.1% ammoniacal nitrogen, 5.5% nitric nitrogen, 6.4% soluble P_2O_5, 19.1% K_2O in the form of K_2SO_4, 4.4% MgO in the form of soluble $MgSO_4$, and microelements (0.17% Fe, 0.27% Mn, 0.18% Cu, 0.045% Zn, 0.09% Mo). Nutrients (nitrogen and phosphorous–present in the soil as ions: NH^{4+}-N, NO_3-N, PO_4-P) were determined using the Perkin Elmer Lambda 40 spectrophotometer (Waltham, MA, USA). The soil pH was corrected by supplementing lime in the amount of 1.0–1.5 g/kg of soil until the optimum pH of 7.5–7.6 was obtained [44]. The prepared AB soil was inoculated with biopreparation A1 (2 dm^3)–prism A, or biopreparation A1 (2 dm^2) with the addition of γ-PGA (2 dm^3 solution Ambiogel®—hereinafter referred to as γ-PGA)—prism B. The prisms were covered with a foil tunnel, which allows to maintain a constant temperature

inside at the level of approx. 20–25 °C, achievable in field conditions in the summer season of a moderate climatic zone and with constant humidity (20–25%). During the treatment process, the content of TPH, BTEX and PAHs in the prism inoculated with the biopreparation A1 (Soil A2, A4, A6) and with the biopreparation A1 with γ-PGA (Soil B2, B4, B6) was monitored, and changes in the level of soil toxicity were determined using the toxicological test package. The process of biodegradation of petroleum hydrocarbons in ex-situ conditions was carried out for 180 days.




Figure 1. Stand for testing the process of biodegradation of petroleum hydrocarbons.

2.4. Respirometric Tests

The respirometric tests consisted in measuring the biological activity in the analyzed soil under aerobic conditions. The research was carried out with the use of the Oxi-Top set (WTW, Weilheim, Germany). Samples weighing 30 g were placed in measuring vessels to which biopreparation A1 (2 cm^3) with a density of 109 cfu mL^{-1} and biopreparation A1 (2 cm^3) with γ-PGA (solution γ-PGA 1 and 2 cm^3) were added. Finally, the humidity of each system was adjusted to 25% by adding distilled water. At the same time, reference samples were prepared in which the analysed soil and inoculant solutions were placed separately. Then all measuring vessels were closed with OxiTop measuring heads, then placed in an incubator and thermostated at 20 °C for 60 days. The Oxi-Top Control system measuring heads read the pressure values in the closed system every two hours. The collected data was transferred via the IR interface to the Oxi-Top OC 110 controller, where it could be processed graphically and statistically using the Achat OC program. The conversion of the measured pressure into the value of consumed oxygen (m_{O_2}) is carried out according to the Formula (1): At the end of the measurement, the data was sent to a PC, where, using Excel, curves of the dependence of the amount of oxygen consumed [mg/dm^3] and the duration of the experiment [days] were prepared. All experiments were performed in triplicate with five replications for each treatment [44,102–104].

$$m_{O_2} = \frac{M(O_2)}{R \times T_m}\left(V_g + \alpha\frac{T_m}{T_0}\right) \times \Delta p \quad (1)$$

where: $M(O_2)$—molar mass of oxygen [kg/mol], V_g—volume of free gas [m^3], R—gas constant [J·mol^{-1}·K^{-1}], T_m—temperature measurement value [K], T_0—reference temperature (273.15 K), α—absorption coefficient (0.03103), Δp—pressure drop in the test [Pa].

2.5. TPH Extraction and Quantification

Isolation of TPH from the soil matrix was carried out using the ultrasound-assisted solvent extraction method. For this purpose, 10 g of air-dry soil sample was placed in an Erlemnajer flask and extracted with 3 portions of solvent (dichloromethane, Avantor Performance Materials Poland S.A., Gliwice, Poland). The soil extraction time was 15 min.

for each portion of the solvent. Extraction was carried out in a Sonoswiss SW 6H ultrasonic bath (Sonoswiss AG, Ramsen, Switzerland) with an ultrasound frequency of 30 kHz [44,105]. The obtained extracts were purified by SPE solid phase extraction on Bakerbond No. 7213-03 columns packed with florisil (J.T.Baker®, Phillipsburg, NJ, USA). The obtained extracts were concentrated on a ChemLand RE100-Pro rotary vacuum (ChemLand, Stargard Szczeciński, Poland evaporator to a volume of 1 cm^3. The efficiency of the ultrasonically assisted solvent extraction was estimated to be 95.9% using the o-terphenyl replacement standard [44,98].

The TPH analysis in soil samples was performed on a Perkin Elmer Clarus 500 GC chromatograph (Waltham, MA, USA) equipped with a flame ionization detector (FID), with the following device parameters: RTX-1 capillary column of fused silica, 30 m × 0.53 mm (Restek, Bellefonte, PA, USA), injector temperature = 290 °C, FID detector temperature = 320 °C, oven temperature programme: 30 °C (2 min. isothermal run), 30–105 °C (temperature increase 10 °C min^{-1}), 105–285 °C (temperature increase 5 °C min^{-1}), and 285 °C (5 min isothermal run), carrier gas (He) flow 20 mL min^{-1} [44]. The total TPH content in soil samples was assessed on the basis of the certified BAM K010 standard (Tusnovic Instruments, Krakow, Poland). Standard mixture paraffin hydrocarbons: (nC_6–nC_{44}) ASTM® D2807 (Supelco, Saint Louis, MO, USA) and certified standard mixture Fuel Oil Degradation Mix n-C_{17}, pristane, n-C_{18}, phytane No A029668: (Restek, Bellefonte, PA, USA) were used to quantitatively the individual n-alkanes included in the petroleum pollutants. A certified standard C_{30} 17α(H), 21β(H)-hopane No 08189 (Sigma-Aldrich, Saint Louis, MO, USA) was used as a biomarker [44].

2.6. BTEX Extraction and Quantification

Headspace gas chromatography was used to determine the content of monoaromatic hydrocarbons (BTEX) in soil [106]. A soil sample with a mass of 6.5 g (as calculated for the dry mass of soil, approx. 5 g) was placed in an ampoule, which was tightly closed with a capping machine and inserted into an autosampler. Before the analysis, the sample was thermostated for 10 min. at 90 °C. The ampoule was filled with a carrier gas (helium) for 3 min, and then a sample of gaseous analytes desorbed to the headspace was collected using a dosing needle heated to the temperature of 95 °C, which was sent via a transfer line (temperature 100 °C) to the injector of the Clarus 500 gas chromatograph with an FID detector, connected to HeadSpace TurboMatrix 16 (Perkin Elmer Inc., Waltham, MA, USA). The injection time was 0.04 min. The operating conditions of the chromatograph were as follows: fused silica capillary column (RT-TCEP: 60 m × 0.25 µm) (Restek, Bellefonte, PA, USA), using the following temperature parameters: injector temperature = 200 °C, FID detector temperature = 280 °C, TCD detector temperature = 150 °C, furnace temperature program: 60 °C—isothermal course for 5 min, 60–100 °C—temperature increase at a rate of 5 °C/min, 100 °C—isothermal course for 10 min. To quantify the total BTEX content, the certified standards of Restek No. 30051 (Restek, Bellefonte, PA, USA) and Supelco No. CRM48026 (Supelco, Saint Louis, MO, USA) [106] were used.

2.7. PAHs Extraction and Quantification

The isolation of analytes from the soil matrix was carried out using the QuEChERS method (Quick, Easy, Cheap, Effective, Rugged, Safe). For this purpose, 5 g of air-dry soil was weighed into a 50 mL vial, and then 5 cm^3 of distilled water and 10 cm^3 of acetonitrile (Avantor Performance Materials Poland S.A., Gliwice, Poland) were added to it. The contents of the sachet with the $MgSO_4$ + NaCl extraction mixture No. SST640 (Interchim, Montluçon, France) were then added to the vial. The samples prepared in this way were shaken for 5 min, and then centrifuged for 10 min at a speed of 3500 rpm [107].

Purification of the extract was performed using the dSPE method using purification vials filled with $MgSO_4$ and PSA No. JO3937 (Interchim, Montluçon, France). 1 cm^3 of the soil extract was transferred to a purification vial, which was then shaken for 5 min and

centrifuged for 10 min at 8000 rpm. The obtained extracts were filtered through a syringe filter with a pore diameter of 25 µm and subjected to chromatographic analysis [107].

The analysis of polycyclic aromatic hydrocarbons (PAHs) in soil samples was performed on a Vanquish Core high pressure liquid chromatograph (HPLC) with the following device parameters: NUCLEODUR C_{18} PAH column, 125 mm × 4 mm, 3 µm (Marcherey-Nagel, Germany), detector: UV-VIS and fluorimetric (FLD) dispenser: automatic, dispensing volume: 10 µL, eluents: A-methanol (Avantor Performance Materials Poland S.A., Gliwice, Poland) 70%, B-acetonitrile (Avantor Performance Materials Poland S.A., Gliwice, Poland), flow: 1.5 mL/min, gradient: 20% B for 1.5 min, 20–50% B in 1.5 min, 50–100% B in 1 min, 100% B for 1 min, 100–0% B for 3 min, 100% A for 3 min. Certified PAH-Mix solution ref. 722393 (Marcherey-Nagel, Düren, Germany) was used to quantify the individual PAHs included in the petroleum pollutants.

2.8. Ecotoxicological Analyses

To evaluate the effectiveness of the biodegradation process of soils contaminated with petroleum substances, a set of toxicological tests used in bioindication analyses was applied. Bioindication analyses are one of the newest methods of environmental toxicity studies in which lethal and sub-lethal effects (morphological changes, diseases) of indicator organisms are observed. During the biodegradation of petroleum hydrocarbons as a result of chemical and microbiological transformations, metabolites of various biological activity may be produced. Toxicological tests quickly provide information on the toxic effects of chemical compounds (both acute and chronic) on a selected indicator organism or population. In this publication, 4 toxicological tests were used to evaluate the effectiveness of the remediation treatments used in the process of soil remediation in ex-situ conditions: Phytotoxkit™ (MicroBioTests Inc., Nazareth, Belgium), Ostracodtoxkit ™ (MicroBioTests Inc., Nazareth, Belgium), Microtox® Solid Phase Test (SDI, Newark, DE, USA) and Ames (Muta-ChromoPlate™ Kit (EBPI, Mississauga, ON, Canada) [44]. The list of toxicological tests used in the research along with the description of the procedures for their performance is presented in Supplementary Materials.

2.9. Mathematical Model of TPH, BTEX and PAHs Biodegradation

In order to work a simplified mathematical model of the biodegradation of petroleum hydrocarbons during *ex-situ* method to normalize the concentrations of TPH analytes (Σ n-C_{10}–n-C_{22} and Σ n-C_{23}–n-C_{40}), BTEX (benzene, toluene, ethylbenzene, xylenes) and PAHs (2-ring PAHs, 3-ring PAHs, 4-ring PAHs, 5-ring PAHs, 6-ring PAHs) the biomarker C3017α (H), 21β (H)-hopan (Supelco, Saint Louis, MO, USA) was used. The biomarker used in biodegradation tests allows for the elimination of analytical errors that may occur during the determination of the analyzed groups of analytes with the chromatographic method. The standardized results of chromatographic determinations performed during biodegradation were used to develop a primary mathematical model describing its course, which was described by Equation (2) [44,100].

$$C/Cx = (C/Cx)0 \exp(-kt) \quad (2)$$

where: C—analyte concentration, Cx—biomarker (Hopane) concentration, k—first-order rate constant [d^{-1}], (C/Cx)0—normalized analyte concentration at the starting point, and t—process duration [d].

The nonlinear regression analysis with the use of Equation (2) made it possible to determine the first-order biodegradation constant (k) and the correlation coefficient (r^2), which determined the fitting of the measuring points to the theoretical curves of the TPH, BTEX and PAHs biodegradation process.

2.10. Data Analysis and Statistical Information

Statistical analysis was performed with Statistica 14.0 (TIBCO Software Inc., Palo Alto, CA, USA). Standard deviation (SD), standard deviation (RSD) (%) and Pearson correlation

coefficient were calculated. All obtained data were previously tested for normal distribution. They were then checked by a one-way ANOVA test. In the case of obtaining significant results in the ANOVA test, Tukey's pairs post-hoc test was performed. The significance was set at $p < 0.05$ [44].

3. Results and Discussion

3.1. Respirometric Tests

Respirometric tests using the OxiTop kit are based on highly accurate, automatic pressure measurements using a piezoresistive electronic pressure sensor (OxiTop Control) in a sealed bottle at a constant temperature. The conducted respirometric tests on control samples (sterile soil not contaminated with hydrocarbons) showed the biological activity of biopreparation A1 and biopreparation A1 with the addition of PGA after 60 days of the test at the level of 526 O_2/dm^3 and 536 O_2/dm^3 (volume ratio of biopreparation A1 and γ-PGA solution was 2:1) and 570 O_2/dm^3 (volume ratio of biopreparation A1 and γ-PGA solution was 1:1), respectively. The oxygen consumption in the AB soil sample, which was not subjected to the inoculation process (Blank sample) after the end of the test, was 62.1 O_2/dm^3. After 60 days of the test, the oxygen consumption in the soil sample inoculated with biopreparation A1 was 1582 mg O_2/dm^3, while in the sample inoculated with biopreparation A1 with the addition of γ-PGA 2040 O_2/dm^3 (volume ratio of biopreparation A1 and γ-PGA solution was 2:1) and 3769 mg O_2/dm^3 (volume ratio of the solutions of biopreparation A1 and γ-PGA solution was 1:1) (Figure 2). The increase in microbiological activity in the reaction environment proves that the microorganisms use carbon contained in aliphatic hydrocarbons, monoaromatic hydrocarbons and polycyclic aromatic hydrocarbons as a food source. The result of the increased biological activity is the increase in oxygen consumption by the examined bacterial consortia over time, which results in the biodegradation of petroleum substances. The obtained results of respirometric tests prove the effectiveness of the biodegradation process of petroleum hydrocarbons with the use of inoculants in the form of biopreparation A1 and biopreparation A1 with the addition of γ-PGA. Based on the conducted respirometric tests, the method of AB soil inoculation with the use of A1 biopreparation A1 and biopreparation A1 with the addition of γ-PGA in the volume ratio of 1:1 solution was selected for further studies of the biodegradation process of petroleum hydrocarbons carried out by the *ex-situ* prism method.

3.2. Assessment of Biodegradation of Petroleum Pollutants on the Basis of Chromatographic Analyses

The effectiveness of the biodegradation process of petroleum hydrocarbons in soils inoculated with the biopreparation A1 and the biopreparation A1 with γ-PGA was determined on the basis of chromatographic analyses.

Chromatographic Analysis of TPH, BTEX and PAHs

As a result of inoculation of soil AB with the biopreparation A1, after 6 months of the purification process, the TPH content was reduced from 19,774.23 mg/kg DM to 6717.33 mg/kg DM, while as a result of inoculation with the biopreparation A1 with the addition of γ-PGA, the TPH content was reduced to 4111.47 mg/kg DM.

The course of TPH biodegradation in the AB soil inoculated with the biopreparation A1, as well as with the biopreparation A1 with the addition of γ-PGA, was similar in the following months of the process, but its pace decreased, and at the end remained at a similar level. Degrees of biodegradation of TPH content during the purification process by inoculation with the biopreparation A1 were: after 2 months 35.03% (Soil A2), after 4 months 53.18% (Soil A4), and after 6 months 66.03% (Soil A6). The degree of TPH reduction during inoculation with the biopreparation A1 with the addition of PGA was higher and was as follows: after 2 months 48.21% (Soil B2), after 4 months 65.73% (Soil B4), and after 6 months 79.21% (Soil B6).




Figure 2. Dependence of the amount of oxygen consumed [mg·dm^{-3}] on time in AB soil inoculated with biopreparation A1 and biopreparation A1 with γ-PGA addition. Control-sterile uncontaminated "pure" soil inoculated with biopreparation A1 and biopreparation A1 with γ-PGA. Blank sample is soil AB with distilled water.

The results of the chromatographic analyses showed that as a result of inoculation with both the biopreparation A1 and the biopreparation A1 with the addition of γ-PGA, the biodegradation of aliphatic hydrocarbons with a carbon chain length of n-C_9–n-C_{21} was the fastest. The degrees of biodegradation of these hydrocarbons in soil after 2, 4 and 6 months were in the range of: 45.60–75.14 (Soil A2), 67.25–87.85 (Soil A4), 86.52–94.59% (Soil A6) and 56.87–85.49% (Soil B2), 88.46–96.67% (Soil B4) and 96.57–99.55% (Soil B6). Hydrocarbons in the n-C_{22}–n-C_{30} range were also satisfactorily biodegradable within the range: Soil A2 (20.87–39.66%), Soil A4 (29.08–64.26%), Soil A6 (44.62–77.77%), Soil B2 (34.43–49.46%), Soil B4 (56.24–79.15%), Soil B6 (80.35–88.35%). Hydrocarbons with more than 30 carbon atoms in the molecule turned out to be the most difficult to biodegrade. The degrees of their biodegradation after 6 months of the treatment process in soil A6 were within the range of 25.13–40.60%, while in soil B6 it was within the range of 46.43–65.73%. The content of unidentified hydrocarbons in Soil A6 decreased by 59.49%, and in Soil B6 by 71.84%.

Chromatographic analysis allowed for a broader look at the course of the process, as it made it possible to determine the biodegradability of individual groups of hydrocarbons. Both in the soil inoculated with the biopreparation A1 and the biopreparation A1 with the addition of γ-PGA, the rate of removal of individual groups of hydrocarbons was in a decreasing order: n-C_{10}–n-C_{22} > n-C_{23}–n-C_{28} > n-C_{28}–n-C_{40}. This removal model is likely related to the chemical structure of the alkanes. Alkanes with a chain length of n-C_{10}–n-C_{21} are the substances most willingly used by bacteria in metabolic processes [36,100,101,108], while alkanes with longer carbon chains are less biodegradable. Straight and short carbon chain hydrocarbons (n-C_{10}–n-C_{22}) present in soil are therefore perceived as the most attractive source of energy by any organism capable of degrading them. The ability of individual bacterial strains (present in the biopreparation) to decompose hydrocarbons also plays an important role in the process of biodegradation of aliphatic hydrocarbons. Bacterial strains capable of decomposing various types of petroleum hydrocarbons were

used in the biopreparation A1. Nevertheless, each bacterial strain may prefer a different type of hydrocarbon. For instance, some bacteria can metabolize specific alkanes, while others break down aromatic hydrocarbons. Taking into account the ability of various bacterial strains to degrade aliphatic hydrocarbons, it should be noted that Dietzia e a broad spectrum of the ability to degrade oil-derived hydrocarbons in the n-C_6–n-C_{40} range) [37]. *Pseudomonas* most effectively degraded n-alkanes in the n-C_{14}–n-C_{30} range, [109], while *Rhodoccocus* in the n-C_{13}–n-C_{17} range [63]. *Gorgonia* had the ability to effectively degrade both linear and branched aliphatic hydrocarbons [38].

A comparison of the content of n-alkanes in the tested soil (Soil AB) inoculated with (a) biopreparation A1; (b) biopreparation A1 with the addition of γ-PGA after 2 (Soil A2/Soil B2), 4 (Soil A4/Soil B4) and 6 (Soil A6/Soil B6) months of the biodegradation process is presented in Figure 3.




Figure 3. Residual content of n-alkanes in soil contaminated petroleum hydrocarbons inoculated (**a**) biopreparation A1, (**b**) biopreparation A1 with the addition of γ-PGA after 2 (Soil A2/Soil B2), 4 (Soil A4/Soil B4) and 6 (Soil A6/Soil B6) months of the biodegradation process carried out in ex-situ conditions. Control was non-inoculated Soil AB. (repetition number $n = 7–10$, $p < 0.05$).

The indicators of biodegradation degree prove that the course of biological decomposition of n-alkanes is satisfactory, both in soil inoculated with the biopreparation A1 and in the biopreparation A1 with the addition of γ-PGA. In A6 soil these indicators decreased: n-C17/Pr from 3.0 to 0.48, n-C18/Ph from 3.14 to 0.53, while in B6 soil n-C17/Pr from 3.0 to 0.20, n-C18/Ph from 3.14 to 0.21. The presented results prove that with proper control of treatment processes (appropriate temperature, humidity, nutrient content, aeration) and extensive monitoring (chromatographic analysis), a satisfactory biodegradation efficiency of aliphatic hydrocarbons can be obtained.

The chromatographic analyses performed showed that the addition of γ-PGA to the biopreparation increases the efficiency of biodegradation of aliphatic hydrocarbons. The comparative analysis carried out on identical samples of the initial soil AB inoculated with the biopreparation A1 and the biopreparation A1 with the addition of γ-PGA showed that the degree of TPH biodegradation after 6 months of the process in B6 soil is 13.18% higher than in A6 soil.

As a result of the biodegradation process carried out under semi-technical conditions (*ex-situ* method) using the biopreparation as an inoculant, after 6 months of the process, the BTEX content was reduced from 17.45 mg/kg DM to 3.48 mg/kg DM. The results of the chromatographic analyses showed that during the inoculation with the biopreparation A1, the biodegradation of toluene took place the fastest and the degrees of its reduction after 2, 4 and 6 months were, respectively: 63.22%, 84.17% and 96.54%. Ethylbenzene and benzene turned out to be slightly less biodegradable, their biodegradation rates were, respectively, 60.47% and 59.17 in Soil A2, 80.08% and 79.98% in Soil A4, and 93.15% and 91.75% in Soil A6. The least biodegradable were xylenes, whose biodegradation rates after the end of the process were at the level of 39.91–46.85% (Soil A2), 55.73–62.29% (Soil A4), and 71.06–78.69% (Soil A6).

Biodegradation of monoaromatic hydrocarbons in the soil inoculated with the biopreparation A1 with the addition of γ-PGA was more effective, as evidenced by the reduction in BTEX content after 6 months of purification to the level of 1.71 mg/kg DM (90.19%). Compared to the soil inoculated with the biopreparation A1, the application of γ-PGA additive increased the degree of BTEX biodegradation by 10.11%. Toluene biodegradation rates over 6 months were 66.06% in Soil B2, 89.88% in Soil B4, and 98.89% in Soil B6. The decrease in the content of ethylbenzene and benzene after 2 months of the experiment was 65.93% and 63.89%, after 4 months it was 86.66% and 84.17%, and after 6 months it was 97.69% and 95.89%. The degrees of biodegradation of the least biodegradable xylenes were within the range of 45.51–50.33% (Soil B2), 66.49–71.19% (Soil B4) and 85.14–89.88% (Soil B6).

In both inoculation methods, the removal rate of particular monoaromatic hydrocarbons decreased in the order of toluene > ethylbenzene > benzene > p-xylene > m-xylene > o-xylene. The available literature data indicate that the degradation rate of particular hydrocarbons from the BTEX group is related to the type of bacterial strain used in the study [110,111]. Research by other scientists shows that the ability to degrade monoaromatic hydrocarbons for *Pseudomonas putida* and *Bacillus cereus* strains decreases in the following order: toluene > ethylbenzene > benzene > xylenes [111], while the data obtained by Shima indicate that for the *Pseudomonas putida* strain this ability decreases in the order toluene > benzene > xylenes > ethylbenzene. In the differences in the order of degradation of individual monoaromatic hydrocarbons, the authors refer to different environmental conditions from which given bacterial strains were isolated [111]. The degradation of BTEX by the *Rhodococcus* sp. decreases in the order of toluene > xylenes > ethylbenzene [112]. By contrast, *Bacillus amyloliquefaciens* degraded monoaromatic hydrocarbons in the order toluene > benzene, ethylbenzene, p-xylene > m-xylene > o-xylene [110]. In the case of inoculation with biopreparations, the rate of removal of individual BTEX by biodegradation may therefore be different depending on the qualitative and quantitative composition of the biopreparation. The use of a mixed culture to inoculate soil contaminated with petroleum hydrocarbons influenced the effectiveness of the BTEX biodegradation process, because in-

terspecies interactions may be necessary for complete biodegradation of a multicomponent mixture of hydrocarbons [111,113].

Moreover, hydrocarbons from the BTEX group belong to volatile organic pollutants, which additionally affects the value of the determination of these compounds in soil samples collected after 2, 4 and 6 months of biodegradation under semi-technical conditions. The use of the aeration system in the construction of the stand to conduct the biodegradation process using the *ex-situ* prism method influenced the speed and efficiency of removing volatile organic pollutants (BTEX) from the soil [114]. The combination of inoculants capable of effectively removing hydrocarbons from the BTEX group and appropriate conditions (constant temperature 20–25 °C, aeration) for the biodegradation process allowed to achieve very high levels of reduction of these compounds after the end of the process (71.06–98.89%).

A comparison of the monoaromatic hydrocarbons content in the tested soil (Soil AB) inoculated with (a) biopreparation A1; (b) biopreparation A1 with the addition of γ-PGA after 2 (Soil A2/Soil B2), 4 (Soil A4/Soil B4) and 6 (Soil A6/Soil B6) months of the biodegradation process is presented in Figure 4.




Figure 4. Residual content of BTEX in soil contaminated petroleum hydrocarbons inoculated (**a**) biopreparation A1, (**b**) biopreparation A1 with the addition of γ-PGA after 2 (Soil A2/Soil B2), 4 (Soil A4/Soil B4) and 6 (Soil A6/Soil B6) months of the biodegradation process carried out in ex-situ conditions. Control was non-inoculated Soil AB. (repetition number $n = 7–10$, $p < 0.05$).

The performed chromatographic analysis allowed to determine the degree of reduction of particular identified PAHs. In the soil inoculated with the biopreparation A1, the highest degree of reduction of the content after the end of the biodegradation process was recorded for naphthalene and it was 49.72% after 2 months, 62.11% after 4 months, and 71.58% after 6 months. A slightly lower degree of biodegradation was obtained for 3-ring PAHs (ACE, FLU, PHEN, ANTH) 31.37–44.44% (Soil A2), 50.00–56.32% (Soil A4), and 63.73–64.44% (Soil A6). In the case of 4-ring PAHs (FLTH, PYR, B[a]A, CHRY), the reduction in their content was much lower and ranged from 14.06% to 30.14% after 2 months, from 23.47% to 38.93% after 4 months, and from 32.33% to 42.78% after 6 months. 5-ring PAHs and 6-ring PAHs were much more difficult to biodegrade, because the efficiency of the biodegradation process after 2 months of the process was determined at the level of 13.98–15.38%, after 4 months in the range of 22.68–20.78%, and after 6 months within the range of 24.62–30.06%. As a result of the biodegradation process carried out in semi-technical conditions (using the *ex-situ* method) with the use of a biopreparation A1, after 6 months of the process, the PAHs content was reduced from 27.03 mg/kg DM to 16.52 mg/kg DM, which is 38.86%. In turn, as a result of inoculation with a biopreparation A1 with the addition of γ-PGA, it

was reduced to 13.19 mg/kg DM (51.18%). Therefore, the use of the γ-PGA addition to the biopreparation A1 increased the efficiency of the PAHs biodegradation process by 12.32%. Inoculation with the biopreparation A1 with the addition of γ-PGA allowed to reduce the content of naphthalene after 2 months by 59.93%, after 4 months by 77.41%, and finally after 6 months by 81.06%. Reduction levels of 3-ring PAHs (ACE, FLU, PHEN, ANTH) were at a level of 45.10–52.22% in Soil B2, 60.78–63.33% in Soil B4 and 68.63–72.22% in Soil B6, while 4-ring PAHs (FLTH, PYR, B[a]A, CHRY) at a level of 26.93–37.05% in Soil B2, 39.11–46.73% in Soil B4 and 44.65–56, 24% in Soil B6. A slightly lower degree of reduction, within the range from 22.37% to 28.87% in Soil B2, from 31.71% to 33.00% in Soil B4 and from 40.30% to 43.17% in Soil B6, was obtained for 5-ring PAHs (B[b]F, B[k]F, B[a]P). The degree of biodegradation of 6-ring PAHs (D[ah]A, B[ghi]P, IND) was the lowest among all PAHs, as it was 15.11–25.40% after 2 months, 25.72–34.72% after 4 months and 6 months after 35.48–37.83%.

PAHs can be adsorbed, volatilized, photolyzed and chemically degraded, but nevertheless microbiological degradation is the main process of their removal from the environment. The degradation of PAHs depends on the environmental conditions, the number and type of microorganisms, the nature and chemical structure of the degraded chemical compound [115]. In general, the biodegradation rate of PAHs decreases with an increase in the number of aromatic rings in the molecule 2-ring PAHs > 3-ring PAHs > 4-ring PAHs > 5-ring PAHs > 6-ring PAHs [4]. 2-ring naphthalene is relatively volatile, soluble and biodegradable. Aromatic hydrocarbons with more rings in the molecule are less biodegradable and less soluble in water. This is because the stability of polycyclic aromatic hydrocarbons increases with increasing molecular weight. Moreover, as the number of aromatic rings in the molecule increases, the vapor pressure of PAHs and their solubility in water decrease, while the resistance to oxidation and reduction increases [12]. It is known that many species of bacteria are capable of decomposing PAHs, and most of them are isolated from contaminated soil. Strains of *Acinetobacter*, *Arthrobacter*, *Burkholderia*, *Dietzia*, *Mycobacterium*/*Mycolicibacterium*, *Rhodococcus Pseudomonas* and *Sphingomonas* are among the commonly known PAHs degraders [115]. The use of A1 biopreparation developed on the basis of indigenous bacteria capable of decomposing a wide range of petroleum pollutants, including PAHs allowed for an effective reduction of the analyzed polycyclic aromatic hydrocarbons in the studied soil. The use of the γ-PGA additive allowed to increase the efficiency of the PAHs biodegradation process, compared to the A1 biopreparation.

A comparison of the content of polycyclic aromatic hydrocarbons in the tested soil (Soil AB) inoculated with (a) biopreparation A1; (b) biopreparation A1 with the addition of γ-PGA after 2 (Soil A2/Soil B2), 4 (Soil A4/Soil B4) and 6 (Soil A6/Soil B6) months of the biodegradation process is presented in Figure 5.

The chromatographic analyses performed showed that the addition of γ-PGA to a biopreparation has a positive effect on the efficiency of the biodegradation process of petroleum hydrocarbons. As a result of inoculation of soil AB with the biopreparation A1 with the addition of γ-PGA over 6 months, higher degrees of biodegradation of all analyzed analytes were obtained than as a result of inoculation with the biopreparation A1. It can therefore be concluded that γ-PGA supports the decomposition of both aliphatic and aromatic hydrocarbons in the presence of the biopreparation. High values of biodegradation degrees of particular groups of petroleum hydrocarbons (aliphatic hydrocarbons, BTEX, WWAs) confirm the supposition that the biopreparation consisted of *Dietzia* sp. *IN133*, *Gordonia* sp. *IN138 Mycolicibacterium frederiksbergense IN53*, *Rhodococcus erythropolis IN119*, *Rhodococcus* sp. *IN136* and *Pseudomonas* sp. *IN132*. in combination with gamma polyglutamic acid, it can be used as an inoculum in bioaugmentation procedures. The use of this type of inoculation allows for the effective biodegradation of petroleum pollutants present in contaminated soils, and is also safe for the environment (Section 3.3). The summary of the content of particular groups of petroleum hydrocarbons in the initial soil AB and after the completion of the biodegradation process (6 months) is provided in Table 3.




Figure 5. Residual content of PAHs in soil contaminated petroleum hydrocarbons inoculated (**a**) biopreparation A1, (**b**) biopreparation A1 with the addition of γ-PGA after 2 (Soil A2/Soil B2), 4 (Soil A4/Soil B4) and 6 (Soil A6/Soil B6) months of the biodegradation process carried out in ex-situ conditions. Control was non-inoculated Soil AB. (repetition number $n = 7–10$, $p < 0.05$). (naphthalene (NAP), acenaphthene (ACE), fluorene (FLU), phenanthrene (PHEN), anthracene (ANTH), fluoranthene (FLTH), pyrene (PYR), benzo[a]anthracene (B[a]A), chrysene (CHRY), benzo[b]fluoranthene (B[b]F), benzo[k]fluoranthene (B[k]F), benzo[a]pyrene (B[a]P), benzo[g,h,i]perylene (B[ghi]P), indeno[1,2,3-c,d]pyrene (IND), and dibenz[a,h]anthracene (D[ah]A).

Table 3. Summary of the content of petroleum hydrocarbons (TPH, BTEX, PAHs) in soil AB and after the completion of the biodegradation process with the use of biopreparation A1 (Soil A6) and biopreparation A1 with the addition of γ-PGA (Soil B6), (repetition number $n = 7–10$, $p < 0.05$).

Hydrocarbons	Content ± SD (mg/kg d.w. Soil)		
	Initial Soil AB	After 180 Days	
		Soil A6 (Inoculated with Biopreparation)	Soil B6 (Inoculated with Biopreparation with PGA)
TPH	19,774.23 ± 988.71	6717.33 ± 671.73	4111.47 ± 411.15
Unidentified aliphatic hydrocarbons	10,196.10 ± 509.81	4130.44 ± 413.04	2871.44 ± 287.14
∑ n-C_9–n-C_{21}	5356.69 ± 535.67	600.42 ± 120.08	154.50 ± 30.90
∑ n-C_{22}–n-C_{30}	3027.63 ± 302.76	1149.09 ± 229.82	451.96 ± 90.39
∑ n-C31–n-C_{40}	683.31 ± 68.33	447.55 ± 89.51	276.31 ± 55.26
BTEX	17.45 ± 1.75	3.48 ± 0.70	1.71 ± 0.34
benzene	3.56 ± 0.53	0.29 ± 0.09	0.15 ± 0.05
toluene	1.11 ± 0.17	0.04 ± 0.02	0.01 ± 0.004
ethylobenzene	0.47 ± 0.09	0.03 ± 0.01	0.01 ± 0.004
xylenes	12.31 ± 1.23	3.11 ± 0.62	1.54 ± 0.31
PAHs	27.03 ± 2.70	16.52 ± 1.62	13.19 ± 1.32
∑ two-ring PAHs	1.37 ± 0.27	0.39 ± 0.12	0.26 ± 0.08
∑ three-ring PAHs	2.65 ± 0.40	0.96 ± 0.14	0.76 ± 0.11
∑ four-ring PAHs	15.19 ± 1.52	9.41 ± 0.94	7.47 ± 0.75
∑ five-ring PAHs	6.01 ± 0.60	4.41 ± 0.44	3.55 ± 0.36
∑ Six-ring PAHs	1.81 ± 0.36	1.36 ± 0.27	1.15 ± 0.23

The course of the biodegradation process of petroleum pollutants (TPH, BTEX, PAHs) during the AB soil inoculation process with biopreparation A1 and biopreparation A1

with the addition of γ-PGA is described by Equation (2). The individual coefficients of Equation (2) are summarized in Table 4.

Table 4. Coefficients of first-order mathematical model describing TPH, BTEX and PAHs hydrocarbons group biodegradation in soil AB. Measurement repetition number $n = 7–10$, $p < 0.05$.

Petroleum Pollutants Group	k [d^{-1}]		$(C/C_H)_0$		Determination Coefficient (r^2)	
	Inoculated Biopreparation A1	Inoculated Biopreparation A1 with γ-PGA	Inoculated Biopreparation A1	Inoculated Biopreparation A1 with γ-PGA	Inoculated Biopreparation A1	Inoculated Biopreparation A1 with γ-PGA
TPH	0.0058 ± 0.0003	0.0083 ± 0.0005	3763 ± 298	3639 ± 201	0.9944	0.9913
∑ n-C_9–n-C_{22}	0.0113 ± 0.0007	0.0186 ± 0.0011	1177 ± 95	10278 ± 89	0.9912	0.9939
∑ n-C_{23}–n-C_{40}	0.0041 ± 0.0002	0.0069 ± 0.0006	634 ± 23	603 ± 16	0.9931	0.9916
BTEX	0.0072 ± 0.0004	0.0112 ± 0.007	3.403 ± 0.182	3.198 ± 0.16	0.9956	0.9906
PAHs	0.0037 ± 0.002	0.0046 ± 0.002	5.095 ± 0.291	4.844 ± 0.27	0.9897	0.9874
∑ two-ring PAHs	0.0061 ± 0.0004	0.0077 ± 0.0004	0.192 ± 0.061	0.176 ± 0.056	0.9953	0.9894
∑ three-ring PAHs	0.0053 ± 0.0003	0.0062 ± 0.0004	0.529 ± 0.032	0.314 ± 0.024	0.9877	0.9842
∑ four-ring PAHs	0.0031 ± 0.0002	0.0036 ± 0.0002	2.964 ± 0.158	2.366 ± 0.128	0.9847	0.9789
∑ five-ring PAHs	0.0018 ± 0.0002	0.0021 ± 0.0002	1.870 ± 0.091	1.560 ± 0.067	0.9961	0.9909
∑ Six-ring PAHs	0.0009 ± 0.0001	0.0011 ± 0.0002	0.352 ± 0.019	0.322 ± 0.015	0.9842	0.9882

The first order biodegradation constants (k) during inoculation of soil AB with biopreparation A1 for aliphatic hydrocarbons are at the following level: TPH (0.0058 d^{-1}), Σ n-C_9–n-C_{22} (0.0113 d^{-1}), Σ n-C_{23}–n-C_{40} (0.0041 d^{-1}), and for mono and polycyclic aromatic hydrocarbons they are, respectively: BTEX (0.0072 d^{-1}) and PAHs (0.0037 d^{-1}). The presented results prove the biodegradability of both aliphatic and aromatic hydrocarbons. The biodegradation rate of polycyclic aromatic hydrocarbons is directly proportional to the number of rings in the molecule and the size of their concentration in the soil (Table 4) The first order biodegradation constants (k) during the inoculation of soil AB with biopreparation A1 with the addition of γ-PGA visibly increase at the level of: TPH (0.0083 d^{-1}), Σ n-C_9–n-C_{22} (0.0186 d^{-1}), Σ n-C_{23}–n-C_{40} (0.0069 d^{-1}), BTEX (0.0112 d^{-1}) and PAHs (0.0046 d^{-1}) (Table 4). The obtained results prove that the addition of γ-PGA increases the degree of biodegradation of both aliphatic and aromatic hydrocarbons.

3.3. Toxicological Assessment

Pollution of the natural environment with petroleum substances such as TPH, BTEX or PAHs contributes to the constant deterioration of the health of living organisms. Therefore, a comprehensive assessment of soil quality should be based not only on the concentration of selected physico-chemical parameters, but also include a bioindication analysis assessing its toxicity to living organisms. The assessment of the effectiveness of the bioremediation treatments used was supplemented with toxicological monitoring aimed at determining the impact of petroleum pollutants (TPH, BTEX, PAHs) and intermediate metabolites formed during remediation treatments on soil biocenosis based on the performed toxicological tests. Toxicological tests allow for the simultaneous determination of the harmful effect of all substances in the tested sample on selected living organisms, taking into account the interactions between all elements of the tested system. The use of living organisms as bioindicators belonging to various taxonomic groups (bacteria, crustaceans and higher plants) representing all trophic groups (producers, consumers and decomposers) allowed for a comprehensive assessment of the effectiveness of the treatment of soil contaminated with petroleum substances. The results of Phytotoxkit™, Ostracodtoxikit F™ and Microtox SPT toxicology tests is listed in Table 5.

Table 5. The results of Phytotoxkit™, Ostracodtoxikit F™ and Microtox SPT toxicology tests.

Tested Organism	Parameter Measured	Control Soil	Soil AB	Soil A3	Soil B3	Soil A6	Soil B6
Test Phytotoxkit™							
Sorghum saccharatum	germination [%]	100	80	100	100	100	100
	average root length [mm]	48	11.4	27.6	31.7	40.8	44.8
	root growth inhibition mean [%]	0	76.25	42.50	33.96	15.00	6.67
Lepidium Sativum	germination [%]	100	60	80	90	100	100
	average root length [mm]	69	11.6	35.1	44.6	56.4	62.9
	root growth inhibition mean [%]	0	83.19	49.13	35.36	18.26	8.84
Sinapis alba	germination [%]	100	70	90	100	100	100
	average root length [mm]	80	20.2	47.5	55.8	68.4	75.5
	root growth inhibition mean [%]	0	74.75	40.63	30.25	14.5	5.63
Test Ostracodtoxikit F™							
Heterocypris incongruens	average mortality [%]	4	81.67	43.33	36.67	18.33	10.00
	average growth inhibition [%]	-	70.6	34.98	29.01	10.11	4.99
Test Microtox SPT							
Vibrio fischeri	*TU*	0	32.3	16.5	14.2	9.4	4.3

On the basis of the results of toxicological tests, it is clearly visible that the initial soil AB was toxic to the tested species of plants, animals and microorganisms.

In the initial AB soil, the observed seed germination inhibition ranged from 20% to 40% depending on the tested organism, while for each of the analyzed plants, more than 74% of root growth was inhibited. *Lepidium sativum* turned out to be the most sensitive to pollutants from the group (TPH, BTEX, PAHs), while *Sorghum saccharatum* was characterized by the highest resistance. Both bioaugmentation variants used improved the quality of the soil by reducing its phytotoxicity compared to the initial AB soil, but differed in the effectiveness of phytotoxicity reduction. The soil inoculated with the biopreparation A1 with the addition of γ-PGA was characterized by lower phytotoxicity after 6 months of the purification process, for which the inhibition of root growth decreased by 69.58–74.35%, while the inhibition of seed germination decreased by 20–40% compared to the starting soil AB. Slightly lower values of the decrease in root growth inhibition after 6 months of biodegradation were observed in the soil inoculated with the biopreparation A1 and amounted to 61.25–64.93% in relation to the AB soil. In general, it can be concluded that the greater the reduction in the content of petroleum hydrocarbons during the biodegradation process, the less phytotoxic the properties of the soil are.

The Ostracodtoxikit F™ direct contact test carried out on the soil before the inoculation treatment showed the toxic properties of the tested soil, as evidenced by the high mortality of *Heterocypris incongruens* (83.33%) and inhibition of the growth of ostracods by 70.60%. The successive reduction of the content of petroleum pollutants through the conducted inoculation procedures resulted in the reduction of the toxicity of the tested soil for spouses, which improved the survival and growth of *Heterocypris incongruens*. The tests performed after 180 days of biodegradation by inoculation with the biopreparation A1 with the addition of γ-PGA showed a 90.00% survival effect of *Heterocypris incongruens*, while the average length of the tested organisms after the end of the test was 764 μm. The second examined parameter, growth inhibition of ostracods, was 4.99%. Compared to the inoculation with the biopreparation A1, the use of γ-PGA supplemented the survival effect of ostracods by 8.33% and the decrease in the average growth inhibition of *Heterocypris incongruens* by 5.12%.

The soil toxicity tests at the trophic level of the decomposers were carried out using the Microtox® SPT test (for the solid phase), which allowed for direct contact of luminescent bacteria (*Vibrio fischeri*) with the tested soil sample. Thanks to this, it is possible to detect

not only water-soluble substances, but also hydrophobic and lipid compounds. Acute toxicity expressed in toxic units (TU) was high in the initial AB soil (Table 4). Conducting the inoculation treatment with the biopreparation A1 and the biopreparation A1 with the addition of γ-PGA resulted in a reduction of the toxicity of the tested soil for microorganisms. Compared to the initial AB soil, inoculation with the biopreparation decreased soil toxicity by 22.8%, and the inoculation with the biopreparation A1 supplemented with γ-PGA by 28%.

In order to determine the presence of mutagenic and carcinogenic compounds in soil samples: raw soil (Soil AB), and after a 6-month inoculation period with biopreparation A1 (Soil A) and biopreparation A1 with addition of γ-PGA (Soil B) Ames test (Figure S2) was run. The study used the TA-100 strain, which is a standard strain of *Salmonella typhimirium*. Its mutations occur on the histidine operon, which is unable to synthesize amino acids. It was noticed that the use of the microsomal S9 fraction as a metabolic activator of promutagenes had a slight effect on the increase in the number of reversible mutations. The results of the tested soil samples are shown in Figure 6 as changes in the number of revertants indicated depending on the concentration of pollutants. In the case of a sample of raw soil (Soil AB) found to be mutagenic, the mutagenicity coefficient was 6.15. After biodegradation of petroleum pollutants as a result of inoculation with the biopreparation A1 (6 months) (soil A6), the mutagenicity coefficient decreased gradually to 2.15, and after inoculation with the biopreparation with the addition of γ-PGA (Soil B6) to 1.54 (Figure 6). The Soil A6 and Soil B6 samples showed no mutagenic properties as the number of non-histidine induced revertants was only slightly higher than the number of spontaneous mutations present in the control sample.




Figure 6. Influence of TPH, BTEX and PAHs concentration of the revertants number ($n = 3$, $p < 0.05$): (**a**) crude soil (Soil AB); (**b**) Soil A6—soil after a 6-month period of biopreparation A1 inoculation; (**c**) Soil B6—soil after a 6-month period of biopreparation A1 with the addition of γ-PGA inoculation.

The conducted toxicological tests confirmed the correlation between the effectiveness of removing petroleum pollutants from the soil matrix and the reduction of its toxicity. In the soil inoculated with the biopreparation with the addition of γ-PGA, a greater reduction in the content of TPH, BTEX and PAHs was observed, and thus a greater improvement in the quality of the tested soil in terms of its toxicity, than in the soil inoculated with the biopreparation A1.

The susceptibility of petroleum substances to microbial degradation depends mainly on their chemical structure. Aliphatic hydrocarbons are usually more easily degraded than that aromatic, long-chain paraffins easier than short-chain, saturated compounds easier than unsaturated and unbranched faster than branched. Cycloalkanes, carbon polycyclic hydrocarbons, compounds with naphthenic-aromatic structures, asphaltenes and other heavy fractions are usually less biodegradable [4]. Biodegradation with the use of biopreparations developed on the basis of indigenous microorganisms capable of decomposing

a wide spectrum of pollutants is currently considered one of the most efficient and environmentally friendly techniques for land remediation. The ability of microorganisms to catabolize various types of organic pollutants is a necessary condition for the proper conduct of the biodegradation process, which is why it has been extensively studied and discussed by researchers from around the world. The challenge now is to improve the efficiency of land remediation processes by biological methods. An interesting solution seems to be the use of various types of additives enriching biopreparations. It is important that the additives used are not toxic. Such an additive in the case of remediation of soils contaminated with petroleum hydrocarbons can be γ-PGA. The conducted research on the use of the γ-PGA polyamide polymer as an additive to a biopreparation A1 has shown that it increases the efficiency of the biodegradation process of petroleum pollutants in the soil matrix. This is due to the special structure of γ-PGA, which makes this compound less susceptible to the action of proteolytic enzymes, which protects the bacterial cell against microbial attack. Moreover, γ-PGA is characterized by strong hygroscopic properties, which in turn allows to maintain favourable conditions for the treatment process of contaminated soils, even in unfavourable weather conditions (drought). It is estimated that in the form of a hydrogel, γ-polyglutamic acid is able to bind water molecules in an amount up to 3500 times its mass [116]. Moreover, depending on the pH of the environment in which it is located, it can be highly soluble in water (pH $\geq$ 6.5) or in organic solvents (pH = 2). Due to these properties, as well as good biodegradability, lack of toxicity and immunogenicity, research into the use of γ-polyglutamic acid in various fields of science is constantly being conducted [117]. In environmental engineering, γ-PGA has been used as a compound that binds harmful substances, which is used in the remediation of soils contaminated with heavy metals. Literature reports indicate that γ-PGA may be released into the environment to sequester toxic metal ions, reduce the salt concentration in the soil environment, provide a carbon source as food for microorganisms, and protect against adverse conditions. In addition, γ-PGA can improve biofilm formation and aid the absorption of essential nutrients from the environment [118]. Research on the biodegradation of petroleum hydrocarbons carried out by the *ex-situ* prism method has shown that γ-polyglutamic acid can be used as an additive to biopreparations in order to increase the efficiency of treating contaminated soils. In addition, research studies have shown that γ-PGA improves the bioavailability of nitrogen in the soil due to microbial transmission in the soil, which increases the possibility of obtaining data by hydro massage. Research by Zhang and his team [119] The study result that γ-PGA may stem from an increase in root biomass. In addition, γ-PGA can be used by plants as a source of nitrogen or carbon, we support their construction. The chromatographic analyses carried out showed that the addition of γ-PGA to the biopreparation A1 increased the biodegradation efficiency of TPH by 13.18%, BTEX by 10.11%, and PAHs by 12.32%. In addition, the toxicological tests carried out showed that the addition of γ-PGA is not toxic and does not increase soil toxicity during the treatment process, and even has a positive effect on its quality. This statement is confirmed by the results of the Phytotoxkit^{TM}, Ostracodtoxikit F^{TM}, Microtox SPT, and Ames tests carried out during biodegradation, which indicate a greater reduction in the toxicity level of soil inoculated with the biopreparation A1 with the addition of γ-PGA. Moreover, the research carried out with the use of the Phytotoxkit^{TM} test confirms the research of Zhang and his team [119] on the effect of the addition of γ-PGA on the biomass of plant roots. Greater root length of *Sorghum saccharatum*, *Lepidium Sativum* and *Sinapis alba* plants was observed in soil inoculated with biopreparation A1 with the addition of γ-PGA than in soil inoculated with biopreparation A1.

4. Conclusions

The aim of the study was to evaluate the possibility of using γ-PGA as an additive to the biopreparation A1 in the process of biodegradation of petroleum hydrocarbons. During the 6-month research, the effectiveness of the biodegradation process of petroleum hydrocarbons (TPH, BTEX, PAHs) was determined as a result of inoculation of contaminated soil

with biopreparation A1 and biopreparation A1 with the addition of γ-PGA. Biodegradation studies carried out by the *ex-situ* prism method with the use of 2 inoculation methods have shown that the addition of γ-PGA increases the efficiency of decomposition of TPH, BTEX and PAHs. The higher microbiological activity observed during respirometric tests and the conducted chromatographic analyses showed that the addition of γ-PGA to the biopreparation increased the biodegradation efficiency of TPH by 13.18%, BTEX by 10.11%, and PAHs by 12.32% compared to the biopreparation A1. The introduction of a set of selected toxicological tests allowed for the observation of changes in the impact of the concentration of petroleum pollutants on various types of bioindicators (plants, animals, microorganisms) representing all trophic groups during the biodegradation process of TPH, BTEX and PAHs. The conducted toxicological tests confirmed the correlation between the effectiveness of removing petroleum pollutants from the soil matrix and the reduction of its toxic effect on the tested bioindicators. This is evidenced by the values of toxicity indicators in the Phytotoxkit^{TM} (germination inhibition, root growth inhibition), Ostracodtoxkit F^{TM} (mortality, growth inhibition) and Microtox SPT (TU) tests, which after the completion of the biodegradation process of petroleum hydrocarbons were lower in the soil inoculated with A1 biopreparation with the addition of γ-PGA rather than A1 biopreparation only. Research on the acceleration of soil bioremediation processes using microorganisms is now an important aspect of environmental engineering. The use of gamma polyglutamic acid as an additive to a biopreparation based on autochthonous bacteria can significantly improve the efficiency of biodegradation processes. Satisfactory results of research carried out in *ex-situ* conditions create the prospect of adopting the method of cleaning soils contaminated with petroleum hydrocarbons with the use of biopreparation A1 with the addition of γ-PGA on an industrial scale.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/ma15020400/s1. Figure S1: Content (%) of individual components of petroleum pollutants in the soil, Ecotoxicological analyses –description of the methodology. Figure S2: Scheme of Ames Test conduct.

Author Contributions: Conceptualization, K.W., T.S. (Teresa Steliga), P.K., T.S. (Tomasz Skalski); funding acquisition K.W., T.S. (Teresa Steliga) and P.K.; formal analysis, K.W., T.S. (Teresa Steliga), P.K. and J.B.; investigation, K.W., T.S. (Teresa Steliga), P.K. and J.B.; methodology, K.W., T.S. (Tomasz Skalski) and J.B.; writing—original draft preparation, K.W., T.S. (Teresa Steliga) and P.K., writing—review and editing, K.W., T.S. (Teresa Steliga), P.K., J.B.; T.S. (Tomasz Skalski). All authors have read and agreed to the published version of the manuscript.

Funding: This research was financially supported by Polish Ministry of Science and Higher Education within statutory funding for Oil and Gas Institute-National Research Institute.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Steliga, T.; Kluk, D. Assessment of the Suitability of Melilotus Officinalis for Phytoremediation of Soil Contaminated with Petroleum Hydrocarbons (TPH and PAH), Zn, Pb and Cd Based on Toxicological Tests. *Toxics* **2021**, *9*, 148. [CrossRef]
2. de Souza, R.B.; Guilherme, T.; Aparecida, C.; Gimenez, T.; Silvia, C. Soil Contamination with Heavy Metals and Petroleum Derivates: Impact on Edaphic Fauna and Remediation Strategies. In *Soil Processes and Current Trends in Quality Assessment*; Hernandez Soriano, M.C., Ed.; InTech: London, UK, 2013; ISBN 9789535110293.
3. Steliga, T.; Jakubowicz, P.; Kapusta, P. Optimisation Research of Petroleum Hydrocarbon Biodegradation in Weathered Drilling Wastes from Waste Pits. *Waste Manag. Res.* **2010**, *28*, 1065–1075. [CrossRef] [PubMed]
4. Steliga, T.; Kluk, D. *Assessment of the Composition of Pollution of Soil Contaminated with TPH and PAHs for the Development of the Bioremediation Technology*; Instytut Nafty i Gazu-Państwowy Instytut Badawczy: Kraków, Poland, 2017; ISBN 9788365649201.

5. Podsiadło, Ł.; Krzyśko-Łupicak, T. Techniki bioremediacji substancji ropopochodnych i metody oceny ich efektywności, Inżynieria i Ochrona Środowiska. *Inżynieria I Ochr. Sr.* **2013**, *16*, 459–476.
6. Hossain, F.; Akter, M.A.; Sohan, S.R.; Sultana, N.; Reza, A.; Hoque, K.M.F. Bioremediation Potential of Hydrocarbon Degrading Bacteria: Isolation, Characterization, and Assessment. *Saudi J. Biol. Sci.* **2021**, *29*, 211–216. [CrossRef]
7. Dandie, C.E.; Weber, J.; Aleer, S.; Adetutu, E.M.; Ball, A.S.; Juhasz, A.L. Assessment of Five Bioaccessibility Assays for Predicting the Efficacy of Petroleum Hydrocarbon Biodegradation in Aged Contaminated Soils. *Chemosphere* **2010**, *81*, 1061–1068. [CrossRef] [PubMed]
8. Xu, X.; Liu, W.; Tian, S.; Wang, W.; Qi, Q.; Jiang, P.; Gao, X.; Li, F.; Li, H.; Yu, H. Petroleum Hydrocarbon-Degrading Bacteria for the Remediation of Oil Pollution Under Aerobic Conditions: A Perspective Analysis. *Front. Microbiol.* **2018**, *9*, 2885. [CrossRef]
9. Souza, E.C.; Vessoni-Penna, T.C.; de Souza Oliveira, R.P. Biosurfactant-Enhanced Hydrocarbon Bioremediation: An Overview. *Int. Biodeterior. Biodegrad.* **2014**, *89*, 88–94. [CrossRef]
10. Cameotra, S.S.; Singh, P. Bioremediation of Oil Sludge Using Crude Biosurfactants. *Int. Biodeterior. Biodegrad.* **2008**, *62*, 274–280. [CrossRef]
11. Leahy, J.G.; Colwell, R.R. Microbial Degradation of Hydrocarbons in the Environment. *Microbiol. Rev.* **1990**, *54*, 305–315. [CrossRef]
12. Abdel-Shafy, H.I.; Mansour, M.S.M. A Review on Polycyclic Aromatic Hydrocarbons: Source, Environmental Impact, Effect on Human Health and Remediation. *Egypt J. Pet.* **2016**, *25*, 107–123. [CrossRef]
13. Kluk, D.; Steliga, T. Efektywna Metoda Identyfikacji Zanieczyszczeń Ropopochodnych (TPH) i Wielopierścieniowych Węglowodorów Aromatycznych (WWA) w Glebach. *Nafta-Gaz* **2017**, *73*, 488–495. [CrossRef]
14. Brakstad, O.G.; Bonaunet, K. Biodegradation of Petroleum Hydrocarbons in Seawater at Low Temperatures (0–5 °C) and Bacterial Communities Associated with Degradation. *Biodegradation* **2006**, *17*, 71–82. [CrossRef]
15. Feitkenhauer, H.; Müller, R.; MAuml, H. Degradation of polycyclic aromatic hydrocarbons and long chain alkanes at 6070 °C by *Thermus* and *Bacillus* spp. *Biodegradation* **2003**, *14*, 367–372. [CrossRef]
16. Ji, Y.; Mao, G.; Wang, Y.; Bartlam, M. Structural Insights into Diversity and N-Alkane Biodegradation Mechanisms of Alkane Hydroxylases. *Front. Microbiol.* **2013**, *4*, 58. [CrossRef]
17. Ghosal, D.; Ghosh, S.; Dutta, T.K.; Ahn, Y. Current State of Knowledge in Microbial Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *Front. Microbiol.* **2016**, *7*, 1369. [CrossRef]
18. Xu, R.; Lau, N.L.A.; Ng, K.L.; Obbard, J.P. Application of a Slow-Release Fertilizer for Oil Bioremediation in Beach Sediment. *J. Environ. Qual.* **2004**, *33*, 1210. [CrossRef] [PubMed]
19. Rosa, A.P.; Triguis, J.A. Bioremediation Process on Brazil Shoreline: Laboratory Experiments. *Env. Sci. Poll. Res. Int.* **2007**, *14*, 470–476. [CrossRef] [PubMed]
20. Steliga, T.; Kapusta, P.; Żak, H. Biodegradacja Substancji Ropopochodnych w Odpadach Kopalnianych z Zastosowaniem Bakterii Autochtonicznych. *Inżynieria Ekol.* **2003**, *8*, 34–42.
21. Ortega-Calvo, J.J.; Lahlou, M.; Saiz-Jimenez, C. Effect of Organic Matter and Clays on the Biodegradation of Phenanthrene in Soils. *Int. Biodeterior. Biodegrad.* **1997**, *40*, 101–106. [CrossRef]
22. Wilkes, H.; Buckel, W.; Golding, B.T.; Rabus, R. Metabolism of Hydrocarbons in ***n***-Alkane-Utilizing Anaerobic Bacteria. *J. Mol. Microbiol. Biotechnol.* **2016**, *26*, 138–151. [CrossRef]
23. Safdari, M.-S.; Kariminia, H.-R.; Rahmati, M.; Fazlollahi, F.; Polasko, A.; Mahendra, S.; Wilding, W.V.; Fletcher, T.H. Development of Bioreactors for Comparative Study of Natural Attenuation, Biostimulation, and Bioaugmentation of Petroleum-Hydrocarbon Contaminated Soil. *J. Hazard. Mater.* **2018**, *342*, 270–278. [CrossRef]
24. Wu, M.; Dick, W.A.; Li, W.; Wang, X.; Yang, Q.; Wang, T.; Xu, L.; Zhang, M.; Chen, L. Bioaugmentation and Biostimulation of Hydrocarbon Degradation and the Microbial Community in a Petroleum-Contaminated Soil. *Int. Biodeterior. Biodegrad.* **2016**, *107*, 158–164. [CrossRef]
25. Zhang, Q.; Zhu, L.; Su, J.; Wang, J.; Xie, H.; Wang, J.; Wang, F. Impacts of Nitrogen and Phosphorus on Atrazine-Contaminated Soil Remediation and Detoxification by Arthrobacter sp. Strain HB-5. *Environ. Earth Sci.* **2014**, *71*, 1465–1471. [CrossRef]
26. Guarino, C.; Spada, V.; Sciarrillo, R. Assessment of Three Approaches of Bioremediation (Natural Attenuation, Landfarming and Bioagumentation—Assistited Landfarming) for a Petroleum Hydrocarbons Contaminated Soil. *Chemosphere* **2017**, *170*, 10–16. [CrossRef]
27. Li, F.; Guo, S.; Wu, B.; Wang, S. Pilot-Scale Electro-Bioremediation of Heavily PAH-Contaminated Soil from an Abandoned Coking Plant Site. *Chemosphere* **2020**, *244*, 125467. [CrossRef]
28. Das, N.; Chandran, P. Microbial Degradation of Petroleum Hydrocarbon Contaminants: An Overview. *Biotechnol. Res. Int.* **2011**, *2011*, 1–13. [CrossRef]
29. Varjani, S.J. Microbial Degradation of Petroleum Hydrocarbons. *Bioresour. Technol.* **2017**, *223*, 277–286. [CrossRef]
30. Mrozik, A.; Piotrowska-Seget, Z. Bioaugmentation as a Strategy for Cleaning up of Soils Contaminated with Aromatic Compounds. *Microbiol. Res.* **2010**, *165*, 363–375. [CrossRef]
31. Ma, Y.-L.; Lu, W.; Wan, L.-L.; Luo, N. Elucidation of Fluoranthene Degradative Characteristics in a Newly Isolated Achromobacter Xylosoxidans DN002. *Appl. Biochem. Biotechnol.* **2015**, *175*, 1294–1305. [CrossRef]
32. Mishra, S.; Jyot, J.; Kuhad, R.C.; Lal, B. Evaluation of Inoculum Addition To Stimulate In Situ Bioremediation of Oily-Sludge-Contaminated Soil. *Appl. Environ. Microbiol.* **2001**, *67*, 1675–1681. [CrossRef]

33. Cerqueira, V.S.; Hollenbach, E.B.; Maboni, F.; Vainstein, M.; Camargo, F.; Peralba, M.D.C.R.; Bento, F.M. Biodegradation Potential of Oily Sludge by Pure and Mixed Bacterial Cultures. *Bioresour. Technol.* **2011**, *102*, 11003–11010. [CrossRef]
34. Janbandhu, A.; Fulekar, M.H. Biodegradation of Phenanthrene Using Adapted Microbial Consortium Isolated from Petrochemical Contaminated Environment. *J. Hazard. Mater.* **2011**, *187*, 333–340. [CrossRef]
35. Eskandari, S.; Hoodaji, M.; Tahmourespour, A.; Abdollahi, A.; Baghi, T.; Eslamian, S.; Ostad-Ali-Askari, K. Bioremediation of Polycyclic Aromatic Hydrocarbons by Bacillus Licheniformis ATHE9 and Bacillus Mojavensis ATHE13 as Newly Strains Isolated from Oil-Contaminated Soil. *JGEESI* **2017**, *11*, 1–11. [CrossRef]
36. Wu, M.; Li, W.; Dick, W.A.; Ye, X.; Chen, K.; Kost, D.; Chen, L. Bioremediation of Hydrocarbon Degradation in a Petroleum-Contaminated Soil and Microbial Population and Activity Determination. *Chemosphere* **2017**, *169*, 124–130. [CrossRef]
37. Wang, X.-B.; Chi, C.-Q.; Nie, Y.; Tang, Y.-Q.; Tan, Y.; Wu, G.; Wu, X.-L. Degradation of Petroleum Hydrocarbons (C6–C40) and Crude Oil by a Novel Dietzia Strain. *Bioresour. Technol.* **2011**, *102*, 7755–7761. [CrossRef]
38. Brown, L.M.; Gunasekera, T.S.; Striebich, R.C.; Ruiz, O.N. Draft Genome Sequence of Gordonia Sihwensis Strain 9, a Branched Alkane-Degrading Bacterium. *Genome Announc.* **2016**, *4*, e00622-16. [CrossRef]
39. Juhasz, A.L.; Naidu, R. Bioremediation of High Molecular Weight Polycyclic Aromatic Hydrocarbons: A Review of the Microbial Degradation of Benzo[a]Pyrene. *Int. Biodeterior. Biodegrad.* **2000**, *45*, 57–88. [CrossRef]
40. Kanaly, R.A.; Harayama, S. Biodegradation of High-Molecular-Weight Polycyclic Aromatic Hydrocarbons by Bacteria. *J. Bacteriol.* **2000**, *182*, 2059–2067. [CrossRef]
41. Kawo, A.H.; Bacha, H.Y. Crude Oil Degradation by *Bacillus* and *Micrococcus* Species Isolated from Soil Compost in Kano, Nigeria. *Bayero J. Pure App. Sci.* **2016**, *9*, 108. [CrossRef]
42. Omotayo, A.E.; Ojo, O.Y.; Amund, O.O. Crude Oil Degradation by Microorganisms in Soil Composts. *Res. J. Microbiol.* **2012**, *7*, 209–218. [CrossRef]
43. Churchill, S.A.; Harper, J.P.; Churchill, P.F. Isolation and Characterization of a *Mycobacterium* Species Capable of Degrading Three- and Four-Ring Aromatic and Aliphatic Hydrocarbons. *Appl. Environ. Microbiol.* **1999**, *65*, 549–552. [CrossRef] [PubMed]
44. Steliga, T.; Wojtowicz, K.; Kapusta, P.; Brzeszcz, J. Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture. *Molecules* **2020**, *25*, 709. [CrossRef] [PubMed]
45. Zhang, L.; Zhang, C.; Cheng, Z.; Yao, Y.; Chen, J. Biodegradation of Benzene, Toluene, Ethylbenzene, and o-Xylene by the Bacterium Mycobacterium Cosmeticum Byf-4. *Chemosphere* **2013**, *90*, 1340–1347. [CrossRef]
46. Zhang, Y.; Wang, F.; Zhu, X.; Zeng, J.; Zhao, Q.; Jiang, X. Extracellular Polymeric Substances Govern the Development of Biofilm and Mass Transfer of Polycyclic Aromatic Hydrocarbons for Improved Biodegradation. *Bioresour. Technol.* **2015**, *193*, 274–280. [CrossRef]
47. Rehmann, K.; Hertkorn, N.; Kettrup, A.A. Fluoranthene Metabolism in Mycobacterium sp. Strain KR20: Identity of Pathway Intermediates during Degradation and Growth. *Microbiology* **2001**, *147*, 2783–2794. [CrossRef]
48. Moody, J.D.; Freeman, J.P.; Doerge, D.R.; Cerniglia, C.E. Degradation of Phenanthrene and Anthracene by Cell Suspensions of *Mycobacterium* sp. Strain PYR-1. *Appl. Environ. Microbiol.* **2001**, *67*, 1476–1483. [CrossRef]
49. Kim, S.-J.; Kweon, O.; Freeman, J.P.; Jones, R.C.; Adjei, M.D.; Jhoo, J.-W.; Edmondson, R.D.; Cerniglia, C.E. Molecular Cloning and Expression of Genes Encoding a Novel Dioxygenase Involved in Low- and High-Molecular-Weight Polycyclic Aromatic Hydrocarbon Degradation in *Mycobacterium Vanbaalenii* PYR-1. *Appl. Environ. Microbiol.* **2006**, *72*, 1045–1054. [CrossRef]
50. Kim, S.-J.; Kweon, O.; Jones, R.C.; Freeman, J.P.; Edmondson, R.D.; Cerniglia, C.E. Complete and Integrated Pyrene Degradation Pathway in *Mycobacterium Vanbaalenii* PYR-1 Based on Systems Biology. *J. Bacteriol.* **2007**, *189*, 464–472. [CrossRef] [PubMed]
51. Kweon, O.; Kim, S.-J.; Jones, R.C.; Freeman, J.P.; Adjei, M.D.; Edmondson, R.D.; Cerniglia, C.E. A Polyomic Approach To Elucidate the Fluoranthene-Degradative Pathway in *Mycobacterium Vanbaalenii* PYR-1. *J. Bacteriol.* **2007**, *189*, 4635–4647. [CrossRef]
52. Yang, R.; Liu, G.; Chen, T.; Li, S.; An, L.; Zhang, G.; Li, G.; Chang, S.; Zhang, W.; Chen, X.; et al. Characterization of the Genome of a Nocardia Strain Isolated from Soils in the Qinghai-Tibetan Plateau That Specifically Degrades Crude Oil and of This Biodegradation. *Genomics* **2019**, *111*, 356–366. [CrossRef] [PubMed]
53. Baek, K.-H.; Yoon, B.-D.; Oh, H.-M.; Kim, H.-S.; Lee, I.-S. Biodegradation of Aliphatic and Aromatic Hydrocarbons by *Nocardia* sp. H17-1. *Geomicrobiol. J.* **2006**, *23*, 253–259. [CrossRef]
54. Yang, Y.; Chen, R.F.; Shiaris, M.P. Metabolism of Naphthalene, Fluorene, and Phenanthrene: Preliminary Characterization of a Cloned Gene Cluster from Pseudomonas Putida NCIB 9816. *J. Bacteriol.* **1994**, *176*, 2158–2164. [CrossRef]
55. Mukherjee, A.K.; Bhagowati, P.; Biswa, B.B.; Chanda, A.; Kalita, B. A Comparative Intracellular Proteomic Profiling of Pseudomonas Aeruginosa Strain ASP-53 Grown on Pyrene or Glucose as Sole Source of Carbon and Identification of Some Key Enzymes of Pyrene Biodegradation Pathway. *J. Proteom.* **2017**, *167*, 25–35. [CrossRef] [PubMed]
56. Kotturi, G.; Robinson, C.W.; Inniss, W.E. Phenol Degradation by a Psychrotrophic Strain of Pseudomonas Putida. *Appl. Microbiol. Biotechnol.* **1991**, *34*, 34. [CrossRef]
57. Meckenstock, R.U.; Boll, M.; Mouttaki, H.; Koelschbach, J.S.; Cunha Tarouco, P.; Weyrauch, P.; Dong, X.; Himmelberg, A.M. Anaerobic Degradation of Benzene and Polycyclic Aromatic Hydrocarbons. *J. Mol. Microbiol. Biotechnol.* **2016**, *26*, 92–118. [CrossRef]
58. Reardon, K.F.; Mosteller, D.C.; Bull Rogers, J.D. Biodegradation Kinetics of Benzene, Toluene, and Phenol as Single and Mixed Substrates ForPseudomonas Putida F1. *Biotechnol. Bioeng.* **2000**, *69*, 385–400. [CrossRef]

59. Huang, X.; Shi, J.; Cui, C.; Yin, H.; Zhang, R.; Ma, X.; Zhang, X. Biodegradation of Phenanthrene by *Rhizobium Petrolearium* SL-1. *J. Appl. Microbiol.* **2016**, *121*, 1616–1626. [CrossRef]
60. Muthukumar, G.; Arunakumari, A.; Mahadevan, A. Degradation of Aromatic Compounds ByRhizobium spp. *Plant Soil* **1982**, *69*, 163–169. [CrossRef]
61. Whyte, L.G.; Slagman, S.J.; Pietrantonio, F.; Bourbonnière, L.; Koval, S.F.; Lawrence, J.R.; Inniss, W.E.; Greer, C.W. Physiological Adaptations Involved in Alkane Assimilation at a Low Temperature by *Rhodococcus* sp. Strain Q15. *Appl. Environ. Microbiol.* **1999**, *65*, 2961–2968. [CrossRef]
62. Hamamura, N.; Olson, S.H.; Ward, D.M.; Inskeep, W.P. Microbial Population Dynamics Associated with Crude-Oil Biodegradation in Diverse Soils. *Appl. Environ. Microbiol.* **2006**, *72*, 6316–6324. [CrossRef] [PubMed]
63. Zhukov, D.V.; Murygina, V.P.; Kalyuzhnyi, S.V. Kinetics of the Degradation of Aliphatic Hydrocarbons by the Bacteria Rhodococcus Ruber and Rhodococcus Erythropolis. *Appl. Biochem. Microbiol.* **2007**, *43*, 587–592. [CrossRef]
64. Choi, K.Y.; Zylstra, G.J.; Kim, E. Benzoate Catabolite Repression of the Phthalate Degradation Pathway in *Rhodococcus* Sp. Strain DK17. *Appl. Environ. Microbiol.* **2007**, *73*, 1370–1374. [CrossRef]
65. Deeb, R.A.; Alvarez-Cohen, L. Temperature Effects and Substrate Interactions during the Aerobic Biotransformation of BTEX Mixtures by Toluene-Enriched Consortia AndRhodococcus Rhodochrous. *Biotechnol. Bioeng.* **1999**, *62*, 526–536. [CrossRef]
66. Kim, D.; Chae, J.-C.; Zylstra, G.J.; Kim, Y.-S.; Kim, S.-K.; Nam, M.H.; Kim, Y.M.; Kim, E. Identification of a Novel Dioxygenase Involved in Metabolism of *o* -Xylene, Toluene, and Ethylbenzene by *Rhodococcus* sp. Strain DK17. *Appl. Environ. Microbiol.* **2004**, *70*, 7086–7092. [CrossRef]
67. Zylstra, G.J.; Kim, E. Aromatic Hydrocarbon Degradation by Sphingomonas Yanoikuyae B1. *J. Ind. Microbiol. Biotechnol.* **1997**, *19*, 408–414. [CrossRef]
68. Shuttleworth, K.L.; Sung, J.; Kim, E.; Cerniglia, C.E. Physiological and Genetic Comparison of Two Aromatic Hydrocarbon-Degrading Sphingomonas Strains. *Mol. Cells* **2000**, *10*, 199–205. [CrossRef]
69. Uyttebroek, M.; Spoden, A.; Ortega-Calvo, J.-J.; Wouters, K.; Wattiau, P.; Bastiaens, L.; Springael, D. Differential Responses of Eubacterial, *Mycobacterium*, and *Sphingomonas* Communities in Polycyclic Aromatic Hydrocarbon (PAH)-Contaminated Soil to Artificially Induced Changes in PAH Profile. *J. Environ. Qual.* **2007**, *36*, 1403–1411. [CrossRef]
70. Van Hamme, J.D.; Singh, A.; Ward, O.P. Recent Advances in Petroleum Microbiology. *Microbiol. Mol. Biol. Rev.* **2003**, *67*, 503–549. [CrossRef]
71. Moscoso, F.; Teijiz, I.; Deive, F.J.; Sanromán, M.A. Efficient PAHs Biodegradation by a Bacterial Consortium at Flask and Bioreactor Scale. *Bioresour. Technol.* **2012**, *119*, 270–276. [CrossRef]
72. Tiwari, B.; Manickam, N.; Kumari, S.; Tiwari, A. Biodegradation and Dissolution of Polyaromatic Hydrocarbons by Stenotrophomonas sp. *Bioresour. Technol.* **2016**, *216*, 1102–1105. [CrossRef]
73. Chen, S.; Yin, H.; Ye, J.; Peng, H.; Liu, Z.; Dang, Z.; Chang, J. Influence of Co-Existed Benzo[a]Pyrene and Copper on the Cellular Characteristics of Stenotrophomonas Maltophilia during Biodegradation and Transformation. *Bioresour. Technol.* **2014**, *158*, 181–187. [CrossRef] [PubMed]
74. Bianco, F.; Race, M.; Papirio, S.; Esposito, G. Phenanthrene Biodegradation in a Fed–Batch Reactor Treating a Spent Sediment Washing Solution: Techno–Economic Implications for the Recovery of Ethanol as Extracting Agent. *Chemosphere* **2022**, *286*, 131361. [CrossRef] [PubMed]
75. Mohd-Kamil, N.-A.-F.; Hussain, N.-H.; Mizad, M.B.; Abdul-Talib, S. Enhancing Performance of Sphingobacterium Spiritivorum in Bioremediation Phenanthrene Contaminated Sand: Enhancing Performance of Sphingobacterium Spiritivorum. *Remediation* **2014**, *24*, 119–128. [CrossRef]
76. Xia, S.; Zhang, L.; Davletshin, A.; Li, Z.; You, J.; Tan, S. Application of Polysaccharide Biopolymer in Petroleum Recovery. *Polymers* **2020**, *12*, 1860. [CrossRef]
77. Bajaj, I.; Singhal, R. Poly (Glutamic Acid)—An Emerging Biopolymer of Commercial Interest. *Bioresour. Technol.* **2011**, *102*, 5551–5561. [CrossRef]
78. Chatterjee, P.; Tiwari, D.; Raval, R.; Dubey, A. Coherent Aspects of Multifaceted Eco-Friendly Biopolymer—Polyglutamic Acid from the Microbes. *J. Pure Appl. Microbiol.* **2019**, *13*, 741–756. [CrossRef]
79. Sung, M.-H.; Park, C.; Kim, C.-J.; Poo, H.; Soda, K.; Ashiuchi, M. Natural and Edible Biopolymer Poly-γ-Glutamic Acid: Synthesis, Production, and Applications. *Chem. Rec.* **2005**, *5*, 352–366. [CrossRef]
80. Wang, L.; Chen, S.; Yu, B. Poly-γ-Glutamic Acid: Recent Achievements, Diverse Applications and Future Perspectives. *Trends Food Sci. Technol.* **2022**, *119*, 1–12. [CrossRef]
81. Park, C.; Choi, J.-C.; Choi, Y.-H.; Nakamura, H.; Shimanouchi, K.; Horiuchi, T.; Misono, H.; Sewaki, T.; Soda, K.; Ashiuchi, M.; et al. Synthesis of Super-High-Molecular-Weight Poly-γ-Glutamic Acid by Bacillus Subtilis Subsp. Chungkookjang. *J. Mol. Catal. B Enzym.* **2005**, *35*, 128–133. [CrossRef]
82. Nabi, I.; Das, S. Poly-Glutamic Acid (PGA)—Structure, Synthesis, Genomic Organization and Its Application: A Review. *IJPSR* **2015**, *6*, 2258–2280. [CrossRef]
83. Gooding, E.A.; Sharma, S.; Petty, S.A.; Fouts, E.A.; Palmer, C.J.; Nolan, B.E.; Volk, M. PH-Dependent Helix Folding Dynamics of Poly-Glutamic Acid. *Chem. Phys.* **2013**, *422*, 115–123. [CrossRef]
84. Tobias, F.; Keiderling, T.A. Role of Side Chains in β-Sheet Self-Assembly into Peptide Fibrils. IR and VCD Spectroscopic Studies of Glutamic Acid-Containing Peptides. *Langmuir* **2016**, *32*, 4653–4661. [CrossRef]

85. Hara, T.; Ueda, S. Regulation of Poly Glutamate Production in *Bacillus Subtilis* (*Natto*): Transformation of High PGA Productivity. *Agric. Biol. Chem.* **1982**, *46*, 2275–2281. [CrossRef]
86. Yanagibashi, T.; Kobayashi, M.; Omori, K. Application of Poly-γ-Glutamic Acid Flocculant to Flocculation–Sedimentation Treatment of Ultrafine Cement Suspension. *Water* **2019**, *11*, 1748. [CrossRef]
87. Li, M.; Zhu, X.; Yang, H.; Xie, X.; Zhu, Y.; Xu, G.; Hu, X.; Jin, Z.; Hu, Y.; Hai, Z.; et al. Treatment of Potato Starch Wastewater by Dual Natural Flocculants of Chitosan and Poly-Glutamic Acid. *J. Clean. Prod.* **2020**, *264*, 121641. [CrossRef]
88. Liang, J.; Shi, W. Poly-γ-Glutamic Acid Improves Water-Stable Aggregates, Nitrogen and Phosphorus Uptake Efficiency, Water-Fertilizer Productivity, and Economic Benefit in Barren Desertified Soils of Northwest China. *Agric. Water Manag.* **2021**, *245*, 106551. [CrossRef]
89. Jeong, D.-Y.; Ryu, M.S.; Yang, H.-J.; Park, S. γ-PGA-Rich Chungkookjang, Short-Term Fermented Soybeans: Prevents Memory Impairment by Modulating Brain Insulin Sensitivity, Neuro-Inflammation, and the Gut–Microbiome–Brain Axis. *Foods* **2021**, *10*, 221. [CrossRef]
90. Araki, R.; Fujie, K.; Yuine, N.; Watabe, Y.; Maruo, K.; Suzuki, H.; Hashimoto, K. The Possibility of Suppression of Increased Postprandial Blood Glucose Levels by Gamma-Polyglutamic Acid-Rich Natto in the Early Phase after Eating: A Randomized Crossover Pilot Study. *Nutrients* **2020**, *12*, 915. [CrossRef]
91. Park, S.-B.; Sung, M.-H.; Uyama, H.; Han, D.K. Poly(Glutamic Acid): Production, Composites, and Medical Applications of the next-Generation Biopolymer. *Prog. Polym. Sci.* **2021**, *113*, 101341. [CrossRef]
92. Pang, X.; Lei, P.; Feng, X.; Xu, Z.; Xu, H.; Liu, K. Poly-γ-Glutamic Acid, a Bio-Chelator, Alleviates the Toxicity of Cd and Pb in the Soil and Promotes the Establishment of Healthy Cucumis Sativus L. Seedling. *Environ. Sci. Pollut. Res.* **2018**, *25*, 19975–19988. [CrossRef]
93. Peng, Y.-P.; Chang, Y.-C.; Chen, K.-F.; Wang, C.-H. A Field Pilot-Scale Study on Heavy Metal-Contaminated Soil Washing by Using an Environmentally Friendly Agent—Poly-γ-Glutamic Acid (γ-PGA). *Environ. Sci. Pollut. Res.* **2020**, *27*, 34760–34769. [CrossRef]
94. Yang, Z.-H.; Dong, C.-D.; Chen, C.-W.; Sheu, Y.-T.; Kao, C.-M. Using Poly-Glutamic Acid as Soil-Washing Agent to Remediate Heavy Metal-Contaminated Soils. *Environ. Sci. Pollut. Res.* **2018**, *25*, 5231–5242. [CrossRef]
95. Zhao, G.; Sheng, Y.; Wang, C.; Yang, J.; Wang, Q.; Chen, L. In Situ Microbial Remediation of Crude Oil-Soaked Marine Sediments Using Zeolite Carrier with a Polymer Coating. *Mar. Pollut. Bull.* **2018**, *129*, 172–178. [CrossRef]
96. Luo, S.G.; Chen, S.C.; Cao, W.Z.; Lin, W.H.; Sheu, Y.T.; Kao, C.M. Application of γ-PGA as the Primary Carbon Source to Bioremediate a TCE-Polluted Aquifer: A Pilot-Scale Study. *Chemosphere* **2019**, *237*, 124449. [CrossRef]
97. Brzeszcz, J.; Kapusta, P.; Steliga, T.; Turkiewicz, A. Hydrocarbon Removal by Two Differently Developed Microbial Inoculants and Comparing Their Actions with Biostimulation Treatment. *Molecules* **2020**, *25*, 661. [CrossRef]
98. Brzeszcz, J.; Steliga, T.; Kapusta, P.; Turkiewicz, A.; Kaszycki, P. R-Strategist versus K-Strategist for the Application in Bioremediation of Hydrocarbon-Contaminated Soils. *Int. Biodeterior. Biodegrad.* **2016**, *106*, 41–52. [CrossRef]
99. Wrenn, B.A.; Venosa, A.D. Selective Enumeration of Aromatic and Aliphatic Hydrocarbon Degrading Bacteria by a Most-Probable-Number Procedure. *Can. J. Microbiol.* **1996**, *42*, 252–258. [CrossRef] [PubMed]
100. Steliga, T.; Jakubowicz, P.; Kapusta, P. Changes in Toxicity during in Situ Bioremediation of Weathered Drill Wastes Contaminated with Petroleum Hydrocarbons. *Bioresour. Technol.* **2012**, *125*, 1–10. [CrossRef] [PubMed]
101. Nwankwegu, A.S.; Orji, M.U.; Onwosi, C.O. Studies on Organic and In-Organic Biostimulants in Bioremediation of Diesel-Contaminated Arable Soil. *Chemosphere* **2016**, *162*, 148–156. [CrossRef] [PubMed]
102. Malińska, K. Application of a Modified OxiTop® Respirometer for Laboratory Composting Studies. *Arch. Environ. Prot.* **2016**, *42*, 56–62. [CrossRef]
103. Steliga, T.; Wojtowicz, K. Wykorzystanie Testów Respirometrycznych Do Oceny Efektywności Biodegradacji Osadów z Instalacji Kopalnianych. *Nafta-Gaz* **2019**, *75*, 29–37. [CrossRef]
104. Steliga, T.; Uliasz, M. Spent Drilling Muds Management and Natural Environment Protection. *Gospod. Surowcami Miner.* **2014**, *30*, 135–155. [CrossRef]
105. Chaîneau, C.H.; Yepremian, C.; Vidalie, J.F.; Ducreux, J.; Ballerini, D. Bioremediation of a Crude Oil-Polluted Soil: Biodegradation, Leaching and Toxicity Assessments. *Water Air Soil Pollut.* **2003**, *144*, 419–440. [CrossRef]
106. Wojtowicz, K. Opracowanie Metodyki Oznaczania BTEX w Próbkach Gleb z Wykorzystaniem Chromatografii Gazowej z Przystawką Headspace. *Nafta-Gaz* **2018**, *74*, 201–207. [CrossRef]
107. Khan, A.I. Analysis of 18 Polycyclic Aromatic Hydrocarbons in Soil Using the QuEChERS Method. Available online: https://tools.thermofisher.com/content/sfs/brochures/AN20677-analysis-of-18-polycyclic-aromatic-hydrocarbons-in-soil-using-the-quechers-method.pdf (accessed on 15 March 2021).
108. Abbasian, F.; Lockington, R.; Megharaj, M.; Naidu, R. A Review on the Genetics of Aliphatic and Aromatic Hydrocarbon Degradation. *Appl. Biochem. Biotechnol.* **2016**, *178*, 224–250. [CrossRef]
109. Sugiura, K.; Ishihara, M.; Shimauchi, T.; Harayama, S. Physicochemical Properties and Biodegradability of Crude Oil. *Environ. Sci. Technol.* **1997**, *31*, 45–51. [CrossRef]
110. Wongbunmak, A.; Khiawjan, S.; Suphantharika, M.; Pongtharangkul, T. BTEX Biodegradation by Bacillus Amyloliquefaciens Subsp. Plantarum W1 and Its Proposed BTEX Biodegradation Pathways. *Sci. Rep.* **2020**, *10*, 17408. [CrossRef]

111. Handayani, S.; Safitri, R.; Surono, W.; Astika, H.; Damayanti, R.; Agung, M. Rukiah Biodegradation of BTEX by Indigenous Microorganisms Isolated from UCG Project Area, South Sumatra. In *IOP Conference Series: Earth and Environmental Science*; IOP Publishing: Bristol, UK, 2019; Volume 308, p. 012017. [CrossRef]
112. Lee, E.-H.; Cho, K.-S. Effect of Substrate Interaction on the Degradation of Methyl Tert-Butyl Ether, Benzene, Toluene, Ethylbenzene, and Xylene by Rhodococcus sp. *J. Hazard. Mater.* **2009**, *167*, 669–674. [CrossRef]
113. Sarkar, P.; Rai, A.R.; Ghosh, S. Degradation of Aromatic Petroleum Hydrocarbons (Btex) by a Solvent Tolerant Bacterial Consortium. *J. Urban Environ. Eng.* **2013**, *7*, 274–279. [CrossRef]
114. Yang, X.; Beckmann, D.; Fiorenza, S.; Niedermeier, C. Field Study of Pulsed Air Sparging for Remediation of Petroleum Hydrocarbon Contaminated Soil and Groundwater. *Environ. Sci. Technol.* **2005**, *39*, 7279–7286. [CrossRef] [PubMed]
115. Haritash, A.K.; Kaushik, C.P. Biodegradation Aspects of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *J. Hazard. Mater.* **2009**, *169*, 1–15. [CrossRef] [PubMed]
116. Starzyńska-Janiszewska, A. Kwas γ-Poliglutaminowy–Bakteryjny Biopolimer z Natto. Available online: https://biomist.pl/biologia/mikroorganizmy/kwas-%CE%B3-poliglutaminowy-bakteryjny-biopolimer-natto/7657 (accessed on 15 March 2021).
117. Wang, G.; Liu, Q.; Wang, Y.; Li, J.; Chen, Y.; Wen, Q.; Zheng, D.; Kang, W.; Quan, H. The Application and Functional Progress of γ-Poly-Glutamic Acid in Food: A Mini-Review. *CPD* **2020**, *26*, 5347–5352. [CrossRef] [PubMed]
118. Luo, Z.; Guo, Y.; Liu, J.; Qiu, H.; Zhao, M.; Zou, W.; Li, S. Microbial Synthesis of Poly-γ-Glutamic Acid: Current Progress, Challenges, and Future Perspectives. *Biotechnol. Biofuels* **2016**, *9*, 134. [CrossRef] [PubMed]
119. Zhang, L.; Yang, X.; Gao, D.; Wang, L.; Li, J.; Wei, Z.; Shi, Y. Effects of Poly-γ-Glutamic Acid (γ-PGA) on Plant Growth and Its Distribution in a Controlled Plant-Soil System. *Sci. Rep.* **2017**, *7*, 6090. [CrossRef] [PubMed]

materials

MDPI

Suplementary materials

Evaluation of the Effectiveness of the Biopreparation in Com-Bination with the Polymer PGA for the Biodegradation of Soils Contaminated with Petroleum Hydrocarbons

Katarzyna Wojtowicz [1,*], Teresa Steliga [1], Piotr Kapusta [1], Joanna Brzeszcz [1] and Tomasz Skalski [2]

[1] Oil and Gas Institute—National Research Institute, Lubicz 25 A, 31-503 Krakow, Poland; teresa.steliga@inig.pl (T.S.); piotr.kapusta@inig.pl (P.K.); joanna.brzeszcz@inig.pl (J.B.)
[2] Biotechnology Centre, Silesian University of Technology, Krzywoustego 8, 44-100 Gliwice, Poland; tomasz.skalski@polsl.pl
* Correspondence: katarzyna.wojtowicz@inig.pl




Figure S1. Content (%) of individual components of petroleum pollutants in the soil. (**a**) TPH; (**b**) BTEX; (**c**) PAHs.

Ecotoxicological analyses–description of the methodology

Phytotoxkit™ test (MicroBioTests Inc., Nazareth, Belgium)

Chronic toxicity assessment test *Phytotoxkit™* is based on the evaluation of germination and early growth of plants (root elongation inhibition measurement. Three types of plants selected according to germination rate and root growth rate are used in the test, which allows making a complete determination in 3 days of incubation: monocotyledonous - sorghum *(Sorghum saccharatum)*, and dicotyledonous-cress *(Lepidium sativum)* and white mustard *(Sinapis alba)*. The determination process was carried out using three repetitions for each test plant. The tests were performed on transparent polystyrene test plates. *Incubation conditions:* temperature T = 25 °C in darkness, incubation time t = 72 h. *Test reaction:* inhibition of germination and early root growth [1–3].

The inhibition in seed germination and root growth was calculated according to a following equation:

$$I = \frac{(A - B)}{B} \cdot 100\%$$

where: I—inhibition (%), A—seed germination or root length in control soil, B—seed germination or root length in test soil

Ostracodtoxkit(F)™ (MicroBioTest Inc., Nazareth, Belgium)
Ostracodtoxkit(F) test [4] belongs to direct contact tests of chronic toxicity estimation with the use of crustaceans *Heterocypris incongruens*. The direct contact test is done with the application of young bottom crustaceans (*Heterocypris incongruens*), which hatched from cysts during 52 h (according to the producer's procedure). The test is carried out on 6-hole polystyrene microplates. A standardised nutrient is an algae suspension, which is applied to the suspension in portions of 2 ml. Portions of 10 ostracods are added to each cell of the microplate. 0.5 g of a control soil is applied to the row A, whereas 0.5 g of the soil sample is added to the consecutive rows. Incubation lasts for 6 days in a temperature of 25 °C. Results reading consists of estimation of life microforms amount in each hole of the microplate and measurement of their length. Growth inhibition of *H. incongruens* was calculated as

$$GI = 100 - \left(\frac{A}{B} \cdot 100\%\right)$$

where: A—ostracods length in tested soil, B—ostracods length in reference soil.

Microtox® Solid Phase Test (SDI, Newark, DE, USA)
Microtox Test created in the USA in 1979 as the first bioindication test, combines typical bioindication and analytic precision. Luminescence bacteria *Vibro fischeri*, which uses about 10% of its metabolism in order to emit light, was used as a bioindicator. In the presence of toxical substances, there is decrease in luminescence and increase in general toxicity of a sample. In an electron transportation system of the bacteria, lucipherasa enzyme (alcane oxygenaze) catalyses oxidation of a reduced substrate (reduced flavine mononucleotide, riboflavin phosphate or flavine adenin dinucleotide) and during this process luminescence, which can be measured with a photometer, takes place. The obtained substrates of this reaction are oxygen and long-chain aldehyde. In the presence of substances that have negative influence on cell metabolism, the decrease in luminescence of bacteria is immediate.
Test Microtox®, produced by SDI Company (USA), enables a direct contact of luminescence bacteria *Vibro fischeri* with a wastewater sample, which leads not only to determination of substances dissolved in water, but also to recognition of lipophilicity systems and poorly dissolved systems in water. Lyophilised bacteria *Vibro fischeri* can be stored in a period of a year in temperature of -20°C and then used for testing immediately after suspending in deionised water. A tests with serial dilutions of the wastewater were done according to a standard procedure and results were obtain in Delta Tox analyser. In this analysis, the concentration of wastewater, which reduces the luminescence by 50%, was determined (half maximal effective concentration, EC_{50}). In this presentation a lower of EC_{50} denotes a higher toxicity. To simplify this interpretation, EC_{50} values were converted to Toxicity Units (TU) in which a higher value denotes a higher toxicity according to the following formula [5–9].

$$\mathrm{TU} = \frac{1}{EC_{50}} \cdot 100$$

AMES (Muta-ChromoPlate™ Kit (EBPI, Mississauga, ON, Canada).
Numerous petroleum contaminants and metabolites created in biodegradation of pollutants include carcinogenic compounds. In order to estimate potentially mutagenic properties of the wastewater, Ames Test was applied. It enables determination of reverse mutations from histidine auxotrophy to prototrophy in mutants, which are test strains of *Salmonella typhimurium*. The test strains of the bacteria can be characterised by various types of mutation in genes, which are responsible for histidine synthesis. Therefore, the presence of histidine in a substratum is required in order to provide the growth of the strains. Generally, TA98 and TA100 strains are used [9–11]. They include proper mutations consisting of so-called reading frame displacement, pair base substitution and two mutagenes (2-nitro-fluoren and sodium

azide). The test strains are mutants, which are unable to histidine synthesis. A mutagenous substance causes reversion in mutation and histidine synthesis can take place, which results in appearance of revertants colonies in the histidine-free basis.

In order to lead Ames Test, a high-tech microplate AMES MPF test was applied. It can be used for detection of genotoxic activity of water contaminated with petroleum substances and metabolites, which are the effects of biodegradation. Delivered microforms of T-98 and T-100 *Salmonella typhimurium* undergo strict quality control (genotype and phenotype). The test results in colour reaction, where yellow means reverse mutation, whereas scarlet—the lack of mutation.




Figure S2. Scheme of Ames Test conduct [12].

Petroleum contaminants from the samples were extracted with the use of dichloromethane, which was then vaporised and the obtained deposit was dissolved in dimethylosulfoxide (DMSO) [10]. As an activator, S9 microsomal fraction from a rat liver was used. The role of the activator is to transform the tested substance, which leads to disclosure of its potentially mutagenous properties.

References

1. Baran, A.; Tarnawski, M. Phytotoxkit/Phytotestkit and Microtox® as Tools for Toxicity Assessment of Sediments. *Ecotoxicology and Environmental Safety* **2013**, *98*, 19–27, doi:10.1016/j.ecoenv.2013.10.010.
2. Blinova, I.; Bityukova, L.; Kasemets, K.; Ivask, A.; Käkinen, A.; Kurvet, I.; Bondarenko, O.; Kanarbik, L.; Sihtmäe, M.; Aruoja, V.; et al. Environmental Hazard of Oil Shale Combustion Fly Ash. *Journal of Hazardous Materials* **2012**, *229–230*, 192–200, doi:10.1016/j.jhazmat.2012.05.095.
3. Mamindy-Pajany, Y.; Hamer, B.; Roméo, M.; Géret, F.; Galgani, F.; Durmiši, E.; Hurel, C.; Marmier, N. The Toxicity of Composted Sediments from Mediterranean Ports Evaluated by Several Bioassays. *Chemosphere* **2011**, *82*, 362–369, doi:10.1016/j.chemosphere.2010.10.005.
4. Niyommaneerat, W.; Nakajima, F.; Tobino, T.; Yamamoto, K. Development of a Chronic Sediment Toxicity Test Using the Benthic Ostracod Heterocypris Incongruens and Their Application to Toxicity Assessments of Urban Road Dust. *Ecotoxicology and Environmental Safety* **2017**, *143*, 266–274, doi:10.1016/j.ecoenv.2017.05.011.
5. Lima, T.M.S.; Procópio, L.C.; Brandão, F.D.; Leão, B.A.; Tótola, M.R.; Borges, A.C. Evaluation of Bacterial Surfactant Toxicity towards Petroleum Degrading Microorganisms. *Bioresource Technology* **2011**, *102*, 2957–2964, doi:10.1016/j.biortech.2010.09.109.

6. Adams, R.H., Kanga-Leyva, K., Guzmán-Osorio, F.J., Escalante-Espinoza, E, .2011. Comparison of moisture management methods for the bioremediation of hydrocarbon contaminated soil. *African Journal of Biotechnology* **2011**, *10*, 394-404.
7. Foucault, Y.; Durand, M.-J.; Tack, K.; Schreck, E.; Geret, F.; Leveque, T.; Pradere, P.; Goix, S.; Dumat, C. Use of Ecotoxicity Test and Ecoscores to Improve the Management of Polluted Soils: Case of a Secondary Lead Smelter Plant. *Journal of Hazardous Materials* **2013**, *246–247*, 291–299, doi:10.1016/j.jhazmat.2012.12.042.
8. Oleszczuk, P.; Jośko, I.; Kuśmierz, M.; Futa, B.; Wielgosz, E.; Ligęza, S.; Pranagal, J. Microbiological, Biochemical and Ecotoxicological Evaluation of Soils in the Area of Biochar Production in Relation to Polycyclic Aromatic Hydrocarbon Content. *Geoderma* **2014**, *213*, 502–511, doi:10.1016/j.geoderma.2013.08.027.
9. Kamber, M.; Fluckiger-Isler, S.; Engelhardt, G.; Jaeckh, R.; Zeiger, E. Comparison of the Ames II and Traditional Ames Test Responses with Respect to Mutagenicity, Strain Specificities, Need for Metabolism and Correlation with Rodent Carcinogenicity. *Mutagenesis* **2009**, *24*, 359–366, doi:10.1093/mutage/gep017.
10. Steliga, T.; Jakubowicz, P.; Kapusta, P. Changes in Toxicity during in Situ Bioremediation of Weathered Drill Wastes Contaminated with Petroleum Hydrocarbons. Bioresource Technology 2012, 125, 1–10, doi:10.1016/j.biortech.2012.08.092.
11. Vijay, U.; Gupta, S.; Mathur, P.; Suravajhala, P.; Bhatnagar, P. Correction: Microbial Mutagenicity Assay: Ames Test. *BIO-PROTOCOL* **2018**, *8*, doi:10.21769/BioProtoc.3016.
12. Available online: http://www.tigret.eu/images/stories/produkty/Toksykologia/Ames_MPF_PL_20191206.pdf (accessed on 15 March 2021).

sustainability

MDPI

Article

Influence of Biosurfactants on the Efficiency of Petroleum Hydrocarbons Biodegradation in Soil

Katarzyna Wojtowicz [1], Teresa Steliga [1,*], Tomasz Skalski [2] and Piotr Kapusta [3]

[1] Department of Production Technology of Reservoir Fluids, Oil and Gas Institute—National Research Institute, Lubicz 25 A, 31-503 Kraków, Poland; katarzyna.wojtowiczk@inig.pl
[2] Biotechnology Centre, Silesian University of Technology, Krzywoustego 8, 44-100 Gliwice, Poland; tomasz.skalski@polsl.pl
[3] Department of Microbiology, Oil and Gas Institute—National Research Institute, Lubicz 25 A, 31-503 Kraków, Poland; piotr.kapusta@inig.pl
* Correspondence: teresa.steliga@inig.pl

Abstract

Soil contamination with petroleum hydrocarbons is a serious environmental issue, necessitating the development of effective and environmentally friendly remediation methods that align with the principles of sustainable development. This study investigated the impact of selected biosurfactants on the efficiency of the biodegradation of total petroleum hydrocarbons (TPH) and polycyclic aromatic hydrocarbons (PAHs) in contaminated soil. Six biosurfactants—poly-γ-glutamic acid (γ-PGA), rhamnolipid, surfactin, a mixture of γ-PGA, rhamnolipids, and surfactin (PSR), as well as two commercial formulations (JBR 425 and JBR 320)—were evaluated in combination with a bacterial consortium. Biodegradation experiments were conducted under laboratory conditions for a 90-day period. The effectiveness of the tested biosurfactants was assessed using respirometric analysis, the chromatographic determination of the residual hydrocarbon content, and toxicity assays. The results showed that the application of a bacterial consortium enriched with a mixture of biosurfactants PSR (a biosurfactant concentration in the inoculating mixture: 5 g/dm^3) was the most effective approach, resulting in an oxygen uptake of 5164.8 mgO_2/dm^3 after 90 days, with TPH and PAH degradation rates of 77.3% and 70.32%, respectively. Phytotoxicity values decreased significantly, with TU values ranging from 6.32 to 4.62 (growth inhibition) and 3.77 to 4.13 (germination). Toxicity also decreased in the ostracodtoxkit test (TU = 4.35) and the Microtox SPT test (TU = 4.91). Among the tested biosurfactants, surfactin showed the least improvement in its bioremediation efficiency. Under the same concentration as in the PSR mixture, the oxygen uptake was 3446.7 mgO_2/dm^3, with TPH and PAH degradation rates of 60.64% and 52.64%, respectively. In the system inoculated with the bacterial consortium alone (without biosurfactants), the biodegradation efficiency reached 44.35% for TPH and 36.97% for PAHs. The results demonstrate that biosurfactants can significantly enhance the biodegradation of petroleum hydrocarbons in soil, supporting their potential application in sustainable bioremediation strategies.

Keywords: biodegradation; bioremediation; biosurfactants; ecotoxicological assessment; microbial consortium; petroleum hydrocarbons

check for updates

Academic Editors: Roberto Mancinelli, Yutao Peng and Chungyu Guan

Received: 18 June 2025
Revised: 9 July 2025
Accepted: 15 July 2025
Published: 16 July 2025

Citation: Wojtowicz, K.; Steliga, T.; Skalski, T.; Kapusta, P. Influence of Biosurfactants on the Efficiency of Petroleum Hydrocarbons Biodegradation in Soil. *Sustainability* **2025**, *17*, 6520. https://doi.org/10.3390/su17146520

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

The contamination of the environment by petroleum hydrocarbons represents a significant environmental concern in the contemporary world [1]. The primary sources of

these pollutants are the activities of the petrochemical industry, oil spills, and the improper storage and transportation of fuels [2]. Consequently, substantial areas of soil and water have been contaminated with toxic compounds that exhibit a high degree of persistence in the environment [3]. These pollutants disrupt the structure and functioning of ecosystems, leading to a decline in biodiversity, alteration of biogeochemical cycles, and toxicity to soil organisms, plants, and animals [4,5].

Petroleum hydrocarbons, such as alkanes, cycloalkanes, monocyclic aromatic hydrocarbons (BTEX), and polycyclic aromatic hydrocarbons (PAHs), are hydrophobic, poorly biodegradable, and toxic compounds [6]. PAHs are of particular concern due to their mutagenic and carcinogenic properties [7,8]. The presence of these substances in the soil has been shown to result in a reduced oxygen availability, the inhibition of microbial activity, and changes in the physicochemical properties of the soil, thereby rendering it less capable of self-cleaning [9,10]. The remediation of petroleum hydrocarbon-contaminated soils is a multifaceted process that employs a range of physicochemical techniques, including thermal treatment [11], solvent extraction [12], solidification/stabilization (S/S) [13], photocatalysis [14], and oxidation processes [15]. While these approaches can be effective, they are often associated with high operational costs, require specialized equipment, and demand a stringent control of reaction conditions, which limits their feasibility for large-scale applications [16]. Consequently, biological methods such as bioremediation have gained increasing attention due to their environmental compatibility, cost-effectiveness, and efficiency under natural field conditions, aligning well with the principles of sustainability and environmentally responsible practices.

Bioremediation, defined as the treatment of the environment with microorganisms capable of breaking down toxic substances, is a highly effective method for the removal of oil pollution [17]. The process is based on the utilization of naturally occurring or introduced microorganisms (bioaugmentation) that facilitate the conversion of hydrocarbons into less harmful products, primarily carbon dioxide, water, and biomass [18]. The efficiency of this process can be significantly increased by the use of biosurfactants, defined as natural surfactant compounds produced by microorganisms [19]. It is evident that, as of 2022, the majority of microorganisms capable of producing biosurfactants are classified into a limited number of well-defined taxonomic groups [20]. Bacteria of the genus *Pseudomonas* constitute the majority of this group, accounting for up to 37% of all identified biosurfactant producers. Of particular note is *Pseudomonas aeruginosa*, which is distinguished by its efficient synthesis of rhamnolipids [21]. These biosurfactants are among the most extensively studied, exhibiting potent surfactant and emulsifying properties. The second most significant group is the *Bacillus* genus, comprising 34% of known biosurfactant producers [20]. It has been demonstrated that species such as *Bacillus subtilis* and *Bacillus licheniformis* are capable of producing biosurfactants [22], which have been shown to be efficacious in the remediation of hydrocarbon-contaminated environments. Among yeasts, a notable proportion (12%) is attributed to representatives of the *Candida* genus, including *Candida bombicola* [23]. Other groups of microorganisms are of lesser quantitative importance, although they are, nevertheless, important from the perspective of biological and metabolic diversity. Bacteria from the *Serratia* genus [24] and the filamentous fungus *Rhizopus arrhizus* [25] each account for 3% of the total number of known biosurfactant producers. The remaining 11% comprise a variety of strains, including consortia of microorganisms that can act synergistically in biosynthesis and biodegradation processes, rendering them a promising area for further biotechnological research [20]. Biosurfactant molecules are composed of both hydrophilic and lipophilic parts [26]. The hydrophilic groups are typically mono- or polysaccharides, amino acids, or peptides, while the hydrophobic part is constituted by long-chain fatty acids, fatty hydroxy acids, or α-alkyl-β-hydroxy acids. The structure

of biosurfactants facilitates their localization at the phase boundary (e.g., oil/water), a process which has been demonstrated to reduce interfacial tensions and increase the contact surface between contaminants and microorganisms [27,28]. In contrast to synthetic surfactants, biosurfactants are characterized by their biodegradability [29], non-toxicity, and stability over a broad spectrum of pH, temperature, and salinity, thereby enhancing their environmental compatibility [30–32]. Biosurfactants can be classified on the basis of their chemical structure, adopting division into five main groups: glycolipids, lipoproteins and lipopeptides, phospholipids and fatty acids, biopolymers, and biological structures [33]. Glycolipids include compounds such as rhamnolipids, mannolipids, sophorolipids, trehalose lipids, and diglucosyl diglycerides [33–35]. Lipoproteins and lipopeptides include compounds like surfactin [36], viscosin, lichenysin, and gramicidin [35]. Phospholipids and fatty acids are represented by compounds such as spiculisporic acid and corynomycolic acids, which differ in the length of their carbon chains. Biopolymers include emulsan, biodispersan, allasan, liposan, polysaccharide–protein complexes, mannoproteins, and carbohydrate–protein complexes [33,35]. Additionally, biosurfactants may occur in the form of biological structures such as vesicles or entire cells that exhibit surface-active properties. An alternative classification of biosurfactants, commonly found in the literature, is based on their molecular weight [37]. This approach distinguishes low-molecular-weight biosurfactants, up to approximately 1500 Da, which mainly include glycolipids (such as rhamnolipids, sophorolipids, and trehalose lipids), lipopeptides (e.g., surfactin, gramicidin S, and polymyxin), as well as certain small proteins [37,38]. In contrast, high-molecular-weight biosurfactants (up to approximately 100 kDa) comprise polysaccharides, lipoproteins, and other complex polymeric molecules [37,39]. This classification reflects not only structural differences, but also the diverse physicochemical and functional properties of these compounds. Biosurfactants have been demonstrated to enhance the biodegradation efficiency by means of several mechanisms [40]. Firstly, biosurfactants can improve the solubility of hydrocarbons in water [41,42]. Secondly, they can mobilize contaminants from the surface of soil particles [43–45]. Thirdly, they can stimulate microbial growth and activity by serving as a carbon source and supplying essential nutrients necessary for proper microbial development [20,45,46]. Moreover, biosurfactants can reduce the toxic effects of hydrocarbons on microbial cells by altering cell surface hydrophobicity and membrane permeability, thereby increasing the cell's tolerance to hydrophobic compounds [47]. Furthermore, biosurfactants have been shown to facilitate the emulsification of hydrophobic organic compounds, thereby significantly enhancing the rate of their biological decomposition [48–51].

2. Materials and Methods

2.1. Materials

The soil in this area has been heavily contaminated with petroleum hydrocarbons, with a TPH fraction concentration of 6062.59 mg/kg d.m. and a polycyclic aromatic hydrocarbon (PAH) content of 12.05 mg/kg d.m. The study material was sourced from the site of a former waste pit located in southeastern Poland. Soil samples were collected from two depth intervals of the near-surface layer: the samples were collected at depths ranging from 0 to 0.25 m and from 0.25 to 1.00 m. Subsequent to collection, the samples were meticulously placed in hermetically sealed, appropriately labeled containers and expedited to the laboratory for immediate analysis. In the context of laboratory experimentation, the samples were subjected to rigorous mixing and homogenization procedures to ensure the generation of a representative test material for subsequent analysis [52].

The bioremediation study utilized a bacterial consortium (MC), which was developed on the basis of indigenous microorganisms isolated from oil hydrocarbon-contaminated

waste pits. The MC consortium comprised the following bacterial strains: *Dietzia* sp. IN118, *Gordonia* sp. IN101, *Mycolicibacterium frederiksbergense* IN53 (formerly *Mycobacterium frederiksbergense* IN53), *Rhodococcus erythropolis* IN119, *Rhodococcus globerulus* IN113, and *Raoultella* sp. IN109. The composition and method of construction of the consortium has been described in detail in earlier works [52,53].

Five biosurfactants of commercial availability and one mixture developed in the laboratory were utilized in the ongoing study of surfactant-assisted bioremediation. Each formulation was tested at two concentrations: 1 g/dm^3 and 5 g/dm^3. The following biosurfactants were included in the analysis: γ-polyglutamate (γ-PGA) (Ambioteco, Polska), rhamnolipids (Sigma-Aldrich, Burlington, MA, USA), surfactin (Sigma-Aldrich, Burlington, MA, USA), and commercial formulations JBR 425 (Janeil Biotechnology, Saukville, WI, USA) and JBR 320 (Janeil Biotechnology, Saukville, WI, USA). Furthermore, a mixture designated PSR was applied comprising γ-PGA, rhamnolipids, and surfactin, prepared at a volume ratio of 1:1:1.

2.2. Methods

Experiment Description

A series of studies were conducted to explore the potential of bioremediation in addressing soil contamination by petroleum hydrocarbons. This research was conducted under controlled laboratory conditions, utilizing the Oxi-Top (WTW, Weilheim, Germany) kit as a primary experimental apparatus. Experiments were conducted on soil samples of 30 g, contaminated with petroleum hydrocarbons (TPH) and polycyclic aromatic hydrocarbons (PAH) under strictly controlled conditions. The temperature was maintained at 20 °C, the soil moisture content at 20%, and the air access and duration of the experiment was 90 days.

In the course of this study, a total of 26 experimental variants were prepared, the distinguishing features of which were the presence of the MC bacterial consortium, the type of biosurfactant utilized, and its respective concentration. The biosurfactants which were the focus of the present study included poly-γ-glutamic acid (γ-PGA), surfactin, rhamnolipids, biosurfactant preparations, JBR 425 and JBR 320, as well as a proprietary mixture designated PSR, consisting of γ-PGA, surfactin, and rhamnolipids in a 1:1:1 ratio. Variants 1–12 comprised samples containing both the bacterial consortium MC and biosurfactant formulations. For each biosurfactant type, measurements were taken at two concentrations (1 g/dm^3 and 5 g/dm^3). The remaining control variants (13–24) contained only the respective biosurfactants at concentrations of 1 g/dm^3 or 5 g/dm^3, without the consortium. Variant 25 was a control sample that was not inoculated, while variant 26 was inoculated with the bacterial consortium MC. Figure 1 shows the experimental design.

The experiment was performed in five replicates. During the course of this study, measurements of oxygen consumption were carried out in the analyzed sample enrichment variants (respirometric analyses), with the aim of reflecting microbial activity in the variants. The detailed procedure of the respirometric analyses is described in the Supplementary Materials. Subsequent to the completion of the measurements, the soil samples were subjected to a process of mixing and homogenization in order to obtain a sample that would be representative of the soil in question for the purposes of this study as a whole. This process included chromatographic analyses and ecotoxicological assessments.

The quantification of the TPH content in soil samples was conducted by gas chromatography, which was performed using a Clarus 500 chromatograph (PerkinElmer, Waltham, MA, USA) [52,53]. The analysis of the PAHs content was conducted by high-performance liquid chromatography (HPLC) on a Vanquish Core Series instrument (Thermo Scientific, Waltham, MA, USA) [54]. Concurrently, an ecotoxicological assessment was conducted

employing standardized Phytotoxkit, Ostracodtoxkit, and Microtox SPT toxicological assays. The detailed procedures for the respirometric and chromatographic analyses and ecotoxicological tests are described in the authors' previous publications [52] and in the Supplementary Materials (Tables S1 and S2).




Figure 1. Scheme of the bioremediation experiment in selected purification process variants. [1] Variant numbers of the purification process are marked on the bottles. [2] A detailed description of the samples corresponding to each variant is provided in the Supplementary Materials.

2.3. Data Analysis and Statistical Information

The statistical analysis was conducted using Statistica 14.0 (TIBCO Software Inc., Palo Alto, CA, USA). Prior to further analysis, the data were tested for normality. Subsequently, one-way analysis of variance (ANOVA) was performed and in instances where significant differences were identified, Tukey's post hoc test was employed for multiple comparisons. Standard deviation (SD), relative standard deviation (RSD, %), and Pearson correlation coefficients were also calculated. Statistical significance was considered at $p < 0.05$ [52,53].

3. Results

3.1. Respirometric Tests

In the 90-day respirometric tests conducted, the microbial activity in the soil samples was evaluated on the basis of oxygen consumption, expressed in mgO_2/dm^3. The results demonstrated significant variations between the analyzed experimental variants (Figure 2).




Figure 2. The temporal variation in oxygen consumption measured in soil samples: (**a**) with biosurfactant (1 g/L) or bacterial consortium MC and biosurfactant (1 g/L), (**b**) with biosurfactant (5 g/L) or the bacterial consortium MC and biosurfactant (5 g/L) (repetition number $n = 5$, $p < 0.05$).

In the blank sample (non-inoculated sample), minimal oxygen consumption (15.51 mgO_2/dm^3) was recorded, confirming the absence of biological activity in this sample. Conversely, in the control sample, comprising soil inoculated with the bacterial consortium, a considerably elevated level of oxygen consumption (1944.8 ± 71.95 mgO_2/dm^3) was detected, signifying an augmentation in microbial activity, associated with the biodegradation of petroleum pollutants by the bacterial strains present in the consortium. The incorporation of selected biosurfactants at a concentration of 1 g/dm^3 into the bacterial consortium resulted in a substantial augmentation in the oxygen consumption within the examined samples. The highest values were recorded for the biosurfactant mixture of PSR (4131.9 ± 148.24 mgO_2/dm^3), JBR425 (3885.3 ± 142.12 mgO_2/dm^3), and JBR320 (3752.8 ± 133.33 mgO_2/dm^3). The bacterial consortium exhibited suboptimal results when enriched with γ-PGA and rhamnolipids, exhibiting oxygen consumption levels of 3486.6 ± 128.31 mgO_2/dm^3 and 3467.6 ± 125.39 mgO_2/dm^3, respectively, after a 90-day period. Conversely, the inoculating mixture that was enriched with surfactin resulted in a variant oxygen consumption of 3332.3 ± 121.08 $mgO_2/dm3$. Research undertaken in analogous conditions on soil samples enrichment only with biosurfactants (i.e., without consortium) at a concentration of 1 g/dm^3 demonstrated significantly diminished oxygen consumption during the experiment. The results of the experiment revealed that the presence of microorganisms is crucial for efficient biodegradation processes, as evidenced by the following measurements: 1081.6 ± 40.69 mgO_2/dm^3 (surfactin), 1052.2 ± 38.95 mgO_2/dm^3 (PSR mixture), 1446.2 ± 53.95 mgO_2/dm^3 (JBR425), and 1281.3 ± 47.40 mgO_2/dm^3 (JBR320) (Figure 2a). The results obtained from samples inoculated with a bacterial consortium enriched in biosurfactants and biosurfactants alone indicate a synergitic effect between microorganisms and biosurfactants, resulting in an intensification of the biological activity (Table S2 in Supplementary Materials).

It was established that an increase in the concentration of biosurfactants in the bacterial consortium to 5 g/dm^3 resulted in a further increase in microbial activity. In a manner analogous to the studies conducted on lower surfactant concentrations, the high-

est oxygen consumption values were observed in the variant inoculated with the bacterial consortium enriched with a mixture of PSR (5164.8 ± 188.85 mgO_2/dm^3), JBR425 (5087.2 ± 182.10 mgO_2/dm^3), and JBR320 (4956.2 ± 179.25 mgO_2/dm^3). The incorporation of γ-PGA, surfactin, and rhamnolipids into the inoculation mixture resulted in oxygen consumption levels of 4440.2 ± 161.36 mgO_2/dm^3, 3446.7 ± 125.22 mgO_2/dm^3 and 4258.3 ± 155.47 mgO_2/dm^3, respectively, in the tested variants after a 90-day period. The values of oxygen consumption in the control samples (following the introduction of biosurfactants alone into the soil at a concentration of 5 g/dm^3) ranged from 1273.2 ± 41.59 to 1749.8 ± 65.77 mgO_2/dm^3 (Figure 2b).

3.2. Chromatographic Analysis

Chromatographic analyses demonstrated that the incorporation of biosurfactants within the inoculation mixture resulted in a substantial enhancement of the biodegradation efficiency of petroleum hydrocarbons within the soil environment. The findings unequivocally demonstrate that the type of surfactant employed, as well as its concentration, exert a substantial influence on the extent of degradation of contaminants. Figure 3 provides a summary of the biodegradation levels of TPH and PAH in the examined variants.




Figure 3. Percentage of TPH and PAH biodegradation in the tested experimental setups, including: a control sample (non-inoculated), a sample inoculated with a bacterial consortium, samples inoculated with the consortium and supplemented with a biosurfactant, and a sample enriched only with the biosurfactant at concentrations of (**a**) 1 g/dm^3 and (**b**) 5 g/dm^3, (repetition number $n = 7–10$, $p < 0.05$).

Tests conducted on a blank sample (non-inoculated sample) showed marginal degradation of the compounds (0.2% TPH and 0.05% PAH), confirming that the spontaneous treatment of soil contaminated with TPH and PAH under these conditions is minimal and requires the support of bioremediation processes. The inoculation of the soil with the bacterial consortium resulted in a decrease in the concentration of TPH from 6062.59 mg/kg dry matter (DM) to 3375.26 mg/kg DM (44.33%) and in the concentration of polycyclic aromatic hydrocarbons (PAH) from 12.05 mg/kg DM to 7.60 mg/kg DM (36.97%). These results indicate the ability of microorganisms to degrade contaminants independently, albeit with less efficiency than in variants enriched with biosurfactants.

The addition of biosurfactants to the bacterial consortium resulted in a substantial enhancement in the efficiency of petroleum substance removal from soil. The highest level of TPH and PAH degradation was achieved for the sample inoculated with the consortium MC and the PSR biosurfactant mixture. At a PSR concentration of 5 g/dm^3, a decline in the

concentration of TPH to 1376.34 mg/kg s.m. (77.30%) and PAH to 3.58 mg/kg DM (70.32%) was observed, representing the most optimal outcome among all the variants evaluated. Variants inoculated with the bacterial consortium enriched with JBR425 and JBR320 at a concentration of 5 g/dm^3 demonstrated equivalent levels of efficiency. The final concentrations for variant 11 were as follows: the TPH concentration was found to be 1554.68 mg/kg, DM (74.36%), while the PAH concentration was determined to be 4.14 mg/kg, DM (65.68%). In variant 12, the TPH concentration was found to be 1641.94 mg/kg DM (72.92%), while the PAH concentration was determined to be 4.37 mg/kg DM (63.71%). The utilization of reduced concentrations of biosurfactants within the inoculating mixture yielded diminished biodegradation efficiency; nevertheless, the outcomes attained following a three-month treatment period remained satisfactory. In soil inoculated with the bacterial consortium enriched with a PSR mixture at a concentration of 1 g/dm^3, the concentrations of TPH and PAH were 2067.09 mg/kg DM and 4.64 mg/kg DM, respectively, corresponding to degradation rates: the mean values for TPH and PAH were found to be 65.90% and 61.53%, respectively. In samples enriched with the bacterial consortium, along with the addition of JBR425 and JBR320 at a concentration of 1 g/dm^3, the degradation efficiency of TPH was observed to range from 61.29% to 69.45%, while that of PAH was found to be between 56.20% and 58.87%.

Tests conducted on samples to which only biosurfactants were added demonstrated a significantly diminished biodegradation efficiency. For samples containing biosurfactants at a concentration of 5 g/dm^3, the range of TPH degradation was from 12.13% to 19.32%, and the range of PAH degradation was from 6.84% to 11.74%. These findings suggest that the use of biosurfactants alone is not as effective as might have been expected. Analogous trends were observed at lower concentrations (1 g/dm^3), where TPH degradation did not exceed 16.32%, and PAH degradation was 10.15%.

A comprehensive analysis of data pertaining to the degradation of TPH across a range of carbon chain lengths (n-C6-n-C9, n-C10-n-C21, n-C22-n-C30, and n-C31-n-C36) has been conducted. The analysis revealed a discernible correlation between the efficiency of biodegradation and the chemical structure of the pollutants. It has been demonstrated that hydrocarbons with shorter carbon chains (n-C6-n-C9 and n-C10-n-C21) are subject to biodegradation at a faster rate than heavier fractions containing longer carbon chains (n-C22-n-C30 and n-C31-n-C36). Figure 4 presents the average degradation percentage of n-alkanes with different carbon chain lengths across various experimental variants.

The highest degradation values were recorded for the n-C10-n-C21 fraction, in which the level of reduction in samples inoculated with the bacterial consortium with the addition of biosurfactants at a concentration of 5 g/dm^3 usually exceeded 90%. The n-C6-n-C9 fraction, which contains light hydrocarbons, also exhibited a notably elevated level of biodegradation (in the range of 75.28–92.62%), despite its share of the total TPH mass typically being less than that of the medium alkane fractions. For the n-C22-n-C30 fraction, degradation values were typically in the range of 61.94–79.71% in variants inoculated with the bacterial consortium, with the addition of biosurfactants (5 g/dm^3). Conversely, for the n-C31-n-C36 fraction, the degradation level rarely exceeded 50%, even in the most efficient variants.

The findings obtained from the experiments involving samples inoculated with biosurfactant-enriched samples at reduced concentrations (1 g/dm^3) and the samples enriched exclusively with biosurfactants substantiate this correlation. However, it is noteworthy that the biodegradation rates in the various fractions were considerably diminished in these instances.




Figure 4. Average degradation percentages of n-alkanes with different carbon chain lengths in soil across various experimental setups (*X*-axis): (**a**) with biosurfactant concentration in the inoculating solution of 1 g/dm^3, (**b**) with biosurfactant concentration of 5 g/dm^3 (repetition number n = 7–10, $p < 0.05$).

In a manner analogous to aliphatic hydrocarbons, a clear relationship was also observed for PAHs between the biodegradation efficiency and the structure of the molecule or, more specifically, the number of aromatic rings. Low-molecular-weight PAHs (containing two-to-three aromatic rings per molecule) have been shown to biodegrade more efficiently than high-molecular-weight PAHs (containing four or more aromatic rings per molecule). Variants incorporating the bacterial consortium and biosurfactants at a concentration of 5 g/dm^3 exhibited the highest degradation efficiencies for low-ring PAHs, with naphthalene demonstrating the greatest degree of degradation (66.11–88.32%). For three-ring PAHs, the values ranged from 49.81 to 66.16%. For the same variants, the degree of degradation of five- and six-ring PAHs was noticeably lower, ranging from 25.84 to 38.22%. In the variants with a lower concentration of biosurfactants (1 g/dm^3), the values of average biodegradation degrees were correspondingly lower, and for low-molecular-weight PAHs, they were in the range of 41.33–77.28%, while for high-molecular-weight PAHs, they were in the range of 21.44–36.60%. In control variants enriched exclusively with biosurfactants, the level of PAH biodegradation was found to be significantly lower for both low- and high-molecular-weight PAHs. However, it was also observed that the biodegradation spheres exhibited a dependence on the number of aromatic rings present in the hydrocarbon molecule. Figure 5 shows the distribution of the average PAH degradation values across different experimental variants (*X*-axis), categorized by the number of aromatic rings in the hydrocarbon molecules.




Figure 5. Average degradation percentages of PAHs with different numbers of aromatic rings in the molecules across various experimental variants (*X*-axis): (**a**) with biosurfactant concentration in the inoculating solution of 1 g/dm^3, (**b**) with biosurfactant concentration of 5 g/dm^3 (repetition number $n = 7$–10, $p < 0.05$).

In evaluating the efficacy of the implemented remediation treatments, the values of the biodegradation coefficients (n-C17/Pr and n-C18/Ph) constitute a pivotal component of the analysis. It was observed that the enrichment of the bacterial consortium with biosurfactants led to a significant reduction in the values of these coefficients compared to the samples inoculated with the bacterial consortium only. This finding suggests that biosurfactants have the capacity to enhance the bioavailability of hydrocarbon contaminants to microorganisms, thereby increasing biodegradation processes. In comparison, the samples containing only biosurfactants exhibited higher values of biodegradation coefficients, which may indicate their limited effectiveness in the absence of microbial activity. The values of the biodegradation coefficients in the soil after the bioremediation process in the tested variants are presented in Table 1.

3.3. *Ecotoxicological Assessment*

The evaluation of the effectiveness of the applied remediation treatments was supplemented by an analysis of soil ecotoxicity after the treatment process in the tested variants. For the purpose of this study, the Phytotoxkit test was employed and the resulting data were converted into toxicity index (TU) values. A summary of the TU values in the Phytotoxkit test obtained for soil samples after bioremediation was completed in the tested variants of the process conduct is provided in Figure 6.

Table 1. Summary of biodegradation rates of n-C17/Pr and n-C18/Ph in soil after bioremediation in the tested variants at biosurfactant concentrations of 1 g/dm^3 and 5 g/dm^3 in the inoculation solution (repetition number n = 7–10, $p < 0.05$).

Sample	n-C17/Pr	n-C18/Ph
Initial soil G6 (control sample)	2.716 ± 0.076	2.575 ± 0.075
Soil G6 + bacterial consortium	1.247 ± 0.027	1.303 ± 0.023
The concentration of biosurfactant in the enriching mixture is 1 g/dm^3		
Soil G6 + MC + γ-PGA	0.781 ± 0.009	0.818 ± 0.010
Soil G6 + MC + surfactin	0.959 ± 0.011	1.004 ± 0.016
Soil G6 + MC + rhamnolipids	0.845 ± 0.015	0.936 ± 0.026
Soil G6 + MC + mixture PSR	0.604 ± 0.016	0.665 ± 0.015
Soil G6 + MC + JBR425	0.802 ± 0.012	0.808 ± 0.022
Soil G6 + MC + JBR320	0.871 ± 0.011	0.925 ± 0.025
Soil G6 + γ-PGA	2.286 ± 0.056	2.350 ± 0.060
Soil G6 + surfactin	2.241 ± 0.059	2.282 ± 0.058
Soil G6 + rhamnolipids	2.273 ± 0.053	2.247 ± 0.053
Soil G6 + mixture PSR	2.192 ± 0.052	2.227 ± 0.057
Soil G6 + JBR425	2.125 ± 0.055	2.232 ± 0.052
Soil G6 + JBR320	2.172 ± 0.048	2.199 ± 0.049
The concentration of biosurfactant in the enriching mixture is 5 g/dm^3		
Soil G6 + MC + γ-PGA	0.257 ± 0.007	0.267 ± 0.003
Soil G6 + MC + surfactin	0.688 ± 0.012	0.719 ± 0.011
Soil G6 + MC + rhamnolipids	0.364 ± 0.006	0.408 ± 0.008
Soil G6 + MC + mixture PSR	0.164 ± 0.004	0.173 ± 0.003
Soil G6 + MC + JBR425	0.243 ± 0.007	0.252 ± 0.008
Soil G6 + MC + JBR320	0.273 ± 0.003	0.284 ± 0.006
Soil G6 + γ-PGA	2.286 ± 0.056	2.225 ± 0.055
Soil G6 + surfactin	2.251 ± 0.059	2.392 ± 0.052
Soil G6 + rhamnolipids	2.184 ± 0.046	2.235 ± 0.045
Soil G6 + mixture PSR	2.108 ± 0.042	2.109 ± 0.041
Soil G6 + JBR425	2.233 ± 0.043	2.211 ± 0.043
Soil G6 + JBR320	2.266 ± 0.056	2.193 ± 0.057

The analyses demonstrated that the enrichment of the MC bacterial consortium with biosurfactants had a clear effect on reducing the soil toxicity, as evidenced by the analysis of key test parameters, including seed germination and root length inhibition, in test organisms (*Sinapis alba*, *Sorghum saccharatum*, and *Lepidium sativum*). For the initial (contaminated) soil sample, prior to remediation, the toxicity coefficient values for germination ranged from 9.02 to 11.24, while for root length inhibition, the values ranged from 8.70 to 12.56. The lowest G6 soil TU values were observed in the variants inoculated with the MC bacterial consortium enriched with the PSR mixture and JBR425 and JBR320 preparations, with a biosurfactant concentration in the inoculating mixture of 5 g/dm^3. In these variants, the post-experiment soil TU values for plant germination were 3.77 for *Sinapis alba* in the variant with PSR, 4.05 with JBR425, and 4.19 with JBR320, respectively. For *Sorghum saccharatum*, the values were 4.13, 4.49, and 4.65, respectively, while for *Lepidium sativum*, they were 4.58, 4.92, and 5.08. In the case of root growth inhibition, reductions in soil TU were observed for *Sinapis alba*, with values of 3.62 (MC + PSR), 3.89 (MC + JBR425), and 4.02 (MC + JBR320). For *Sorghum saccharatum*, the corresponding values were 4.19, 4.52, and 4.67, while for *Lepidium sativum* they were 4.62, 5.01, and 5.20, respectively (Figure 7b). For the lower concentration of biosurfactants in the mixture with the bacterial consortium of 1 g/dm^3, slightly higher toxicity coefficient values were observed. For germination,

the G6 soil TU values against *Sinapis alba* were 4.79 in the PSR variant, 5.16 with JBR425, and 5.31 with JBR320. For *Sorghum saccharatum*, the corresponding values were 5.44, 5.92, and 6.11, respectively. For *Lepidium sativum*, the values were marginally higher at 5.83, 6.29, and 6.46, respectively. For root growth inhibition, G6 soil TU values against *Sinapis alba* were 4.61 (MC + PSR), 4.97 (MC + JBR425), and 5.11 (MC + JBR320), respectively, for *Sorghum saccharatum* 5.39, 5.83, and 6.00, and for Lepidium sativum, 6.07, 6.62, and 6.83, respectively. Surfactin was found to be the least effective biosurfactant in this study. Its presence in the inoculating mixture, at a concentration of 1 g/dm^3, resulted in a decline in G6 soil TU, ranging from 5.75 to 7.01 (germination) and 5.53 to 7.48 (root length inhibition). An increase in the surfactin concentration to 5 g/dm^3 yielded only marginally superior outcomes; TU values decreased by 5.20–6.35 (germination) and 5.01–6.69 (root length inhibition) (Figure 7a).

Tests conducted on control samples enriched only in biosurfactants, without the MC bacterial consortium, showed only a slight decrease in soil toxicity to test organisms. The toxicity coefficient values for these samples for germination ranged between 7.94 and 10.29, and for root growth inhibition, between 7.66 and 11.57. The findings of this study demonstrate that the variants incorporating a blend of bacterial consortium and biosurfactants emerged as the most efficacious in mitigating soil toxicity. The efficacy of this mitigation was found to be contingent upon both the nature of the biosurfactant employed and its concentration.

Another test employed to evaluate the ecotoxicity of the soil following the treatment process in the variants examined was the Ostracodtoxkit test, in which the test organism is *Heterocypris incongruens*. In a manner analogous to the Phytotoxkit test, the results of the test, including the mortality and growth inhibition of bivalve mollusks, were converted into toxicity index (TU) values. A summary of the TU values in the Ostracodtoxkit test obtained for soil samples after bioremediation was completed in the tested variants is provided in Figure 7a.

The initial soil (G6) exhibited the highest toxicity value (TU) of 10.5, signifying the most pronounced toxic effect against the bioindicators employed among all the samples examined. The inoculation of the soil with the bacterial consortium (MC) alone resulted in a decrease in toxicity to TU = 8.05, suggesting partial detoxification by microbial action. The lowest TU values (and, therefore, the lowest toxicity) were obtained after the simultaneous application of the bacterial consortium MC and biosurfactants. When the soil was inoculated with the consortium enriched with biosurfactants at a concentration of 5 g/dm^3, a range of TU values was observed, ranging from 4.35 to 6.42. The lowest toxicity was found for the combination of the bacterial consortium MC + PSR mixture, and the highest for bacterial consortium MC + surfactin. Slightly higher TU values were obtained with a lower concentration of biosurfactants (1 g/dm^3), whose values ranged from 5.81 to 7.18. In soil samples inoculated with biosurfactants only, total toxicity values (TU) remained high, ranging from 9.79 to 10.15, indicating that biosurfactants used in isolation did not significantly reduce the soil toxicity.

In order to complement the ecotoxicological analysis, an evaluation of the toxicity of soil samples was conducted using the Microtox SPT test. In this test, the indicator organism is the bioluminescent bacterium *Vibrio fischeri*.




Figure 6. Summary of toxicity unit (TU) values obtained in the Phytotoxkit phytotoxicity test for the examined variants, with the application of biosurfactants at concentrations of: (**a**) 1 g/dm^3 and (**b**) 5 g/dm^3, (repetition number n = 7–10, $p < 0.05$).




Figure 7. Summary of toxicity unit (TU) values obtained in (**a**) the Ostracodtoxkit and (**b**) the Microtox test for the examined variants, (repetition number $n = 7–10$, $p < 0.05$).

As with the other tests, the initial soil sample (G6) exhibited the highest level of toxicity (TU = 13.05), indicating the presence of substances that are harmful to aquatic organisms. Inoculating the soil with the MC bacterial consortium alone resulted in a marked decrease in toxicity (TU = 8.87) due to the bacteria breaking down xenobiotics. Significantly lower TU values were recorded in samples inoculated with both the MC bacterial consortium and biosurfactants at a concentration of 5 g/dm^3. The TU values in this group ranged from 4.91 to 7.05. The soil sample with the lowest toxicity level was inoculated with the MC bacterial consortium enriched with the PSR mixture (TU = 4.91), and for the G6 soil sample, inoculated with the bacterial consortium with the addition of surfactin (TU = 7.05). At a lower concentration of biosurfactants (1 g/dm^3) in the inoculating mixture, the TU values of the soil samples ranged from 6.42 to 7.87. Tests conducted on control samples enriched only with biosurfactants at concentrations of 1 and 5 g/dm^3 showed a slight decrease in TU values relative to the initial sample. These values were in the ranges of 11.6–12.05 and 11.33–11.98, respectively, confirming the low effectiveness of biosurfactants alone in treating soils contaminated with petroleum hydrocarbons. A summary of the Microtox SPT test results for all soil samples after the experiment is provided in Figure 7b.

4. Discussion

Contamination of the soil and groundwater environment with petroleum hydrocarbons, including fractions of TPH and PAHs, is one of the key challenges of modern environmental protection [55]. Due to their persistence, low solubility, and high affinity for organic matter, these compounds exhibit limited bioavailability, making their biodegradation much more difficult [56,57]. In response to these challenges, combinatorial methods, such as, among others, biosurfactant-assisted bioaugmentation, are receiving increasing attention [58]. The employment of meticulously selected consortia of microorganisms, endowed with the capacity to degrade pollutants, in conjunction with biosurfactants that augment the bioavailability of hydrophobic compounds, has the potential to markedly enhance the efficacy of bioremediation processes [31,59]. These methods have been proven to enhance the efficiency of contaminant elimination whilst concurrently fostering sustainable development by minimizing the environmental impact, eschewing the use of deleterious chemicals,

curtailing secondary pollution, and preserving ecosystem integrity. Biosurfactant-assisted techniques exemplify the balance between technological efficiency and environmental responsibility, rendering them particularly relevant in modern, sustainability-oriented environmental management.

The experimental results obtained unequivocally confirm the effectiveness of using a biosurfactant-assisted bioaugmentation strategy in the bioremediation of soils historically contaminated with petroleum hydrocarbons. The application of the selected bacterial consortium, comprising strains from the genera *Rhodococcus*, *Dietzia*, *Mycolicibacterium*, *Gordonia*, and *Raoultella*, to soil through inoculation resulted in a substantial enhancement of the degradation parameters, particularly in the presence of biosurfactants (γ-PGA, surfactin, rhamnolipids, the PSR mixture, and the commercially available preparations JBR425 and JBR320). This process led to an increase in the bioavailability of the contaminants.

Respirometric and chromatographic analyses indicated that biosurfactants played a key role in enhancing the efficiency of the biodegradation of oil-derived pollutants by microorganisms present in the MC consortium. In the course of respirometric analyses, the highest oxygen consumption, which is an indicator of intensive microbial activity, was recorded in samples enriched with the MC consortium and PSR mixture, which contain three biosurfactants (γ-PGA, surfactin, and rhamnolipids), and MC with the commercial preparations JBR425 and JBR320. Conversely, the lowest level of activity was exhibited by the variants containing the MC consortium and surfactin. The findings suggest a clear relationship between the capacity of TPH and PAH biodegradation and the type of biosurfactant employed. It can be assumed that mixtures of different surfactants, characterized by a more complex composition, interact more effectively with the diverse microorganisms present in the bacterial consortium, providing a broader spectrum of nutrients necessary for their growth and activity. Such conditions promote the intensification of metabolic processes of microflora in complex systems, which in turn may result in the increased bioavailability of contaminants and accelerated biodegradation [60,61]. This observation is corroborated by the finding that, in variants inoculated with the bacterial consortium alone or with bio-surfactants alone, significantly lower oxygen consumption was observed during the experiment. The observations made during respirometric analyses were corroborated by the results of chromatographic analyses, which were used to determine the concentrations of petroleum hydrocarbons (TPH, aliphatic hydrocarbons, and PAHs) in the initial soil sample and in the soil samples after bioremediation in the tested treatment variants. The optimal conditions for achieving maximum hydrocarbon biodegradation were determined to be variants augmented with the bacterial consortium and biosurfactants. The application of biosurfactants resulted in a substantial decrease in the concentrations of TPH and PAHs in the soil, in comparison to the baseline soil sample and the control sample inoculated solely with the biosurfactant. The highest efficiency was obtained with the PSR biosurfactant mixture at a higher concentration (5 g/dm^3) in combination with the MC consortium, with the reduction of contaminants exceeding 70%, indicating a strong synergistic effect of these components. It was also observed that equivalent results were attained with reduced concentrations of PSR, though the efficacy was marginally diminished. Conversely, samples with surfactin exhibited reduced biodegradation efficiency, particularly at lower concentrations, indicating the limited efficacy of this biosurfactant under the examined conditions. For the purpose of comparison, the use of the bacterial consortium alone allowed for a moderate reduction in contaminants. Conversely, experiments with biosurfactants alone, without the participation of microorganisms, showed the lowest efficiency, especially with regard to TPH and PAHs, where reductions did not exceed 20% and 12%, respectively. The findings of the present research were consistent with those reported by others. For instance, in a study conducted by Zhang, the application of rhamnolipids in PAH-contaminated

soil increased the degradation efficiency by more than 50% compared to the control sample [62]. A further study demonstrated that the incorporation of surfactin resulted in the acceleration of the degradation of phenanthrene and pyrene in PAH-contaminated soils [63]. Owsianiak et al. noted that the use of rhamnolipids in the biodegradation of hydrocarbons increased the degradation efficiency to 77% after 7 days, compared to 58% in a sample without biosurfactants [64]. In turn, Zeng et al. showed that the use of monoramnolipids significantly improved hexadecane biodegradation in a *Candida* yeast monoculture, achieving 93% degradation after 4 days, while only 78% was achieved in the sample without biosurfactant [65]. Husein's observations confirmed the conjecture that biosurfactant mixtures allow better pyrene degradation efficiency. In their study, they showed that the use of a mixture of rhamnolipids, emulsane, and indigenous biosurfactants resulted in 98% pyrene degradation, while the absence of emulsane in the biosurfactant mixture resulted in a reduction in pyrene biodegradation to 91% [66,67]. These results are in line with our observations, suggesting the advantage of biosurfactant mixtures over their individual forms, due to a possible synergistic effect. In their study, Lai et al. compared the effectiveness of various biosurfactants, including surfactin and rhamnolipids, in removing total petroleum hydrocarbons (TPH) from soils contaminated with petroleum hydrocarbons. At the same concentrations of biosurfactants, the efficiency of TPH removal from soil for rhamnolipids was in the range of 23–63%, while for surfactin, it was in the range of 14–62%, indicating that surfactin was less effective [68]. The present study investigates the relationship between the intensity of biodegradation processes and the concentration of the biosurfactant in the inoculating mixture. This relationship was investigated through the conducting of tests with two variants of biosurfactant concentrations (1 g/dm^3 and 5 g/dm^3). However, it is important to consider whether the achieved biodegradation effect is commensurate with the increased financial outlay associated with a higher biosurfactant concentration. It is important to note that high efficiency does not necessarily equate to the optimization of the process from the point of view of operating costs. This is a particularly salient consideration in the context of industrial-scale implementations of bioremediation technologies. The experimental findings suggest that the optimal equilibrated dosage of biosurfactants can significantly impact the efficiency and cost-effectiveness of the overall bioremediation process by biosurfactant-enhanced bioaugmentation [69].

Furthermore, the findings of chromatographic analyses demonstrated a correlation between the chemical structure of organic pollutants and their biodegradability [56,70]. This relationship was observed for aliphatic carbon donors, where fractions with shorter carbon chains (n-C6-n-C9 and n-C10-n-C21) were degraded much faster than long-chain fractions (n-C22-n-C29 and n-C30-n-C36) [71–73]. A comparable phenomenon was observed in the context of polycyclic aromatic hydrocarbons, wherein compounds comprising fewer aromatic rings per molecule (<four aromatic rings per molecule) exhibited significantly higher biodegradability in comparison to compounds containing four or more aromatic rings [4,74–76]. The underlying mechanisms responsible for this trend can be elucidated by examining the physicochemical properties of the constituent compounds. Shorter-chain aliphatic hydrocarbons and low-molecular-weight PAHs exhibit higher volatility, enhanced water solubility, and greater bioavailability to microorganisms [77]. Consequently, these compounds are more rapidly transported into microbial cells and are more easily metabolized. In contrast, higher-molecular-weight hydrocarbons exhibit reduced mobility, diminished solubility, and an increased propensity for a strong affinity with the organic fraction of the soil or sediment. This, in turn, limits their bioavailability and consequently slows their rate of biodegradation [56,57,78,79]. The employment of biosurfactants, which enhance the solubility and bioavailability of contaminants, in conjunction with the introduction of selected consortia of microorganisms with high metabolic activity to a given type of

compound [31,80], thus ensures optimal treatment outcomes. The effectiveness of biodegradation was also confirmed by an analysis of the n-C17/Pr and n-C18/Ph indices [80–82]. In the variants containing biosurfactants at a concentration of 5 g/dm^3, the values of these indices fell into the ranges of 0.16–0.70 and 0.17–0.74, respectively, compared to the values in soil inoculated only with the consortium (1.22 and 1.28). This finding suggests that the presence of biosurfactants can significantly enhance the rate of biodegradation. In the control samples, the values remained close to the baseline, confirming their limited effectiveness without the participation of microorganisms.

Ecotoxicological analyses, conducted utilizing Phytotoxkit, Ostracodtoxkit, and Microtox SPT tests, constituted a significant component in the comprehensive evaluation of the efficacy of the bioremediation process. The lowest toxicity (TU) values were obtained in variants in which the bacterial consortium and biosurfactants (especially PSR, JBR425, and JBR320 at a concentration of 5 g/dm^3) were applied simultaneously. TU values for phytotoxicity parameters (germination and root growth) fell into the range of 3.62–4.61, in the Ostracod-toxkit test to 4.35, while in the Mcrotx SPT test to 4.91, which is a significant reduction from the baseline values (8.70–12.56, 10.5, and 13.05, respectively). A further significant finding is that no secondary toxic by-products were identified in the tested variants following the remediation process. Concurrently, the findings substantiated the notion of a direct correlation between the concentration of TPH and PAHs and soil toxicity [69,83,84]. The outcomes of both chromatographic analyses and toxicological tests indicated that a decrease in the pollutant concentration was proportionate to a decrease in the TU values. Furthermore, the variants comprising of solely biosurfactants, devoid of microbial involvement, did not attain substantial enhancement in the soil quality parameters. This underscores the imperativeness for a synergistic interplay between biosurfactants and microorganisms in achieving efficacious environmental purification.

5. Conclusions

The accumulated data clearly indicate that the use of a composite method such as biosurfactant-assisted biodegradation is an effective strategy in the remediation of soils contaminated with petroleum hydrocarbons. The findings from the respirometric, chromatographic, and toxicological analyses substantiate that this integrated approach considerably enhances the biodegradation efficiency of both total petroleum hydrocarbons (TPH) and polycyclic aromatic hydrocarbons (PAHs). Furthermore, the toxicity unit (TU) values obtained in Phytotoxkit, Ostracodtoxkit, and Microtox tests indicate an enhancement in the overall quality of the soil environment and a reduction in potential ecological risks. The findings substantiate the assertion that the judicious selection of the bacterial consortium, in congruence with a biosurfactant, is paramount to the efficacy of remediation processes. This approach is a sustainable and economically viable solution for the remediation of industrial sites, supporting long-term environmental recovery while minimizing harmful side effects, which is in line with the core objectives of sustainable development.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/su17146520/s1. A description of the samples is presented in Figure 1; Methods description; Table S1: characteristics of bioindication ecotoxicological tests applied for toxicity assessment; Table S2: synergistic index (SI) for different experimental setups and biosurfactant concentrations.

Author Contributions: Conceptualization, K.W., T.S. (Teresa Steliga) and P.K.; Methodology, K.W., T.S. (Teresa Steliga), T.S. (Tomasz Skalski) and P.K.; Validation, K.W., T.S. (Teresa Steliga), T.S. (Tomasz Skalski) and P.K.; Formal analysis, K.W., T.S. (Teresa Steliga), T.S. (Tomasz Skalski) and P.K.; Investi-

gation, K.W.; Writing—original draft preparation, K.W.; Writing—Review and Editing, K.W. and P.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was financially supported by the Polish Ministry of Science and Higher Education within statutory funding for the Oil and Gas Institute—the National Research Institute DK:4100-19-23.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Dataset available on request from the authors.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Song, X.; Wu, X.; Song, X.; Shi, C.; Zhang, Z. Sorption and Desorption of Petroleum Hydrocarbons on Biodegradable and Nondegradable Microplastics. *Chemosphere* **2021**, *273*, 128553. [CrossRef] [PubMed]
2. Ambaye, T.G.; Chebbi, A.; Formicola, F.; Prasad, S.; Gomez, F.H.; Franzetti, A.; Vaccari, M. Remediation of Soil Polluted with Petroleum Hydrocarbons and Its Reuse for Agriculture: Recent Progress, Challenges, and Perspectives. *Chemosphere* **2022**, *293*, 133572. [CrossRef] [PubMed]
3. Mekonnen, B.A.; Aragaw, T.A.; Genet, M.B. Bioremediation of Petroleum Hydrocarbon Contaminated Soil: A Review on Principles, Degradation Mechanisms, and Advancements. *Front. Environ. Sci.* **2024**, *12*, 1354422. [CrossRef]
4. Sharma, K.; Shah, G.; Singh, H.; Bhatt, U.; Singhal, K.; Soni, V. Advancements in Natural Remediation Management Techniques for Oil Spills: Challenges, Innovations, and Future Directions. *Environ. Pollut. Manag.* **2024**, *1*, 128–146. [CrossRef]
5. Boopathy, R. Factors Limiting Bioremediation Technologies. *Bioresour. Technol.* **2000**, *74*, 63–67. [CrossRef]
6. Claro, E.M.T.; Cruz, J.M.; Montagnolli, R.N.; Lopes, P.R.M.; Júnior, J.R.M.; Bidoia, E.D. Microbial Degradation of Petroleum Hydrocarbons: Technology and Mechanism. In *Microbial Action on Hydrocarbons*; Kumar, V., Kumar, M., Prasad, R., Eds.; Springer: Singapore, 2018; pp. 125–141. ISBN 9789811318399.
7. Speight, J.G. Organic Chemistry. In *Environmental Organic Chemistry for Engineers*; Elsevier: Amsterdam, The Netherlands, 2017; pp. 43–86, ISBN 9780128044926.
8. Abdel-Shafy, H.I.; Mansour, M.S.M. A Review on Polycyclic Aromatic Hydrocarbons: Source, Environmental Impact, Effect on Human Health and Remediation. *Egypt. J. Pet.* **2016**, *25*, 107–123. [CrossRef]
9. Kakde, P.; Sharma, J. Microbial Bioremediation of Petroleum Contaminated Soil: Structural Complexity, Degradation Dynamics and Advanced Remediation Techniques. *J. Pure Appl. Microbiol.* **2024**, *18*, 2244–2261. [CrossRef]
10. Koshlaf, E.; Ball, A.S.; Koshlaf, E.; Ball, A.S. Soil Bioremediation Approaches for Petroleum Hydrocarbon Polluted Environments. *AIMS Microbiol.* **2017**, *3*, 25–49. [CrossRef] [PubMed]
11. Vidonish, J.E.; Zygourakis, K.; Masiello, C.A.; Sabadell, G.; Alvarez, P.J.J. Thermal Treatment of Hydrocarbon-Impacted Soils: A Review of Technology Innovation for Sustainable Remediation. *Engineering* **2016**, *2*, 426–437. [CrossRef]
12. Silva, A.; Deleruematos, C.; Fiuza, A. Use of Solvent Extraction to Remediate Soils Contaminated with Hydrocarbons. *J. Hazard. Mater.* **2005**, *124*, 224–229. [CrossRef] [PubMed]
13. Roslee, N.F.; Kamil, N.A.F.M.; Kadir, A.A.; Jalil, A.R.; Hamzah, N.; Noor, N.M.; Sandu, A.V. New Ray on Remediation of High Rings Polycyclic Aromatic Hydrocarbons: Remediation of Raw Petroleum Sludge Using Solidification and Stabilization Method. *Arch. Metall. Mater.* **2022**, *67*, 1201–1208. [CrossRef]
14. Diao, Z.-H.; Pu, S.-Y.; Qian, W.; Liang, S.; Kong, L.-J.; Xia, D.-H.; Lei, Z.-X.; Du, J.-J.; Liu, H.; Yang, J.-W. Photocatalytic Removal of Phenanthrene and Algae by a Novel Ca-Ag3PO4 Composite under Visible Light: Reactivity and Coexisting Effect. *Chemosphere* **2019**, *221*, 511–518. [CrossRef] [PubMed]
15. Arroyo, S.; Rosano-Ortega, G.; Martínez-Gallegos, S.; Pérez-Armendariz, B.; Vega-Lebrún, C.A. Reduction of Hydrocarbons in Contaminated Soil through Paired Sorption and Advanced Oxidation Processes. *Soil Secur.* **2021**, *4*, 100013. [CrossRef]
16. Smith, J.W.N. Debunking Myths about Sustainable Remediation. *Remediat. J.* **2019**, *29*, 7–15. [CrossRef]
17. Kuppan, N.; Padman, M.; Mahadeva, M.; Srinivasan, S.; Devarajan, R. A Comprehensive Review of Sustainable Bioremediation Techniques: Eco Friendly Solutions for Waste and Pollution Management. *Waste Manag. Bull.* **2024**, *2*, 154–171. [CrossRef]
18. Elshafei, A.M.; Mansour, R. Microbial Bioremediation of Soils Contaminated with Petroleum Hydrocarbons. *Discov. Soil* **2024**, *1*, 9. [CrossRef]

19. Parus, A.; Ciesielski, T.; Woźniak-Karczewska, M.; Ślachciński, M.; Owsianiak, M.; Ławniczak, Ł.; Loibner, A.P.; Heipieper, H.J.; Chrzanowski, Ł. Basic Principles for Biosurfactant-Assisted (Bio)Remediation of Soils Contaminated by Heavy Metals Petroleum Hydrocarbons—ACritical Evaluation of the Performance of Rhamnolipids. *J. Hazard. Mater.* **2023**, *443*, 130171. [CrossRef] [PubMed]
20. Eras-Muñoz, E.; Farré, A.; Sánchez, A.; Font, X.; Gea, T. Microbial Biosurfactants: A Review of Recent Environmental Applications. *Bioengineered* **2022**, *13*, 12365–12391. [CrossRef] [PubMed]
21. Henkel, M.; Müller, M.M.; Kügler, J.H.; Lovaglio, R.B.; Contiero, J.; Syldatk, C.; Hausmann, R. Rhamnolipids as Biosurfactants from Renewable Resources: Concepts for next-Generation Rhamnolipid Production. *Process Biochem.* **2012**, *47*, 1207–1219. [CrossRef]
22. Gudiña, E.J.; Teixeira, J.A. Bacillus Licheniformis: The Unexplored Alternative for the Anaerobic Production of Lipopeptide Biosurfactants? *Biotechnol. Adv.* **2022**, *60*, 108013. [CrossRef] [PubMed]
23. Da Silva, R.R.; Santos, J.C.V.; Meira, H.M.; Almeida, S.M.; Sarubbo, L.A.; Luna, J.M. Microbial Biosurfactant: Candida Bombicola as a Potential Remediator of Environments Contaminated by Heavy Metals. *Microorganisms* **2023**, *11*, 2772. [CrossRef] [PubMed]
24. Clements, T.; Ndlovu, T.; Khan, S.; Khan, W. Biosurfactants Produced by *Serratia* Species: Classification, Biosynthesis, Production and Application. *Appl. Microbiol. Biotechnol.* **2019**, *103*, 589–602. [CrossRef] [PubMed]
25. Pele, M.A.; Ribeaux, D.R.; Vieira, E.R.; Souza, A.F.; Luna, M.A.C.; Rodríguez, D.M.; Andrade, R.F.S.; Alviano, D.S.; Alviano, C.S.; Barreto-Bergter, E.; et al. Conversion of Renewable Substrates for Biosurfactant Production by *Rhizopus arrhizus* UCP 1607 and Enhancing the Removal of Diesel Oil from Marine Soil. *Electron. J. Biotechnol.* **2019**, *38*, 40–48. [CrossRef]
26. Badmus, S.O.; Amusa, H.K.; Oyehan, T.A.; Saleh, T.A. Environmental Risks and Toxicity of Surfactants: Overview of Analysis, Assessment, and Remediation Techniques. *Environ. Sci. Pollut. Res.* **2021**, *28*, 62085–62104. [CrossRef] [PubMed]
27. Bjerk, T.R.; Severino, P.; Jain, S.; Marques, C.; Silva, A.M.; Pashirova, T.; Souto, E.B. Biosurfactants: Properties and Applications in Drug Delivery, Biotechnology and Ecotoxicology. *Bioengineering* **2021**, *8*, 115. [CrossRef] [PubMed]
28. Dini, S.; Bekhit, A.E.-D.A.; Roohinejad, S.; Vale, J.M.; Agyei, D. The Physicochemical and Functional Properties of Biosurfactants: A Review. *Molecules* **2024**, *29*, 2544. [CrossRef] [PubMed]
29. Patel, S.; Kharawala, K. Biosurfactants Their Biodegradability: A Review Examination. *Int. J. Eng. Adv. Technol.* **2022**, *11*, 4–11. [CrossRef]
30. Romero Vega, G.; Gallo Stampino, P. Bio-Based Surfactants and Biosurfactants: An Overview and Main Characteristics. *Molecules* **2025**, *30*, 863. [CrossRef] [PubMed]
31. Banat, I.M.; Franzetti, A.; Gandolfi, I.; Bestetti, G.; Martinotti, M.G.; Fracchia, L.; Smyth, T.J.; Marchant, R. Microbial Biosurfactants Production, Applications and Future Potential. *Appl. Microbiol. Biotechnol.* **2010**, *87*, 427–444. [CrossRef] [PubMed]
32. Kugaji, M.; Ray, S.K.; Parvatikar, P.; Raghu, A.V. Biosurfactants: A Review of Different Strategies for Economical Production, Their Applications and Recent Advancements. *Adv. Colloid Interface Sci.* **2025**, *337*, 103389. [CrossRef] [PubMed]
33. GreyViews Global Biosurfactants Market Growth 2021–2028. Available online: https://greyviews.com/reports/global-biosurfactants-market/25 (accessed on 2 June 2025).
34. Mnif, I.; Ellouz-Chaabouni, S.; Ghribi, D. Glycolipid Biosurfactants, Main Classes, Functional Properties and Related Potential Applications in Environmental Biotechnology. *J. Polym. Environ.* **2018**, *26*, 2192–2206. [CrossRef]
35. Ahmadi-Ashtiani, H.-R.; Baldisserotto, A.; Cesa, E.; Manfredini, S.; Sedghi Zadeh, H.; Ghafori Gorab, M.; Khanahmadi, M.; Zakizadeh, S.; Buso, P.; Vertuani, S. Microbial Biosurfactants as Key Multifunctional Ingredients for Sustainable Cosmetics. *Cosmetics* **2020**, *7*, 46. [CrossRef]
36. Phulpoto, I.A.; Yu, Z.; Hu, B.; Wang, Y.; Ndayisenga, F.; Li, J.; Liang, H.; Qazi, M.A. Production and Characterization of Surfactin-like Biosurfactant Produced by Novel Strain Bacillus Nealsonii S2MT and It's Potential for Oil Contaminated Soil Remediation. *Microb. Cell Factories* **2020**, *19*, 145. [CrossRef] [PubMed]
37. Rosenberg, E.; Ron, E.Z. High- and Low-Molecular-Mass Microbial Surfactants. *Appl. Microbiol. Biotechnol.* **1999**, *52*, 154–162. [CrossRef] [PubMed]
38. Lourenço, M.; Duarte, N.; Ribeiro, I.A.C. Exploring Biosurfactants as Antimicrobial Approaches. *Pharmaceuticals* **2024**, *17*, 1239. [CrossRef] [PubMed]
39. Simões, C.R.; Da Silva, M.W.P.; De Souza, R.F.M.; Hacha, R.R.; Merma, A.G.; Torem, M.L.; Silvas, F.P.C. Biosurfactants: An Overview of Their Properties, Production, and Application in Mineral Flotation. *Resources* **2024**, *13*, 81. [CrossRef]
40. Sah, D.; Rai, J.P.N.; Ghosh, A.; Chakraborty, M. A Review on Biosurfactant Producing Bacteria for Remediation of Petroleum Contaminated Soils. *3 Biotech* **2022**, *12*, 218. [CrossRef] [PubMed]
41. Bordoloi, N.K.; Konwar, B.K. Bacterial Biosurfactant in Enhancing Solubility and Metabolism of Petroleum Hydrocarbons. *J. Hazard. Mater.* **2009**, *170*, 495–505. [CrossRef] [PubMed]
42. Souza, E.C.; Vessoni-Penna, T.C.; De Souza Oliveira, R.P. Biosurfactant-Enhanced Hydrocarbon Bioremediation: An Overview. *Int. Biodeterior. Biodegrad.* **2014**, *89*, 88–94. [CrossRef]

43. Kuyukina, M.S.; Ivshina, I.B.; Makarov, S.O.; Litvinenko, L.V.; Cunningham, C.J.; Philp, J.C. Effect of Biosurfactants on Crude Oil Desorption and Mobilization in a Soil System. *Environ. Int.* **2005**, *31*, 155–161. [CrossRef] [PubMed]
44. Pacwa-Płociniczak, M.; Płaza, G.A.; Piotrowska-Seget, Z.; Cameotra, S.S. Environmental Applications of Biosurfactants: Recent Advances. *Int. J. Mol. Sci.* **2011**, *12*, 633–654. [CrossRef] [PubMed]
45. Markam, S.S.; Raj, A.; Kumar, A.; Khan, M.L. Microbial Biosurfactants: Green Alternatives and Sustainable Solution for Augmenting Pesticide Remediation and Management of Organic Waste. *Curr. Res. Microb. Sci.* **2024**, *7*, 100266. [CrossRef] [PubMed]
46. Ron, E.Z.; Rosenberg, E. Biosurfactants and Oil Bioremediation. *Curr. Opin. Biotechnol.* **2002**, *13*, 249–252. [CrossRef] [PubMed]
47. Kaczorek, E.; Pacholak, A.; Zdarta, A.; Smułek, W. The Impact of Biosurfactants on Microbial Cell Properties Leading to Hydrocarbon Bioavailability Increase. *Colloids Interfaces* **2018**, *2*, 35. [CrossRef]
48. Bustamante, M.; Durán, N.; Diez, M.C. Biosurfactants Are Useful Tools for the Bioremediation of Contaminated Soil: A Review. *J. Soil Sci. Plant Nutr.* **2012**, *12*, 667–687. [CrossRef]
49. Karlapudi, A.P.; Venkateswarulu, T.C.; Tammineedi, J.; Kanumuri, L.; Ravuru, B.K.; Dirisala, V.R.; Kodali, V.P. Role of Biosurfactants in Bioremediation of Oil Pollution-a Review. *Petroleum* **2018**, *4*, 241–249. [CrossRef]
50. Zhuang, X.; Wang, Y.; Wang, H.; Dong, Y.; Li, X.; Wang, S.; Fan, H.; Wu, S. Comparison of the Efficiency and Microbial Mechanisms of Chemical- and Bio-Surfactants in Remediation of Petroleum Hydrocarbon. *Environ. Pollut.* **2022**, *314*, 120198. [CrossRef] [PubMed]
51. Mnif, I.; Sahnoun, R.; Ellouz-Chaabouni, S.; Ghribi, D. Application of Bacterial Biosurfactants for Enhanced Removal and Biodegradation of Diesel Oil in Soil Using a Newly Isolated Consortium. *Process Saf. Environ. Prot.* **2017**, *109*, 72–81. [CrossRef]
52. Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J. Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (*Zea mays*). *Molecules* **2023**, *28*, 6104. [CrossRef] [PubMed]
53. Wojtowicz, K.; Steliga, T.; Kapusta, P. Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using Echinacea Purpurea. *Appl. Sci.* **2023**, *13*, 13077. [CrossRef]
54. Wojtowicz, K. Opracowanie Metodyki Oznaczania WWA w Próbkach Gleb z Wykorzystaniem Chromatografii Cieczowej HPLC. *Nafta-Gaz* **2022**, *78*, 141–153. [CrossRef]
55. Logeshwaran, P.; Megharaj, M.; Chadalavada, S.; Bowman, M.; Naidu, R. Petroleum Hydrocarbons (PH) in Groundwater Aquifers: An Overview of Environmental Fate, Toxicity, Microbial Degradation and Risk-Based Remediation Approaches. *Environ. Technol. Innov.* **2018**, *10*, 175–193. [CrossRef]
56. Haritash, A.K.; Kaushik, C.P. Biodegradation Aspects of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *J. Hazard. Mater.* **2009**, *169*, 1–15. [CrossRef] [PubMed]
57. Gundlapalli, M.; Sivagami, K.; Gopalakrishnan, M.; Harshini, P.; Janjaroen, D.; Ganesan, S. Biodegradation of Low Molecular Weight Polycyclic Aromatic Hydrocarbons in Soil: Insights into Bacterial Activities and Bioremediation Techniques. *Sustain. Chem. Environ.* **2024**, *7*, 100146. [CrossRef]
58. Vaishnavi, J.; Parthipan, P.; Arul Prakash, A.; Sathishkumar, K.; Rajasekar, A. Biosurfactant-Assisted Bioremediation of Crude Oil/Petroleum Hydrocarbon Contaminated Soil. In *Handbook of Assisted and Amendment: Enhanced Sustainable Remediation Technology*; Prasad, M.N.V., Ed.; Wiley, John Wiley & Sons, Inc.: Hoboken, NJ, USA, 2021; pp. 313–329. ISBN 9781119670360.
59. Wojtowicz, K.; Steliga, T. Wpływ Mieszaniny Biosurfaktantów Na Efektywność Bioremediacji Gruntu Zanieczyszczonego Węglowodorami Ropopochodnymi. *Nafta-Gaz* **2024**, *80*, 269–278. [CrossRef]
60. Abdoli, S.; Asgari Lajayer, B.; Bagheri Novair, S.; Price, G.W. Unlocking the Potential of Biosurfactants in Agriculture: Novel Applications and Future Directions. *Sustainability* **2025**, *17*, 2110. [CrossRef]
61. Abo Elsoud, M.M.A. Classification and Production of Microbial Surfactants. In *Microbial Biosurfactants*; Inamuddin Ahamed, M.I., Prasad, R., Eds.; Springer: Singapore, 2021; pp. 65–89, ISBN 9789811566066.
62. Zhang, Y.; Zhu, Y.-G.; Houot, S.; Qiao, M.; Nunan, N.; Garnier, P. Remediation of Polycyclic Aromatic Hydrocarbon (PAH) Contaminated Soil through Composting with Fresh Organic Wastes. *Environ. Sci. Pollut. Res.* **2011**, *18*, 1574–1584. [CrossRef] [PubMed]
63. Lu, X.-Y.; Zhang, T.; Fang, H.H.-P. Bacteria-Mediated PAH Degradation in Soil and Sediment. *Appl. Microbiol. Biotechnol.* **2011**, *89*, 1357–1371. [CrossRef] [PubMed]
64. Owsianiak, M.; Szulc, A.; Chrzanowski, Ł.; Cyplik, P.; Bogacki, M.; Olejnik-Schmidt, A.K.; Heipieper, H.J. Biodegradation and Surfactant-Mediated Biodegradation of Diesel Fuel by 218 Microbial Consortia Are Not Correlated to Cell Surface Hydrophobicity. *Appl. Microbiol. Biotechnol.* **2009**, *84*, 545–553. [CrossRef] [PubMed]
65. Zeng, G.; Liu, Z.; Zhong, H.; Li, J.; Yuan, X.; Fu, H.; Ding, Y.; Wang, J.; Zhou, M. Effect of Monorhamnolipid on the Degradation of N-Hexadecane by Candida Tropicalis and the Association with Cell Surface Properties. *Appl. Microbiol. Biotechnol.* **2011**, *90*, 1155–1161. [CrossRef] [PubMed]
66. Husain, S. Effect of Surfactants on Pyrene Degradation by Pseudomonas Fluorescens 29L. *World J. Microbiol. Biotechnol.* **2008**, *24*, 2411–2419. [CrossRef]

67. Ławniczak, Ł.; Marecik, R.; Chrzanowski, Ł. Contributions of Biosurfactants to Natural or Induced Bioremediation. *Appl. Microbiol. Biotechnol.* **2013**, *97*, 2327–2339. [CrossRef] [PubMed]
68. Lai, C.-C.; Huang, Y.-C.; Wei, Y.-H.; Chang, J.-S. Biosurfactant-Enhanced Removal of Total Petroleum Hydrocarbons from Contaminated Soil. *J. Hazard. Mater.* **2009**, *167*, 609–614. [CrossRef] [PubMed]
69. Mendes Da Silva Santos, E.; Alvares Da Silva Lira, I.R.; Moraes Meira, H.; Dos Santos Aguiar, J.; Diniz Rufino, R.; Germano De Almeida, D.; Casazza, A.A.; Converti, A.; Asfora Sarubbo, L.; Moura De Luna, J. Enhanced Oil Removal by a Non-Toxic Biosurfactant Formulation. *Energies* **2021**, *14*, 467. [CrossRef]
70. Das, N.; Chandran, P. Microbial Degradation of Petroleum Hydrocarbon Contaminants: An Overview. *Biotechnol. Res. Int.* **2011**, *2011*, 1–13. [CrossRef] [PubMed]
71. Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J.; Skalski, T. Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil. *Materials* **2022**, *15*, 400. [CrossRef] [PubMed]
72. Steliga, T.; Jakubowicz, P.; Kapusta, P. Changes in Toxicity during in Situ Bioremediation of Weathered Drill Wastes Contaminated with Petroleum Hydrocarbons. *Bioresour. Technol.* **2012**, *125*, 1–10. [CrossRef] [PubMed]
73. Wu, M.; Li, W.; Dick, W.A.; Ye, X.; Chen, K.; Kost, D.; Chen, L. Bioremediation of Hydrocarbon Degradation in a Petroleum-Contaminated Soil and Microbial Population and Activity Determination. *Chemosphere* **2017**, *169*, 124–130. [CrossRef] [PubMed]
74. Teliga, T.; Kluk, D. *Assessment of the Composition of Pollution of Soil Contaminated with TPH and PAHs for the Development of the Bioremediation Technology*; Instytut Nafty i Gazu-Państwowy Instytut Badawczy: Kraków, Poland, 2017; ISBN 9788365649201.
75. Kanaly, R.A.; Harayama, S. Biodegradation of High-Molecular-Weight Polycyclic Aromatic Hydrocarbons by Bacteria. *J. Bacteriol.* **2000**, *182*, 2059–2067. [CrossRef] [PubMed]
76. Wilson, S.C.; Jones, K.C. Bioremediation of Soil Contaminated with Polynuclear Aromatic Hydrocarbons (PAHs): A Review. *Environ. Pollut.* **1993**, *81*, 229–249. [CrossRef] [PubMed]
77. Van Hamme, J.D.; Singh, A.; Ward, O.P. Recent Advances in Petroleum Microbiology. *Microbiol. Mol. Biol. Rev.* **2003**, *67*, 503–549. [CrossRef] [PubMed]
78. Abbasian, F.; Lockington, R.; Megharaj, M.; Naidu, R. A Review on the Genetics of Aliphatic and Aromatic Hydrocarbon Degradation. *Appl. Biochem. Biotechnol.* **2016**, *178*, 224–250. [CrossRef] [PubMed]
79. Cai, D.; Wang, Y.; Zhao, X.; Zhang, C.; Dang, Q.; Xi, B. Regulating the Biodegradation of Petroleum Hydrocarbons with Different Carbon Chain Structures by Composting Systems. *Sci. Total Environ.* **2023**, *903*, 166552. [CrossRef] [PubMed]
80. Ghosal, D.; Ghosh, S.; Dutta, T.K.; Ahn, Y. Current State of Knowledge in Microbial Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *Front. Microbiol.* **2016**, *7*, 1369. [CrossRef] [PubMed]
81. Wang, Z.; Fingas, M.; Blenkinsopp, S.; Sergy, G.; Landriault, M.; Sigouin, L.; Foght, J.; Semple, K.; Westlake, D.W.S. Comparison of Oil Composition Changes Due to Biodegradation and Physical Weathering in Different Oils. *J. Chromatogr. A* **1998**, *809*, 89–107. [CrossRef] [PubMed]
82. Venosa, A.D.; Zhu, X. Biodegradation of Crude Oil Contaminating Marine Shorelines and Freshwater Wetlands. *Spill Sci. Technol. Bull.* **2003**, *8*, 163–178. [CrossRef]
83. Steliga, T.; Wojtowicz, K.; Kapusta, P.; Brzeszcz, J. Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture. *Molecules* **2020**, *25*, 709. [CrossRef] [PubMed]
84. Steliga, T.; Jakubowicz, P.; Wojtowicz, K.; Kluk, D. Zastosowanie Testów Toksykologicznych w Przemyśle Naftowym. *Nafta-Gaz* **2018**, *74*, 684–689. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

sustainability

MDPI

Suplementary materials

Influence of biosurfactants on the efficiency of petroleum hydrocarbons biodegradation in soil

Katarzyna Wojtowicz[1], Teresa Steliga[1*], Tomasz Skalski[2] and Piotr Kapusta[3]

[1] Department of Production Technology of Reservoir Fluids, Oil and Gas Institute – National Research Institute, Lubicz 25 A, 31-503 Kraków, Poland
[2] Biotechnology Centre, Silesian University of Technology, Krzywoustego 8, 44-100 Gliwice, Poland
[3] Department of Microbiology, Oil and Gas Institute – National Research Institute, Lubicz 25 A, 31-503 Kraków, Poland
* Correspondence: teresa.steliga@inig.pl

Academic Editor: Firstname Lastname

Received: date
Revised: date
Accepted: date
Published: date

Citation: To be added by editorial staff during production.

Copyright: © 2025 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

Description of the samples presented in the Figure 1:

Samples inoculated with the bacterial consortium MC and enriched with a selected biosurfactant (1 g/dm³):

1. soil G6 + MC + γ-PGA
2. soil G6 + MC + surfactin
3. soil G6 + MC + rhamnolipids
4. soil G6 + MC + mixture PSR
5. soil G6 + MC + JBR425
6. soil G6 + MC + JBR320

Samples inoculated with the bacterial consortium MC and enriched with a selected biosurfactant (5 g/dm³):

7. soil G6 + MC + γ-PGA
8. soil G6 + MC + surfactin
9. soil G6 + MC + rhamnolipids
10. soil G6 + MC + mixture PSR
11. soil G6 + MC + JBR425
12. soil G6 + MC + JBR320

Control samples enriched only with the selected biosurfactant (1 g/dm³), without inoculation:

13. soil G6 + γ-PGA
14. soil G6 + surfactin
15. soil G6 + rhamnolipids
16. soil G6 + mixture PSR
17. soil G6 + JBR425
18. soil G6 + JBR320

Control samples enriched only with the selected biosurfactant (5 g/dm³), without inoculation:

19. soil G6 + γ-PGA
20. soil G6 + surfactin
21 soil G6 + rhamnolipids
22. soil G6 + mixture PSR
23. soil G6 + JBR425
24. soil G6 + JBR320

Reference samples:

25. blanc sample – non-inoculated, no additives
26. soil G6 + bacterial consortium MC – control sample inoculated only with the MC consortium

Methods description

Respirometric test

Respirometric measurements were used to assess the aerobic biodegradation potential by monitoring oxygen consumption in soil samples under different enrichment conditions [1]. The experiments were conducted using the OxiTop® Control system. For the measurements, 30 g of soil (sample G6) was placed in sealed respirometric bottles. Depending on the experimental setup, the bottles were supplemented with: a bacterial consortium enriched with selected biosurfactants (at concentrations of 1 or 5 g/dm³), biosurfactants alone (1 or 5 g/dm³) (control samples), the bacterial consortium alone (control sample), or sterile water (blank sample). The bottles were then tightly sealed with measurement heads and incubated in a thermostatic chamber at 20°C for 90 days. The OxiTop® Control system recorded the pressure in the closed system every two hours. The data were transferred via an IR interface to the OxiTop OC 110 controller for graphical and statistical analysis using Achat OC software. Oxygen consumption (expressed in mg O_2/dm³) was calculated based on the pressure decrease using the following equation [2-4]:

$$m_{O_2} = \frac{M(O_2)}{RT_m} \cdot (V_g + \alpha \frac{T_m}{T_0}) \cdot \Delta p$$

Where: $m(O_2)$ – mass of consumed oxygen [mg/dm³], Δp – pressure drop in the vessel [Pa], Vg – volume of free gas [m³], $M(O_2)$ – molar mass of oxygen [kg/mol], R – gas constant [J·mol^{-1}·K^{-1}], T_m – measurement temperature [K], T_0 – reference temperature (273.15 K),

α – absorption coefficient (0.03103).

At the end of the 90-day incubation period, the obtained oxygen consumption data were used to evaluate the potential synergistic effect between the bacterial consortium and the biosurfactants. For this purpose, the Synergitic Index (SI) was calculated using the following formula:

$$\text{Synergitic Index (SI)} = \text{Synergitic index (SI)} \ \frac{\text{Effect(MC+B)}}{\text{Effect(MC)}+\text{Effect(B)}}$$

Where: *Effect(MC+B)* – measured biological activity (oxygen consumption) in the system inoculated with both the bacterial consortium (MC) and biosurfactants, *Effect(MC)* – activity in the system inoculated with the bacterial consortium MC alone, *Effect(B)* – activity in the system with biosurfactants alone.

SI values greater than 1 indicate a synergistic interaction between the components. This parameter was used to identify combinations showing enhanced biodegradation potential due to cooperative action.

Chromatographic analyses

TPH Extraction and Quantification

For the analysis of TPH in soil, 10 g of air-dried soil were placed in Erlenmeyer flasks and subjected to solvent extraction using dichloromethane (POCH S.A., Poland). The extraction was performed in three successive steps (3 × 20 mL), each with 15-minute ultrasonic agitation in a Sonoswiss SW 6H ultrasonic bath (30 kHz). The use of ultrasound-assisted extraction improved recovery rates, achieving approximately 95.9%, as verified using o-tert-phenyl as an internal recovery standard [4-5].

Following extraction, polar co-extracts were removed via solid-phase extraction (SPE) using Bakerbond Florisil® columns (No. JTB- 7213-03). The eluates were concentrated under reduced pressure using a rotary evaporator, redissolved in 1 mL of dichloromethane, and subjected to chromatographic analysis.

Quantitative and qualitative analysis TPH, individual n-alkanes (n-C6–n-C44) and isoprenoid hydrocarbons (pristane and phytane) was carried out using a Perkin Elmer Clarus 500 gas chromatograph equipped with a flame ionization detector (FID). The separation was performed on a fused silica RTX-1 capillary column (30 m × 0.53 mm; Restek, USA), with helium as the carrier gas at a constant flow rate of 20 mL·min^{-1}. The oven temperature program was as follows: initial isothermal hold at 30 °C for 2 min; ramp to 105 °C at 10 °C·min^{-1}; increase to 285 °C at 5 °C·min^{-1}; isothermal hold at 285 °C for 5 min. [4-7]. The injector and detector temperatures were set at 290 °C and 320 °C, respectively. Calibration for TPH quantification was based on a certified standard (BAM K010; Tusnovic Instruments). Quantification of individual hydrocarbons was performed using certified standard mixtures: Supelco D2807 (n-C6–n-C44) and A029668 (Fuel Oil Degradation Mix: n-C17, pristane, n-C18, phytane). The method showed good repeatability, with standard deviations ranging between 1.53 and 4.32%. Based on the concentrations of TPH and individual n-alkanes determined in the soil samples, the biodegradation rate of these compounds was calculated using the following equation:

$$\textbf{Biodegradation [\%]} = \frac{Cn{-}x_0 - Cn{-}x}{Cn{-}x_0} * \mathbf{100}$$

Where: $Cn\text{-}x_0$ - initial concentration of TPH or n-alkane in the soil before bioremediation [mg/kg DM]. C_n - concentration of TPH or n-alkane in the soil after the bioremediation [mg/kg DM].

In addition, biodegradation coefficients C17/Pr and C18/Ph were calculated as indicators of biodegradation, using the following formulas:

$$\text{n-C17/Pr}=\frac{c(n-C17)}{c(Pr)}$$

Where: c(n-C17) – concentration of n-c17 in soil [mg/kg DM], c(Pr) – concentration of pristane in soil [mg/kg DM],

$$\text{n-C18/Ph}=\frac{c(n-C18)}{c(Ph)}$$

Where: c(n-C18) – concentration of n-c18 in soil [mg/kg DM], c(Ph) – concentration of phytane in soil [mg/kg DM],

PAHs Extraction and Quantification

The quantification of polycyclic aromatic hydrocarbons (PAHs) in soil samples was carried out using a modified QuEChERS extraction procedure, as described in the study by Wojtowicz (2022) [8], which also provides detailed information on method validation, analyte recovery, and statistical treatment of the data.

For each analysis, 10 g of air-dried soil were transferred into 50 mL polypropylene centrifuge vials. Subsequently, 5 mL of distilled water and 10 mL of acetonitrile

(Chempur, Poland) were added, followed by a salt mixture containing magnesium sulfate ($MgSO_4$) and sodium chloride (NaCl) (SST640, Interchim, France). The vials were sealed and manually shaken for 5 minutes to ensure efficient extraction. The samples were then centrifuged at 3500 rpm for 10 minutes using an MPW-223e centrifuge (MPW, Poland).
The organic supernatant was purified using a Clean-up Kit (No. JO3937, Interchim, France), followed by filtration through 0.2 µm nylon syringe filters [6-7].
Chromatographic analysis of PAHs was conducted on a Vanquish Core HPLC system (Thermo Scientific, USA) equipped with a UV-VIS detector and a fluorescence detector (FLD), operated in series and supported by an autosampler. Separation was achieved on a NUCLEODUR C18 PAH column (125 × 4 mm, 3 µm; Macherey-Nagel, Germany). The injection volume was 10 µL. The mobile phases used were methanol (eluent A, 70% purity, POCH S.A., Poland) and acetonitrile (eluent B, 99% purity, POCH S.A., Poland). The analysis was conducted with a flow rate of 1.5 mL/min, using the following gradient program: 20–50% B (1.5 min), 50–100% B (1 min), hold at 100% B (1 min), decrease to 0% B (3 min), and final hold at 100% A (3 min) [6-8].
For the quantitative determination of PAHs, a certified standard solution (PAH Standard, Ref. No. 722393, Macherey-Nagel, Germany) was used. Analyses were performed on contaminated soil samples spiked with a certified reference material (PAHs by HPLC 40g, SQC017-40G, Sigma Aldrich, USA), ensuring high accuracy and comparability of results across matrices. Based on the concentrations of total and individual PAHs determined in the soil samples, the biodegradation rate of these compounds was calculated using the following equation:

$$\textbf{Biodegradation [\%]} = \frac{Cn{-}x_0 - Cn{-}x}{Cn{-}x_0} * \mathbf{100}$$

Where: $Cn\text{-}x_0$ - initial concentration of given PAH or total PAHs in the soil before bioremediation [mg/kg DM]. C_n - concentration of given PAH or total PAHs in the soil after the bioremediation [mg/kg DM].

toxicological analyses

Phytotoxkit™ Test (MicroBioTests Inc., Belgium)
The Phytotoxkit™ chronic toxicity assay evaluates the inhibition of seed germination and early root elongation. Three plant species, selected based on their germination and root growth rates, are used to ensure comprehensive toxicity assessment within a 72-hour incubation period. These species include the monocotyledon *Sorghum saccharatum* and the dicotyledons *Lepidium sativum* (cress) and *Sinapis alba* (white mustard). Tests were conducted in triplicate using transparent polystyrene plates under controlled conditions (25°C, darkness) [6-7, 9-12]. Inhibition of germination and root growth was calculated as:

$$I = \frac{(A - B)}{B} \cdot 100\%$$

Where: **I** – inhibition (%), **A** – seed germination or root length in control soil, **B** – seed germination or root length in test soil

Ostracodtoxkit(F)™ Test (MicroBioTests Inc., Belgium)
The Ostracodtoxkit(F)™ test is a direct contact chronic toxicity bioassay using juvenile *Heterocypris incongruens* crustaceans. Juveniles hatched from cysts over 52 hours were exposed to soil samples in six-well polystyrene microplates. Each well contained 0.5 g of soil and 2 mL of standardized algal suspension as food. Ten ostracods were introduced per well. Controls were conducted using reference soil. Incubation occurred at 25°C for

six days [6-7, 12-14]. Endpoints included survival and growth, with growth inhibition (I) calculated as:

$$I = 100 - \left(\frac{A}{B} \cdot 100\%\right)$$

Where: A- ostracod length in test soil, B=- ostracod length in control soil.

Microtox® Solid Phase Test (SDI, USA)

The Microtox® test employs the luminescent bacterium *Vibrio fischeri* to evaluate sample toxicity through bioluminescence inhibition. Bioluminescence is catalyzed by luciferase during bacterial metabolism and decreases in the presence of toxic substances. Lyophilized *V. fischeri* bacteria were stored at -20°C and rehydrated before testing. The assay involved exposing bacteria to serial dilutions of wastewater samples, with luminescence measured using the Delta Tox analyzer. The effective concentration reducing luminescence by 50% (EC_{50}) was determined; lower EC_{50} values indicate higher toxicity [6-7, 15-17]. To facilitate interpretation, EC_{50} values were converted to Toxicity Units (TU) using:

$$TU = \frac{100}{EC_{50}}$$

Higher TU values correspond to increased toxicity.

To allow for comparison across different bioassays, toxicity results expressed as inhibition percentages (Phytotoxkit™, Ostracodtoxkit(F)™) and effective concentrations (EC_{50}; Microtox®) were converted to Toxicity Units (TU). For tests reporting EC_{50} values, TU was calculated as:

$$TU = \frac{100}{EC_{50}}$$

For tests where inhibition (%) was directly measured, TU was derived from inhibition values (I, in %) as:

$$TU=I$$

where I represents the percentage inhibition of seed germination, root growth, or organism growth. This approach enables the integration of toxicity endpoints on a common scale, facilitating comparison between assays.

The key information regarding the applied bioassays is summarized in Table S1.

Table S1 Characteristics of bioindication ecotoxicological tests applied for toxicity assessment

Test	Test Organism	Type of Toxicity	Experimental Parameters	Toxicity Indicators
Phytotoxkit™	*Sorghum saccharatum*, *Lepidium sativum*, *Sinapis alba*	Acute	- Incubation time: 72 h - Temperature: 25°C - Darkness, polystyrene test plates	EC_{50} for seed germination and root elongation Converted to Toxic Units (TU)

Test	Test Organism	Type of Toxicity	Experimental Parameters	Toxicity Indicators
Ostracodtoxkit F™	*Heterocypris incongruens* (freshwater crustacean)	Chronic	- Exposure duration: 6 days - Temperature: 25°C - Carried out in 6-well polystyrene plates	Mortality (%) Growth inhibition (based on body length) Converted to Toxic Units (TU)
Microtox® Solid Phase	*Vibrio fischeri* (marine bacterium)	Acute	- Exposure time: 15 min - Temperature: ambient - Based on luciferase-catalyzed bioluminescence inhibition	EC_{50} (50% luminescence reduction) Converted to Toxic Units (TU)

Table 2 Synergitic Index (SI) for different experimental setups and biosurfactant concentrations

Sample	**Synergitic Index** (SI) Concentration of biosurfactant in the inoculating mixture (1 g/dm³)	**Synergitic Index** (SI) Concentration of biosurfactant in the inoculating mixture (5 g/dm³)
soil G6+MC+γ-PGA	1,152	1,365
soil G6+MC+surfactin	1,112	1,071
soil G6+MC+rhamnolipids	1,146	1,309
soil G6+MC+mixture PSR	1,218	1,398
soil G6+MC+JBR425	1,204	1,461
soil G6+MC+JBR320	1,195	1,455

*** All SI values are greater than 1, which indicates the presence of a synergitic effect between the bacterial consortium and the applied biosurfactants.**

References:

1. Malińska, K. Application of a Modified OxiTop® Respirometer for Laboratory Composting Studies. *Archives of Environmental Protection* **2016**, *42*, 56–62, doi:10.1515/aep-2016-0007.
2. Steliga, T.; Wojtowicz, K. Wykorzystanie Testów Respirometrycznych Do Oceny Efektywności Biodegradacji Osadów z Instalacji Kopalnianych. *Nafta-Gaz* **2019**, *75*, 29–37, doi:10.18668/NG.2019.01.05.
3. Steliga, T.; Uliasz, M. Spent Drilling Muds Management and Natural Environment Protection. *Gospodarka Surowcami Mineralnymi* **2014**, *30*, 135–155, doi:10.2478/gospo-2014-0011.
4. Steliga, T.; Wojtowicz, K.; Kapusta, P.; Brzeszcz, J. Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture. *Molecules* **2020**, *25*, 709, doi:10.3390/molecules25030709.
5. Steliga, T.; Jakubowicz, P.; Kapusta, P. Changes in Toxicity during in Situ Bioremediation of Weathered Drill Wastes Contaminated with Petroleum Hydrocarbons. *Bioresource Technology* **2012**, *125*, 1–10, doi:10.1016/j.biortech.2012.08.092.
6. Wojtowicz, K.; Steliga, T.; Kapusta, P. Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using Echinacea Purpurea. *Applied Sciences* **2023**, *13*, 13077, doi:10.3390/app132413077.
7. Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J. Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (Zea Mays). *Molecules* **2023**, *28*, 6104, doi:10.3390/molecules28166104.

8. Wojtowicz, K. Opracowanie Metodyki Oznaczania WWA w Próbkach Gleb z Wykorzystaniem Chromatografii Cieczowej HPLC. *Nafta-Gaz* **2022**, *78*, 141–153, doi:10.18668/NG.2022.02.06.
9. Mamindy-Pajany, Y., Hamer, B., Roméo, M., Géret, F., Galgani, F., Durmiši, E., Hurel, C., & Marmier, N. (2011). The toxicity of composted sediments from Mediterranean ports evaluated by several bioassays. *Chemosphere, 82*, 362–369. https://doi.org/10.1016/j.chemosphere.2010.10.005
10. Blinova, I., Niskanen, J., Kajankari, P., Kanarbik, L., Käkinen, A., Tenhu, H., Penttinen, O. P., & Kahru, A. (2013). Toxicity of two types of silver nanoparticles to aquatic crustaceans Daphnia magna and Thamnocephalus platyurus. *Environmental Science and Pollution Research, 20*(5), 3456–3463.
11. Baran, A., & Tarnawski, M. (2013). Phytotoxkit/Phytotestkit and Microtox® as tools for toxicity assessment of sediments. *Ecotoxicology and Environmental Safety, 98*, 19–27. https://doi.org/10.1016/j.ecoenv.2013.10.010Baran, A., & Tarnawski, M. (2013). Phytotoxkit/Phytotestkit and Microtox® as tools for toxicity assessment of sediments. *Ecotoxicology and Environmental Safety, 98*, 19–27. https://doi.org/10.1016/j.ecoenv.2013.10.010
12. Steliga, T., Jakubowicz, P., Wojtowicz, K., Kluk, D., 2018a. Zastosowanie testów toksykologicznych w przemyśle naftowym. Nafta-Gaz 74, 684–689. https://doi.org/10.18668/NG.2018.09.07
13. Rouvalis, A., Theodoropoulos, C., & Georgudaki, J. I. (2013). Assessment of toxicity of the untreated and Pleurotus ostreatus treated olive mill wastewater by using microbiotests. *International Journal of Environmental Engineering, 5*, 373. https://doi.org/10.1504/IJEE.2013.056623
14. . Niyommaneerat, W., Nakajima, F., Tobino, T., & Yamamoto, K. (2017). Development of a chronic sediment toxicity test using the benthic ostracod Heterocypris incongruens and their application to toxicity assessments of urban road dust. *Ecotoxicology and Environmental Safety, 143*, 266–274. https://doi.org/10.1016/j.ecoenv.2017.05.011
15. Lima, T.M.S., Procópio, L.C., Brandão, F.D., Leão, B.A., Tótola, M.R., Borges, A.C., 2011. Evaluation of bacterial surfactant toxicity towards petroleum degrading microorganisms. Bioresource Technology 102, 2957–2964. https://doi.org/10.1016/j.biortech.2010.09.109
16. Foucault, Y., Durand, M.-J., Tack, K., Schreck, E., Geret, F., Leveque, T., Pradere, P., Goix, S., & Dumat, C. (2013). Use of ecotoxicity test and ecoscores to improve the management of polluted soils: Case of a secondary lead smelter plant. *Journal of Hazardous Materials, 246–247*, 291–299. https://doi.org/10.1016/j.jhazmat.2012.12.042
17. Oleszczuk, P., Jośko, I., Kuśmierz, M., Futa, B., Wielgosz, E., Ligęza, S., & Pranagal, J. (2014). Microbiological, biochemical and ecotoxicological evaluation of soils in the area of biochar production in relation to polycyclic aromatic hydrocarbon content. *Geoderma, 213*, 502–511. https://doi.org/10.1016/j.geoderma.2013.08.027

molecules

MDPI

Article

Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (*Zea mays*)

Katarzyna Wojtowicz *, Teresa Steliga, Piotr Kapusta and Joanna Brzeszcz

Oil and Gas Institute—National Research Institute, ul. Lubicz 25 A, 31-503 Krakow, Poland; teresa.steliga@inig.pl (T.S.); piotr.kapusta@inig.pl (P.K.); joanna.brzeszcz@inig.pl (J.B.)
* Correspondence: wojtowiczk@inig.pl

Abstract: Biological methods are currently the most commonly used methods for removing hazardous substances from land. This research work focuses on the remediation of oil-contaminated land. The biodegradation of aliphatic hydrocarbons and PAHs as a result of inoculation with biopreparations B1 and B2 was investigated. Biopreparation B1 was developed on the basis of autochthonous bacteria, consisting of strains *Dietzia* sp. IN118, *Gordonia* sp. IN101, *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN119, *Rhodococcus globerulus* IN113 and *Raoultella* sp. IN109, whereas biopreparation B2 was enriched with fungi, such as *Aspergillus sydowii*, *Aspergillus versicolor*, *Candida* sp., *Cladosporium halotolerans*, *Penicillium chrysogenum*. As a result of biodegradation tests conducted under ex situ conditions for soil inoculated with biopreparation B1, the concentrations of TPH and PAH were reduced by 31.85% and 27.41%, respectively. Soil inoculation with biopreparation B2 turned out to be more effective, as a result of which the concentration of TPH was reduced by 41.67% and PAH by 34.73%. Another issue was the phytoremediation of the pre-treated G6-3B2 soil with the use of *Zea mays*. The tests were carried out in three systems (system 1—soil G6-3B2 + *Zea mays*; system 2—soil G6-3B2 + biopreparation B2 + *Zea mays*; system 3—soil G6-3B2 + biopreparation B2 with γ-PGA + *Zea mays*) for 6 months. The highest degree of TPH and PAH reduction was obtained in system 3, amounting to 65.35% and 60.80%, respectively. The lowest phytoremediation efficiency was recorded in the non-inoculated system 1, where the concentration of TPH was reduced by 22.80% and PAH by 18.48%. Toxicological tests carried out using Phytotoxkit™, Ostracodtoxkit™ and Microtox® Solid Phase tests confirmed the effectiveness of remediation procedures and showed a correlation between the concentration of petroleum hydrocarbons in the soil and its toxicity. The results obtained during the research indicate the great potential of bioremediation practices with the use of microbial biopreparations and *Zea mays* in the treatment of soils contaminated with petroleum hydrocarbons.

Keywords: biodegradation; biopreparation; ecotoxicology; microorganisms; petroleum contaminants; phytoremediation; γ-PGA-polyglutamic acid

check for updates

Citation: Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J. Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (*Zea mays*). *Molecules* **2023**, *28*, 6104. https://doi.org/10.3390/molecules28166104

Academic Editor: Fahrul Huyop

Received: 6 July 2023
Revised: 14 August 2023
Accepted: 15 August 2023
Published: 17 August 2023

Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

Human activity has been altering the environment for ages, reducing its resources, changing its natural properties and introducing foreign substances not present in the natural conditions. Oil and gas mining uses up valuable minerals, involving large amounts of various materials and energy and producing a significant amount of waste and causing irreversible changes in the natural environment. Petroleum products are one of the main sources of soil pollution appearing due to oil mining work. Petroleum waste has an adverse effect on plant production and is dangerous to people and animals. The introduction of petroleum waste into the ecosystems disturbs the natural cycles of matter and energy, reducing the water storage capacity of the soil and hindering air exchange between the soil and the atmosphere. Additionally, the large number of different specific properties of petroleum

pollutants and the products of their biological and physico-chemical transformations in the natural environment make it very difficult to identify the environmental consequences of environmental pollution and select the correct remediation technology and the methods for determining and monitoring the progress of the remediation of contaminated soil [1]. At the moment, the most popular methods for the remediation of petroleum-contaminated soil include biological methods, which have a wide range of applications and are highly efficient and fairly inexpensive. However, biological methods have their limitations, including the fact that high concentrations of pollutants in the soil may exceed the levels tolerated by the organisms used in the process. For this reason, in some cases, it is necessary to conduct staged remediation of soil using multiple remediation methods.

Biodegradation using biopreparations made with microorganisms isolated from areas naturally occurring in the remediated region is one of the most effective and frequently used methods for remediating petroleum-contaminated soil. It is estimated that out of the native microbiological population, approx. 10^5 to 10^6 have properties, which enable the biodegradation of petroleum hydrocarbons [2]. However, it should be noted that most bacteria, fungi and yeasts metabolise petroleum products to a limited extent. *Desulfosarcina* sp. [3], *Mycobacterium* sp., *Pseudomonas butanovora* [4], *Pseudonocardia* sp. [5] bacteria are effective at breaking down C_2–C_8 alkanes, whereas *Acinetobacter baumannii* [6], *Alcanivorax borkumensis* [7], *Collimonas* sp., *Dietzia* sp. [8], *Flavobacterium* sp., *Pseudomonas* sp. [9], *Geobacillus thermodenitrificant* [10], *Gordonia sihwensis* [11], *Mycobacterium* sp. [12], *Mycobacterium frederiksbergense* IN53, *Acinetobacter* sp. IN47 [13,14], *Nocardioides albus*, *Ochrobactrum* sp., *Oleispira antarctica* [15], *Pseudomonas aeruginosa* [16–20], *Pseudomonas* spp. *Alphaproteobacteria* and *Betaproteobacteria* [21], *Pseudomonas citronellolis* [22], *Pseudomonas stutzeri* [15,23], *Rhodococcus* sp. [24], *Rhodococcus coprophilus* [25], *Rhodococcus ruber* [26] or *Stenotrophomonas maltophilia* are used in biodegradation tests of alkanes with a carbon chain length of C_{10}–C_{36}, oils and asphaltenes. In addition, it was confirmed that the fungal strains *Aspergillus fumigatus*, *Aspergillus, terreus*, *Fusarium solani*, *Gliomastix* sp., *Paecilomyces* sp. and *Verticillum* sp. have the ability to degrade short-chain n-alkanes [27,28]. Hydrocarbons with a carbon chain length of C_{10}–C_{36} are also broken down by fungi *Penicillium chrysogenum* [29], *Penicillium Citrinum* [27], *Candida* sp. [30], *Candida lipolytica* [28], *Candida parapsilosis*, *Candida krusei*, *Candida famata* and *Rhodotorula* spp. [31], *Aspergillus flavus*, *Fusarium* sp., *Scopulariopsis* sp. [27].

The most popular microorganisms capable of breaking down polycyclic aromatic hydrocarbons include the *Achromobacter xylosoxidans* [32], *Aeribacillus pallidus* [33], *Alcaligenes faecalis*, *Agmenellum quadruplicatum* [34], *Bacillus Licheniformis*, *Bacillus Mojavensis* [35], *Bacillus thuringiensis*, *Brevundimonas* sp. [36], *Burkholderia* sp., *Cycloclasticus* sp. [37], *Gordona sputi* [38,39], *Micrococcus* sp. [40], *Mycobacterium* sp. [41,42], *Mycobacterium vanbaalenii* [43–45], *Neptunomonas naphthovoran* [46], *Nocardioides* sp., *Nocardia*, *Oscillatoria* [47], *Pleurotus cornucopiae* [48], *Pseudomonas* sp. [49,50], *Pseudomonas aeruginosa* [50], *Pseudomonas putida* [51], *Pseudomonas stutzeri* [52], *Rhodococcus* sp. [53], *Rhodococcus erythropolis* [54–56], *Sphingobium* and *Novosphingobium*, *Sphingomonas* sp. [57,58], *Sphingomonas yanoikuyae* [59], *Staphylococcus warneri* and *Bacillus pumilus* [60], *Stenotrophomonas* sp. [61], *Stenotrophomonas maltophilia* [62], *Streptomyces* sp. [63] bacteria and fungi *Aspergillus versicolor* [64], *Aspergillus sydowii* [65], *Candida* sp. [30], *Clodosporium* sp. [66], *Penicillium* sp., *Fusarium solani* [67,68]. For this reason, it is preferred to use biopreparations containing several to several dozen strains of bacteria and fungi to ensure the broadest possible spectrum of action. The microorganisms contained in the biopreparations should be specially selected from the natural environment, which significantly improves the efficiency of pollutant removal [69,70] and helps avoid antagonistic reactions of the native micro-flora of the soil to the foreign microbial cultures. It is also necessary to bear in mind that the microbes introduced into the soil cannot be pathogenic to plants and animals, and they should not produce toxic substances, should not be resistant to antibiotics and should not be involved in the frequent exchange of genes, which encode undesirable properties [71–74]. At the moment, there are more and more studies combining several methods of soil bioremediation to ensure the highest possible efficiency. Interesting solutions, which have been successfully applied, include the com-

bination of biodegradation of petroleum products by inoculation with a biopreparation with phytoremediation.

Biological methods using plants (phytoremediation) mainly rely on the tolerance for high concentrations of toxic substances, absorption, accumulation and metabolism of hazardous substances in large quantities (in the plant's own organs) and stabilisation of pollutants in the soil or transformation of toxic substances in the environment into non-toxic compounds [75]. Organic pollutants are absorbed by the plants together with water by simple diffusion. The diffusion of petroleum products into the plant, however, is hindered if most ingredients of crude oil are hydrophobic compounds, which are highly lipophilic and have large molecular masses. For this reason, the absorption of lipophilic substances by plants is positively correlated with the lipid content of the root epidermis. However, plants may release surface-active agents, such as low-molecular-weight organic acids or polypeptides, which reduce surface tension and emulsify hydrophobic substances, increasing bioavailability [76–83]. The plants used for phytoremediation should be tolerant of high concentrations of xenobiotics, a high degree of accumulation or biodegradation of pollutants, even with a fairly low pollution level, the ability to accumulate multiple pollutants at the same time, quick growth, high biomass production and resistance to diseases and pests and difficult environmental conditions [75,84]. The most often cited species of haptophytes include the tall sedge (*Carex hirta*), Kentucky bluegrass (*Poa pratensis*), perennial ryegrass (*Lolium perenne*), broadleaf plantain (*Plantago major*) and common nettle (*Urtica dioica*), which tolerate petroleum pollutants in the soil in the range of 204–504 g/kg, 112–320 g/kg, 132–166 g/kg, up to 470 g/kg and up to 506 g/kg, respectively [85]. Other haptophytes include the common toadflax (*Linaria vulgaris*) [86], European dewberry (*Rubus caesius*), tufted grass (*Holcus lanatus*) [87], white clover (*Trifolium repens*) [88], sweet yellow clover (*Melilotus officinalis*) [89] and tall fescue (*Festuca arundinacea*) [90]. Further plant species, which can successfully be used in the phytoremediation of petroleum compounds, include the marvel of Peru (*Mirabilis jalapa*) [91], alfalfa (*Medicago sativa*) [92], perennial ryegrass (*Lolium perenne, Lolium multiflorum*) [93,94], common bird's-foot trefoil (*Lotus corniculatus*) [94,95], sorghum (*Sorghum bicolor*) [96], maize (*Zea mays*) [97–99], Bermuda grass (*Cynodon dactylon*) [100], nodding beggarticks (*Bidens cernua*) [101], common blanketflower (*Gaillardia aristata*) [102], eastern purple coneflower (*Echinacea purpurea*) [102,103] and grasses and leguminous plants [104–106]. Since the plants used for phytoremediation should grow fast, have a deep and extensive root system and fairly high biomass, trees such as poplars (*Populus* L.) and birches (*Salix* L.) [107,108] are increasingly often considered as plants, which can be used for the bioremediation of petroleum-contaminated soil. Additionally, the effectiveness of phytoremediation largely depends on the soil conditions, which affect the growth of plant organisms. For this reason, recent years saw increased research on the ways of improving the efficiency of phytoremediation by combining it with biodegradation using biopreparations made with native microorganisms and biopreparation additives. One of the compounds, which can increase the efficiency of phytoremediation of petroleum-contaminated soil is γ-PGA.

γ-PGA, i.e., polyglutamic acid, is a chemical compound produced by bacteria of the *Bacillus* genus, including in particular *Bacillus amyloliquefaciens, Bacillus licheniformis, Bacillus megaterium, Bacillus subtilis* and *Bacillus anthracis* [109]. γ-PGA is highly soluble in water, non-toxic to living organisms and readily biodegradable, and it can be used as a natural additive to fertilisers [110–113]. Additionally, γ-PGA is a readily available source of carbon, which significantly affects the growth of plant organisms. The satisfactory effects of the addition of γ-PGA to biopreparations made with native bacteria on the efficiency of biodegradation of petroleum hydrocarbons in the soil in areas of drilling waste pits [55,114]; the impact of γ-PGA on the removal of oil from marine sediments [115] and trichloroethane from the aquifer [116]; and the use of γ-PGA for the remediation of soil contaminated with heavy metals [117–119] were the reasons for the decision to experimentally test the possibilities of using this compound in the phytoremediation process.

2. Results

2.1. Biodegradation of Hydrocarbons Using Biopreparations

Biodegradation tests of petroleum pollutants using the ex situ prism method demonstrated that the addition of fungi to the biopreparation increased the efficiency of biodegradation of both aliphatic hydrocarbons and aromatic hydrocarbons. After three months of tests, the TPH content was reduced from 4527.70 mg/kg ds to 3090.60 mg/kg ds (in soil inoculated with biopreparation B1) and to 2640.83 mg/kg ds (in soil inoculated with biopreparation B2).

Chromatographic analyses showed that both in the case of soil inoculated with the biopreparation made with bacteria and the biopreparation made with bacteria, yeasts and fungi, aliphatic hydrocarbons with a carbon chain length of n-C_{10}–n-C_{21} were the most readily biodegradable. Their degrees of biodegradation in soil inoculated with biopreparation B1 ranged from 18.49% to 31.00% (after 1 month), from 26.93% to 50.37% (after 2 months) and from 34.78% to 64.58% (after 3 months), and in soil inoculated with biopreparation B2, the degrees of biodegradation of n-C_{10}–n-C_{21} hydrocarbons were 26.83% to 43.86% (after 1 month), 42.63% to 69.69% (after 2 months) and 54.08% to 78.00% (after 3 months), respectively. A slightly smaller biodegradation efficiency was recorded for light hydrocarbons with a carbon chain length of n-C_6–n-C_9, which ranged from 53.96% to 63.99% in G6-3B1 soil and from 78.50% to 65.17% in G6-3B2 soil after the tests. Significant differences in the biodegradation of aliphatic hydrocarbons in soil inoculated with biopreparations B1 and B2 were recorded for hydrocarbons with a carbon chain length of n-C_{22}–n-C_{30} and n-C_{31}–n-C_{36}. Their biodegradation in soil inoculated with the biopreparation made with bacteria was much slower than in soil inoculated with the biopreparation made with bacteria, yeasts and fungi. The degree of biodegradation of hydrocarbons with a carbon chain length of n-C_{22}–n-C_{30} ranged from 4.76% to 16.93% (soil G6-1B1), 7.24% to 25.76% (soil G6-2B1), 10.25% to 32.44% (soil G6-3B1), 12.28% to 23.34% (soil G6-1B2), 19.52% to 37.08% (soil G6-2B2) and 22.05% to 47.63% (soil G6-3B2). Heavy hydrocarbons containing more than 30 carbon atoms per molecule were broken down the most. Inoculation with biopreparation B1 decreased their content in the soil by 2.87% to 4.41% (after 1 month), 5.54% to 7.14% (after 2 months) and 9.23% to 10.67% (after 3 months). The biodegradation of heavy hydrocarbons was much more efficient in soil inoculated with biopreparation B2. The content of hydrocarbons in the n-C_{31}–n-C_{36} range was reduced by 7.79% to 9.65% after 1 month, by 12.38% to 15.33% after 2 months and by 14.77% to 18.65% after 3 months. The content of unidentified hydrocarbons after 3 months of biodegradation tests using the prism method decreased by 31.44% (soil G6-3B1) and 37.97% (soil G6-3B2). The specific progress of the biodegradation of aliphatic hydrocarbons in soil G6 inoculated with biopreparations B1 and B2 is shown in Figures 1 and 2.

Another important parameter, which can be used to determine the efficiency of biodegradation of petroleum pollutants in soil, is the change in n-C_{17}/Pr and n-C_{18}/F ratios. After the test, the n-C_{17}/Pr ratio was lower in soil inoculated with biopreparation B2 than in soil inoculated with biopreparation B1 by 0.29. Additionally, the n-C_{17}/Pr ratio in G6-3B2 soil was lower by 0.30 in comparison with G6-3B1 soil. The n-C_{17}/Pr and n-C_{18}/F biodegradation ratios are indicative of a higher efficiency of biodegradation of petroleum hydrocarbons in soil inoculated with biopreparation B2. The values of n-C_{17}/Pr and n-C_{18}/F biodegradation ratios are shown in Figure 3.

The PAH biodegradation tests in soil inoculated with biopreparations B1 and B2 showed that the presence of fungi in the biopreparation increased the efficiency of PAH biodegradation. This is confirmed by the results of the chromatographic analyses, which showed that the concentration of naphthalene in soil inoculated with biopreparation B1 in subsequent months of the tests decreased by 18.54% (after 1 month), 26.54% (after 2 months) and 34.49% (after 3 months), and in soil inoculated with biopreparation B2 by 25.96% (after 1 month), 36.11% (after 2 months) and 43.81% (after 3 months). Thus, the difference in the degrees of biodegradation of naphthalene in soil inoculated with biopreparations B1 and B2 after the tests amounted to 9.32%, which is a significant level, considering the short

duration of the tests. Additionally, the chromatographic analyses showed that the degree of biodegradation decreased along with the increase in the number of rings in the PAH molecule. The 3-ring PAHs (acenaphthene, fluorene, phenanthrene, anthracene) turned out to be slightly less biodegradable than naphthalene, and their degrees of biodegradation ranged from 10.02% to 13.86% (soil G6-1B1), from 17.73% to 23.44% (soil G6-2B1), from 24.40% to 30.82% (soil G6-3B1), from 12.08% to 16.70% (soil G6-1B2), from 22.44% to 29.67% (soil G6-2B2) and from 30.12% to 38.05% (soil G6-3B2). The degrees of reduction in 4-ring PAHs (fluoranthene, pyrene, benzo[a]anthracene, chrysene) were much lower, and after the tests, they ranged from 15.22% to 21.71% for the soil inoculated with biopreparation B1 and from 18.79% to 26.80% for the soil inoculated with biopreparation B2. The 5-ring and 6-ring PAHs were the least biodegradable. The degrees of biodegradation of 5-ring PAHs in soil inoculated with biopreparations B1 and B2 after 3 months of tests ranged from 11.42% to 11.46% and from 16.56% to 16.61%, respectively, and the degrees for 6-ring PAHs ranged from 7.96% to 8.37% (soil G6-3B1) and from 14.77% to 15.27% (soil G6-3B2), respectively. The specific progress of biodegradation of PAHs in soil G6 inoculated with biopreparations B1 and B2 is shown in Figure 4.




Figure 1. Concentrations of n-alkanes in soil inoculated with biopreparation B1 over 3 months of observation (repetition number $n = 7–10$, $p < 0.05$).




Figure 2. Concentrations of n-alkanes in soil inoculated with biopreparation B2 over 3 months of observation (repetition number $n = 7–10$, $p < 0.05$).




Figure 3. (**a**) n-C17/Pr and (**b**) n-C18/F ratios in soil inoculated with biopreparations B1 and B2 over subsequent months of remediation using the ex situ method (repetition number $n = 7–10$, $p < 0.05$).




Figure 4. List of PAH concentrations in soil inoculated with biopreparations (**a**) B1 and (**b**) B2 over 3 months (ex situ method) (repetition number $n = 7–10$, $p < 0.05$). N—naphthalene, Ac—acenaphthene, Fluo—fluorene, Phen—phenanthrene, A—anthracene, F—Fluoranthene, Pir—pyrene, BaA—benzo(a)anthracene, CH—chrysene, BbF—benzo(b)fluoranthene, BkF—benzo(k)fluoranthene, BaP—benzo(a)pyrene, DahA—dibenzo(a,h)anthracene, BghiP—benzo(g,h,i)perylene, Ind123P—indeno(1,2,3-c,d)pyrene.

The content of the individual groups of hydrocarbons before and after the biodegradation tests of petroleum products by inoculation with biopreparations made with native microorganisms (biopreparations B1 and B2) is shown in Supplementary Materials (Table S1).

2.2. Phytoremediation

The first stage of phytoremediation tests was the morphological analysis of plant organisms in the tested systems. Significant differences were observed in the development of the individual specimens of maize in soil not inoculated with biopreparations, soil inoculated with biopreparation B2 and soil inoculated with biopreparation B2 with added γ-PGA. An analysis of the morphological properties of maize in the first and sixth month of phytoremediation tests of petroleum-contaminated soil shows that the plants grew best in the system inoculated with biopreparation B2 with added γ-PGA. This is indicated by the best growth of the plants, the colour of the leaves and the development of the root. Additionally, all the maize seeds sown germinated in the first month of the tests. After 6 months of phytoremediation in system 3, the individual maize specimens reached sizes from 90 to 120 cm, and their root system was well developed. Additionally, the size of the individual leaves ranged from 30 to 50 cm, and they had an intensely green colour. Additionally, maize sown in soil inoculated with biopreparation B2 with added γ-PGA was able to develop more leaves than in soil inoculated with the biopreparation alone and in soil, which was not inoculated. In system 2, six out of eight seeds germinated, but the growth of the individual specimens was much slower. Additionally, in comparison with the system inoculated with biopreparation B2 with added γ-PGA, the individual maize specimens were less developed (smaller leaves, poorly developed root system). The maize in soil inoculated with biopreparation B2 after 6 months of bioremediation tests reached a height of 50 to 80 cm and had a moderately developed root system. The leaves of the plant reached a length of 20 to 30 cm and had a light green colour. The development of maize was the slowest by a wide margin in system 1. Out of eight seeds, only three germinated, and their sizes were much smaller than the sizes of specimens in systems 2 and 3. The sizes of the individual plants at the end of phytoremediation in system 1 ranged from 30 to 50 cm, and their root systems were very poorly developed, which hindered the efficient remediation of petroleum-contaminated soil. The appearance of the selected maize specimens in soil G6-3B2(F1), G6-3B2(F2) and G6-3B2(F3) is shown in Figures 5–7.



Figure 5. Growth and degree of germination of Zea mays after 1 month of research on phytoremediation of soil G6-3B2 under the tested systems: (**a**) system 1, (**b**) system 2, (**c**) system 3.

The morphological analysis only enables an indicative determination of the effectiveness of phytoremediation. The observations of the development of maize suggest that inoculation with biopreparation B2 increases the efficiency of phytoremediation of petroleum-contaminated soil, and the use of the γ-PGA additive with biopreparation B2 increases it further. This hypothesis is confirmed by the results of chromatographic analyses, which enable a specific determination of the changes in the concentration of hydrocarbons under the tested systems.




Figure 6. Appearance of *Zea mays* after 6 months of research on phytoremediation of soil G6-3B2 under the studied systems: (**a**) system 1, (**b**) system 2, (**c**) system 3.




Figure 7. Morphology of selected *Zea mays* individuals after completion of G6-3B2 soil phytoremediation studies under the studied systems: (**a**) system 1, (**b**) system 2, (**c**) system 3.

The developed chromatographic methods for the determination of aliphatic hydrocarbons in contaminated soil enable the identification of n-alkanes in contaminated soil after phytoremediation. As a result of the tests conducted, the highest degree of biodegradation of TPH and PAHs was achieved in the system inoculated with biopreparation B2 with added γ-PGA, and the least efficient biodegradation was recorded in the system not inoculated with biopreparations. This indicates that phytoremediation alone only slightly increases the efficiency of remediation of petroleum-contaminated soil, whereas the combination of inoculation with biopreparation with phytoremediation significantly improves its efficiency.

As a result of phytoremediation using maize, the concentration of TPH in the tested systems after 6 months of tests decreased by 22.81% (system 1), 49.08% (system 2) and 65.35% (system 3), whereas the efficiency of PAH phytoremediation was 18.48% in soil G6-3B2(F1), 40.74% in soil G6-3B2(F2) and 60.80% in soil G6-3B2(F3). The TPH and PAH concentrations in the tested systems at the end of phytoremediation are shown in Figure 8.




Figure 8. Concentration of (**a**) TPH and (**b**) PAHs in soils after completion of the phytoremediation process using *Zea mays* (repetition number $n = 7–10$, $p < 0.05$).

The results of chromatographic analyses showed that during phytoremediation, the highest efficiency was recorded for the biodegradation of aliphatic hydrocarbons with a carbon chain length of n-C_{10}–n-C_{21}, for which the degrees of biodegradation ranged from 32.99% to 53.93% (soil G6-3B2(F1)), from 70.39% to 80.21% (soil G6-3B2(F2)) and from 87.51% to 92.36% (soil G6-3B2(F3)). A satisfactory degree of biodegradation was also observed for light hydrocarbons with a carbon chain length of n-C_6–n-C_9 and hydrocarbons in the n-C_{22}–n-C_{30} range. In both cases, phytoremediation using maize was most efficient in system 3, i.e., soil inoculated with a mixture of biopreparation B2 and γ-PGA. The degrees of biodegradation of n-C_6–n-C_9 ranged from 80.00% to 92.97% and for n-C_{22}–n-C_{30}—from 59.67% to 78.54%. Hydrocarbons with fewer than 30 carbon atoms per molecule were broken down to a smaller extent, and their degrees of biodegradation for systems 1, 2 and 3 ranged from 9.58% to 11.86%, 21.36% to 27.54% and 35.52% to 51.34%, respectively. The content of unidentified hydrocarbons after 6 months of phytoremediation decreased by 21.69% in soil G6-3B2(F1), by 48.47% in soil G6-3B2(F2) and by 60.67% in soil G6-3B2(F3). A comparison of changes in the content of aliphatic hydrocarbons in soil G6-3 after phytoremediation using maize is shown in Figure 9.

The efficiency of the removal of polycyclic aromatic hydrocarbons from the soil by phytoremediation using maize was the highest in the system inoculated with biopreparation B2 with added γ-PGA, slightly lower in the system inoculated with biopreparation alone and significantly lower in the system, which was not inoculated. Out of the determined polycyclic aromatic hydrocarbons, the highest degree of biodegradation after 6 months of the experiment was recorded for naphthalene, and it amounted to 29.39% in soil G6-3B2(F1), 54.34% in soil G6-3B2(F2) and 78.87% in soil G6-3B2(F3). Slightly worse biodegradation was recorded for the 3-ring PAHs (Ac, Fluo, Fen, A), whose degrees of biodegradation ranged from 12.99% to 18.91% in system 1, from 41.70% to 49.10% in system 2 and from 60.99% to 72.84% in system 3. Through biodegradation combined with phytoremediation, the concentrations of 4-ring PAHs (F, Pir, BaA, CH) were reduced in systems 1, 2 and 3 by 5.21% to 13.98%, 20.02% to 30.55% and 32.20% to 51.50%, respectively. The 5-ring and 6-ring PAHs were less readily biodegradable, and their degrees of biodegradation ranged from 3.39% to 4.67% in soil G6-3(1), from 17.97% to 19.92% in soil G6-3(2) and from 26.67% to 31.09% in soil G6-3(3). The specific progress of biodegradation of PAHs in soil G6-3 subjected to phytoremediation under the tested systems is shown in Figure 10.




Figure 9. N-alkanes content in the tested soils after completion of the phytoremediation process. Soil G6-3B2 was the control (repetition number $n = 7–10$, $p < 0.05$).




Figure 10. PAHs content in the tested soils after completion of the phytoremediation process. Soil G6-3B2 was the control (repetition number $n = 7–10$, $p < 0.05$).

The content of the tested groups of petroleum hydrocarbons in soil at the end of the phytoremediation process (6 months) under three tested systems is shown in Supplementary Materials (Table S2).

2.3. *Toxicological Analyses*

Toxicological tests were conducted on soil samples before the remediation process (initial soil—G6), at the end of biodegradation using biopreparations B1 (soil G6-3B1) and B2 (soil G6-3B2) (3 months) and at the end of phytoremediation under the tested process systems (phytoremediation, inoculation with biopreparation B2 and inoculation

with a mixture of biopreparation B2 and γ-PGA solution) (soil G6-3B2(F1), G6-3B2(F2), G6-3B2(F3)). The results enabled the determination of changes in the toxic properties of soil during subsequent remediation stages.

2.3.1. Soil Phytotoxicity Tests Using the Phytotoxkit™ Test

The phytotoxicity tests conducted showed that the sample of the petroleum-contaminated soil (soil G6) before biodegradation was toxic to the tested plants: *Sorghum saccharatum*, *Lepidium sativum* and *Sinapis alba*. Out of this group, the plant most sensitive to pollution (TPH and PAHs) was cress, for which the root growth was inhibited by 43.1% after 72 h of the test. A slightly higher resistance to pollution was recorded for sorghum (40.2%) and mustard (39.7%). The second parameter tested, i.e., the percentage of germinated seeds, ranged from 60% to 90%.

At the end of biodegradation tests performed with the prism method using bioaugmentation with biopreparations B1 and B2, the content of petroleum hydrocarbons in the soil decreased, reducing its toxicity, as confirmed by the results of the second phytotoxicity test. For soil G6-3B1, the average increase in root length ranged from 21.5 to 35.7 mm for *Sorghum saccharatum*, from 30.2 to 47.2 mm for *Lepidium sativum* and from 40.2 to 59.8 mm for *Sinapis alba*. In the case of soil inoculated with biopreparation B2, the average root length was 39.9 mm for *Sorghum saccharatum*, 52.4 mm for *Lepidium sativum* and 66.3 mm for *Sinapis alba*. The inhibition of root growth ranged from 25.25% to 31.59% for soil G6-3B1 and from 16.88% to 24.06% for soil G6-3B2. Additionally, in both soils, *Sorghum saccharatum* and *Sinapis alba* germinated 100%, whereas the percentage of germination of *Lepidium sativum* was 70% in soil inoculated with biopreparation B1 and 80% in soil inoculated with biopreparation B2.

As a result of phytoremediation, the best reduction in toxic soil properties was achieved in the system inoculated with biopreparation B2 with added γ-PGA, where the average root length was 47.8 mm for *Sorghum saccharatum*, 68.5 mm for *Lepidium sativum* and 79.2 mm for *Sinapis alba*. Root growth inhibition ranged from 0.42% to 1%. Slightly worse test results were achieved in soil G6-3B2(F2), where the average root length of the tested plants ranged from 45.7 mm (*Sorghum saccharatum*) to 76.8 mm (*Sinapis alba*), and root growth inhibition ranged from 4.00% to 5.36%. The effect of soil remediation by phytoremediation was worst by a wide margin in system 1, where the average root length of *Sorghum saccharatum*, *Lepidium sativum* and *Sinapis alba* was 41.2 mm, 55.7 mm and 69.45 mm, respectively, and the average growth inhibition ranged from 10.6 to 19.28. A summary of the Phytotoxkit™ test results conducted during the remediation of G6 soil is included in Figure 11.




Figure 11. Results of the Phytotoxkit™ test (**a**) carried out for the original soil and after biodegradation tests with the use of biopreparations B1 and B2, (**b**) carried out for soil after completion of phytoremediation tests under the three tested systems (repetition number $n = 7–10$, $p < 0.05$).

2.3.2. Soil Toxicity Tests Using the Ostracodtoxikit^{FTM} Test

The Ostracodtoxikit^{FTM} tests showed that the initial soil G6 had toxic properties, as evidenced by the increase in the mortality of Heterocypris incongruens crustaceans to 46.67% relative to the control soil (clean soil) and inhibition of the growth of ostracods by 37.1%. The bioremediation of soil by inoculation with biopreparations made with native microorganisms reduced its toxicity to ostracods. In soil G6-3B1, the mortality of the test organisms was recorded at 25.00%, and the average growth inhibition decreased to 16.3%, while in soil G6-3B2, the mortality of ostracods decreased to 18.33%, and the average growth inhibition decreased to 11.7%.

Soil toxicity tests using the Ostracodtoxikit^{FTM} test conducted after the completion of phytoremediation of the soil using *Zea mays* showed the highest decrease in the average mortality of *Heterocypris incongruens* and average growth inhibition in soil G6-3B2(F3) of 3.33% and 0.52%, respectively. Slightly higher ostracod mortality and growth inhibition were recorded in soil G6-3B2(F2), namely 6.67% and 4.25%, respectively, while the parameters tested in soil G6-3B2(F1) were 13.33% (mortality) and 9.24% (average growth inhibition). A summary of the results of the Ostracodtoxkit^{FTM} test conducted during the remediation of G6 soil is included in Figure 12.




Figure 12. Results of the Ostracodtoxkit^{FTM} test (**a**) carried out for the original soil and after biodegradation tests with the use of biopreparations B1 and B2, (**b**) carried out for soil after completion of phytoremediation tests under the three tested systems (repetition number $n = 7–10$, $p < 0.05$).

2.3.3. Soil Toxicity Tests Using the Microtox®SPT Test

Soil toxicity tests on the trophic level of decomposers were conducted using the Microtox®SPT test (for solid phase), enabling direct contact of luminescent bacteria (*Vibrio fischeri*) with the tested soil sample. The tests showed that soil contaminated with petroleum products (TPH, PAHs) before the biodegradation process was highly toxic, as indicated by the toxic units TU = 15.7. As a result of inoculation with biopreparation B1, TU was reduced in the Microtox®SPT test to 10.1, while inoculation with biopreparation B2 reduced TU to 8.7. The phytoremediation treatments performed under the tested systems additionally reduced TU to 4.9 in soil G6-3B2(F1), 2.7 in soil G6-3B2(F2) and 1.2 in soil G6-3B2(F3). A summary of the results of the Microtox®SPT test conducted during the remediation of G6 soil is included in Figure 13.




Figure 13. Results of the Microtox®SPT test **(a)** carried out for the original soil and after biodegradation tests with the use of biopreparations B1 and B2, **(b)** carried out for soil after completion of phytoremediation tests under the three tested systems (repetition number $n = 7–10$, $p < 0.05$).

3. Discussion

The bioremediation tests using biopreparations made with native microorganisms showed differences in the efficiency of biodegradation of both aliphatic hydrocarbons and polycyclic aromatic hydrocarbons, depending on the composition of the applied biopreparation. The use of biopreparation B2 enhanced with selected species of fungi resulted in a greater reduction in aliphatic hydrocarbons in the soil than the use of B1 bacterial biopreparation [69,70]. The presence of fungi in the biopreparation had the biggest impact on the efficiency of biodegradation of aliphatic hydrocarbons with a carbon chain length of n-C_{22}–n-C_{30} and heavy hydrocarbons with a carbon chain length of n-C_{31}–n-C_{36} [27,55]. This is confirmed by studies by other researchers, indicating that *Penicillium* sp., *Candida* sp., *Aspergillus* sp. are good degraders of aliphatic hydrocarbons with carbon chain lengths of C_{10}–C_{36} [27–31]. The available research indicates that most microorganisms can metabolise hydrocarbons to a limited extent. Consequently, in order to increase the efficiency of bioremediation of soil contaminated with various groups of hydrocarbons (aliphatic hydrocarbons, PAH), multi-component mixtures with strains of bacteria and fungi with a broad spectrum of action should be used. Additionally, the efficiency of biodegradation of aliphatic hydrocarbons depends on the size and structure of the molecule [55,120]. Aliphatic hydrocarbons with a carbon chain length of n-C_{10}–n-C_{21} were the most readily biodegradable as a result of inoculation with biopreparations B1 and B2, whereas heavy hydrocarbons were the most difficult to break down. Hydrocarbons with shorter carbon chains are, therefore, more readily used by microorganisms in metabolic processes. It was generally assumed that the rate of removal of selected groups of aliphatic hydrocarbons from soil inoculated with both the bacterial biopreparation and biopreparation made with bacteria and fungi decreased in the following order: nC_{10}–nC_{21}> nC_6–nC_9 > nC_{22}–nC_{30} > nC_{31}–nC_{36} [120–123]. The situation was the same with polycyclic aromatic hydrocarbons. The inoculation of the soil tested with biopreparation B2 produced a higher decrease in PAH concentration after 3 months of tests than inoculation with biopreparation B1. Considering the structure of the hydrocarbon molecule, it is assumed that biodegradation efficiency decreases along with the increase in the number of aromatic rings in the PAH molecule. The rate of removal of polycyclic aromatic hydrocarbons from soil inoculated with biopreparations B1 and B2 decreased in the following order: 2-ring-PAH > 3-ring-PAH > 4-ring-PAH > 5-ring-PAH > 6-ring-PAH [55]. The sequence of removal of individual

groups of hydrocarbons from the soil matrix was consistent with the results obtained in previous studies and with the results obtained by other researchers [55,120–123].

The phytoremediation tests conducted using *Zea mays* demonstrated that the use of plants for the remediation of petroleum-contaminated land enabled a significant decrease in TPH and PAH concentrations in the tested soil. *Zea mays* is tolerant to high concentrations of xenobiotics and has a high rate of pollution accumulation, grows quickly, has large biomass production and is resistant to diseases and pests [124]. Beneficial results were also obtained during the use of maize in phytoremediation tests of heavy metals and petroleum hydrocarbons [97–99,124–128]. Most researchers conducting research using Zea mays in bioremediation treatments focus only on the phytoremediation process, not taking into account combined techniques, e.g., phytoremediation supported by inoculation [98,99]. Since the tests were conducted under three systems (system 1—phytoremediation; system 2—phytoremediation enhanced with inoculation with biopreparation B2; and system 3—phytoremediation enhanced with inoculation with biopreparation with added γ-PGA), it was possible to determine the optimum conditions for the remediation of soil from waste pit areas and take into account the impact of bioaugmentation on the effectiveness of the phytoremediation process. The least optimal method was the use of phytoremediation in isolation using *Zea mays*. After 6 months of tests, the efficiency of TPH biodegradation was 22.81%, and the efficiency of PAH biodegradation was 14.48%. The combination of phytoremediation with inoculation with biopreparation B2 enabled a significant increase in the efficiency of biodegradation of TPH and PAHs. The degree of reduction in TPH and PAHs in system 2 was higher than in system 1 by 26.99% and 22.26%, respectively. The most optimal solution was the use of phytoremediation using *Zea mays* in combination with inoculation with biopreparation B2 with added γ-PGA. The use of the γ-PGA additive with biopreparation B2 enabled an additional increase in the efficiency of soil remediation of 16.27% (TPH) and 20.06% (PAHs) relative to system 2. Additionally, the best plant growth was observed in system 3, as indicated by the best development of the root, stem and leaves. The results obtained from chromatographic analyses after the biodegradation process are consistent with the current knowledge, and the use of different systems of phytoremediation tests using *Zea mays* enabled an improvement of the existing methods for remediation of petroleum-contaminated soil using plant organisms [97]. Based on the chromatographic analyses conducted and observations of the morphology of the plants, it was found that the use of microorganisms and additives such as γ-PGA had a significant impact on the efficiency of soil remediation. The microorganisms present in the biopreparation can increase the bioavailability for the pollutant and the conversion of pollutants into forms more readily absorbed by the plants; they can also reduce the concentration of the pollutants to a level tolerated by the plants, as well as the presence of symbiotic interactions between the plant and the bacteria or fungi in the biopreparation [88,90,94–97]. Additionally, the better growth of plants in systems 2 and 3 may be indicative of the impact of biopreparation on the epuvalisation process, which consists of the plant using the products of bacterial breakdown to increase biomass production [129]. It should also be noted that the efficiency of phytoremediation was significantly affected by the presence of γ-PGA in the biopreparation. The available research indicates that due to its specific structure, γ-PGA is resistant to the action of microorganisms and has strong hygroscopic properties, which enables the maintenance of appropriate soil moisture—e.g., during a drought—has properties, which bind harmful substances, and can be used as an additional source of carbon [110–114].

The efficiency of the applied bioremediation treatments was complemented by toxicological monitoring intended to determine the impact of petroleum pollutants (TPH, PAHs) and indirect metabolites produced during remediation treatments on soil biocenosis based on the toxicological tests conducted. Soil inoculated with biopreparation B2 was less toxic to the indicator organisms after the tests in comparison with soil inoculated with biopreparation B1. The toxicological tests conducted on soil subjected to phytoremediation using *Zea mays* under three tested systems showed the highest decrease in soil toxicity relative to the indicator organisms tested under the system inoculated with biopreparation

B2 with added γ-PGA (system 3). The smallest decrease in toxicity in soil subjected to phytoremediation was recorded for system 1. The toxicological tests conducted confirmed the correlation between the efficiency of the removal of petroleum pollutants from the soil matrix and the decrease in its toxicity.

4. Materials and Methods

4.1. Materials

The biodegradation tests of petroleum products used soil collected from the G-6 waste pit located in south-eastern Poland (N: 49°40′16.22″, E: 22°04′42.99″). The soil tested had a high concentration of TPH (4527.70 mg/kg ds) and PAHs (10.48 mg/kg ds). The analyses of soil samples for the presence of heavy metals did not indicate any breach of the limits. The content of the individual hydrocarbon groups in the soil tested is shown in Figure 14.




Figure 14. Content (%) of individual components of petroleum pollutants in the soil: (**a**) aliphatic hydrocarbons, (**b**) PAHs (repetition number $n = 7–10$, $p < 0.05$).

For soil inoculation, two types of biopreparations developed on the basis of autochthonous microorganisms were used in the research. Bacterial biopreparation B1 consisted of the bacterial strains *Dietzia* sp. IN118, *Gordonia* sp. IN101, *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN119, *Rhodococcus globerulus* IN113 and *Raoultella* sp. IN109, and biopreparation B2 was enriched with *Aspergillus sydowii*, *Aspergillus versicolor*, *Candida* sp., *Cladosporium halotolerans*, *Penicillium chrysogenum*. The Ambiogel® (Ambioteco Sp. z o.o., Staszów, Poland) containing 5% of pure poly-γ-glutamic acid with varying molecular mass (2000–1000 kDa—30%; 1000–140 kDa—50%; 100 kDa–300 Da—15%; <300 Da—5%) was used as a γ-PGA additive [130]. The phytoremediation tests used seeds of maize (*Zea mays*) (Legutko, Poland) planted in pre-treated soil (after completing biodegradation tests using inoculation with biopreparations) in May 2022.

4.2. Methods

4.2.1. Construction of Biopreparations

The bacterial and fungal strains used in this study (bacteria: *Dietzia* sp. IN118, *Gordonia* sp. IN101, *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN119, *Rhodococcus globerulus* IN113 and *Raoultella* sp. IN109; fungi: *Aspergillus sydowii*, *Aspergillus versicolor*, *Candida sp.*, *Cladosporium halotolerans*, *Penicillium chrysogenum*) came from the hydrocarbon-degrading microbial collection of the Department of Microbiology (at the Oil and Gas Institute—National Research Institute, Poland). Some of them were previously tested, and their hydrocarbon-degrading capabilities are well established [55,56]. Bacterial

strains were phylogenetically identified following 16S rDNA sequencing analysis, as described elsewhere [121]. Strain diagnostic features were determined based on microscopic observations, morphology, growth on the selective agar media and biochemical profile (API tests, bioMerieux). The 16S rDNA sequences of bacterial strains were deposited in the NCBI database, and their accession numbers are shown in Table 1 (together with their capabilities of using selected hydrocarbons as a sole carbon and energy source).

Table 1. Hydrocarbon-degrading capabilities of bacterial strains and fungal strains comprising biopreparations B1 and B2.

Strain	NCBI Accession Number	$nC_{18}H_{38}$	Izo-$C_{19}H_{40}$	Tol + Xyl	N	A	Phen	Fluo	F	Pir
Dietzia sp. IN118	KT923327	+	+	+/−	+	−	+	−	−	+
Gordonia sp. IN101	KT923337	+	+	+	+	−	+/−	−	−	−
Mycolicibacterium frederiksbergense IN53	JN572675	+	+	−	+	+	+	−	−	+
Raoultella sp. IN109	KT923339	+	+	+/−	+	−	+	−	−	+/−
Rhodococcus erythropolis IN119	KT923331	+	+	+	+	−	−	−	−	−
Rhodococcus globerulus IN113	KT923347	+	+	+	+	+/−	+/−	+/−	−	+/−
Aspergillus sydowii	n/a	+	n/t	−	−	−	+	n/t	n/t	n/t
Aspergillus versicolor	n/a	+	n/t	+	+	+	+	n/t	n/t	n/t
Candida sp.	n/a	+	n/t	+/−	+	−	−	n/t	n/t	n/t
Cladosporium halotolerans	n/a	+	n/t	−	+	+	+	n/t	n/t	n/t
Penicillium chrysogenum	n/a	+	n/t	+	+	−	−	n/t	n/t	n/t

iso-$C_{19}H_{40}$: pristane, Tol: toluene, Xyl: mixture of xylenes, N: naphthalene, A: anthracene, Phen: phenanthrene, Fluo: fluorene, F: fluoranthene, Pir: pyrene, +: growth, −: no growth, +/−: ambiguous observation, n/a: not applicable, n/t: not tested.

The hydrocarbon-metabolising capabilities of all microbial strains were examined based on growth on the mineral medium supplemented with the compounds tested, as described in our previous work and according to the methods of Wrenn and Venosa [131]. Each bacterial strain was grown individually in a nutrient broth supplemented with sodium acetate (0.2%, *w*/*v*), while each fungal strain was grown in a Czapek–Dox broth (both BD Difco™) and incubated at room temperature with shaking at 150 rpm for 24–72 h to obtain density 10^8–10^9 cfu/mL. The biopreparations used in this study were constructed by mixing equal volumes of each strain. Then, the obtained mixed culture was kept in a nutrient broth supplemented with sodium acetate (0.2%, *w*/*v*) and sucrose and occasionally examined for the presence of all introduced strains. Table 2 presents the putative genes involved in petroleum hydrocarbon degradation found in genomes of the closest relatives of all bacterial strains used in this study [55,56].

Table 2. Identification of bacterial strains used in biopreparations B1 and B2.

Strain	The Closest Relative Based on 16S rRNA Accession Number (% of Identity) *	The Closest Relative for Which the Genome Sequence Is Available in NCBI GenBank, Accession Number (% of Identity) *	Putative Gene Encoding for the Enzymes Catalysing the Breakdown of Hydrocarbons
Dietzia sp. IN118	*Dietzia* sp. LJ3 MG049763 99.93%	*Dietzia kunjamensis* 313 CP099712 99.78%	Alkane 1-monooxygenase, aromatic ring-hydroxylating dioxygenase subunit alpha
Gordonia sp.IN101	*Gordonia* sp. Tm-B24 MT533993 99.63%	*Gordonia terrae* RL-JC02 CP049836 99.63%	Alkane 1-monooxygenase, pentachlorophenol monooxygenase, naphthalene 1,2-dioxygenase subunit alpha (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Mycolicibacterium frederiksbergense IN53	*Mycolicibacterium frederiksbergense* DSM 44346 (typical strain) NR_025393.1 99.58%	*Mycolicibacterium frederiksbergense* LB 501T 99.58%	Alkane 1-monooxygenase (2 copies), pentachlorophenol monooxygenase, naphthalene 1,2-dioxygenase subunit alpha (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Rhodococcus erythropolis IN119	*Rhodococcus erythropolis* KD-1 CP050124 99.42%	*Rhodococcus erythropolis* KD-1 CP050124 99.42%	Alkane 1-monooxygenase (5 copies), pentachlorophenol monooxygenase, cyclohexanone monooxygenase (2 copies), biphenyl 2,3-dioxygenase (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Rhodococcus globerulus IN113	*Rhodococcus globerulus* D757 CP079698 99.86%	*Rhodococcus globerulus* D757 CP079698 99.86%	Alkane 1-monooxygenase (3 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies), aromatic ring-hydroxylating dioxygenase subunit alpha (7 copies)
Raoultella sp. IN109	*Raoultella planticola* SCLZS62 CP082168 99.08%	*Raoultella planticola* SCLZS62 CP082168 99.08%	aromatic ring-hydroxylating dioxygenase subunit alpha (2 copies)

* the data presented concern the data gathered in the NCBI GenBank database (June 2023). Therefore, strain IN118 is presented here as *Dietzia* sp. instead of *Dietzia maris* (as it was originally identified).

4.2.2. Identification of Fungal Strains by MALDI-TOF MS

Matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry (MALDI-TOF MS) was used to identify specific fungal strains and was performed at the Jagiellonian Centre of Innovation (Kraków, Poland). A colony from a fresh overnight culture was used for sample preparation to obtain the mass spectrometer measurement. The material was suspended in 300 µL double-distilled water, following which, 900 µL ethanol was added, and the components were mixed well; thereafter, the mixture was centrifuged (2 min, 9000× *g*), and the supernatant was removed. For sample extraction, 40 µL of formic acid (70% in water) was added to the fungal pellet, following which, the components were mixed thoroughly, and 40 µL of acetonitrile was added. After performing centrifugation at 9000× *g* for 2 min, 1 µL of the supernatant containing the fungal extract was transferred to a well of the 96-well MALDI Biotarget plate (Bruker Daltonics, Bremen, Germany) and was allowed to dry at room temperature. Subsequently, the sample was overlaid with 2 µL of the MALDI matrix solution (α-cyano-4-hydroxy-cinnamic acid) and air-dried again. The measurements were performed using the MALDI BioTyper 2.0 Microflex LT

system (Bruker), using the manufacturer's recommended settings. In order to perform microorganisms identification, the raw protein spectral data were imported into the MALDI BioTyper 2.0 software (Bruker) and were analysed via standard pattern matching (with default parameter settings) against the spectral data in the BioTyper database (Bruker). Parameters such as the mass-to-charge ratio and peak intensity were considered to produce a matching score, which was subsequently used to rank the results. In these studies, the identification criteria were as follows: a score of 2.000 and above was considered sufficient for correct species identification; score values of 1.700 to 1.999 confirmed genus identification; and scores lower than 1.700 indicated no identification.

4.2.3. TPH and PAH Analyses

The determination of TPH and PAH was carried out according to the procedures developed by the Oil and Gas Institute—State Research Institute (Kraków, Poland).

The TPH analysis was conducted using the gas chromatography method, employing the Clarus 500 chromatograph (Perkin Elmer, Waltham, MA, USA) with a flame ionisation detector (FID). Before the chromatographic analysis could be conducted, it was necessary to correctly prepare the soil samples, including the following:

- drying soil samples to an air-dry condition;
- isolating the analytes using ultrasound-assisted extraction with dichloromethane (Chempur, Piekary Śląskie, Poland);
- purifying the extract using the dSPE method, employing Florisil columns Chromabond No. 30081 (Macherey-Nagel, Düren, Germany);
- concentrating the extract to a volume of 1 mL in a rotary evaporator (Chemland, Stargard, Poland).

Then, the produced extract was subjected to a chromatographic analysis using the following operating parameters of the chromatograph: capillary column of fused silica (RTX-1:30 m × 0.53 mm) (Restek, Bellefonte, PA, USA); using the following temperature parameters: injector temperatures = 290 °C, FID temperature = 320 °C; using the following oven temperature programme: 30 °C—isothermal run for 2 min, 30 °C to 105 °C—temperature increase at a rate of 10 °C/min, 105 °C to 285 °C—temperature increase at a rate of 5 °C/min, 285 °C—isothermal run for 5 min [55,56]. The diagram of the TPH analysis is shown in Figure 15.

The PAH analysis was conducted using the HPLC liquid chromatography method, employing the Vanquish Core chromatography set (Thermo Scientific, Waltham, MA, USA). The analysis of the quantitative and qualitative determination of 16 PAHs in the soil samples required the preparation of an analytical sample, which included the following:

- drying soil samples to an air-dry condition;
- isolating the analytes (PAHs) using the QuEChERS method;
- purifying the extract using the dSPE method with $MgSO_4$ and PSA vials No. JO3937 (Interchim, Montluçon Cedex, France).

Then, the produced extract was subjected to a chromatographic analysis with the following operating parameters of the equipment: NUCLEODUR C_{18} PAH column, 125 × 4 mm, 3 µm (Marcherey-Nagel, Düren, Germany); detector: UV-VIS; fluorometric (FLD) dispenser: automatic; dispensing volume: 10 µL; eluents: A-methanol (POCH S.A., Poland) 70%, B-acetonitrile (POCH S.A., Gliwice, Poland); flow rate: 1.5 mL/min; gradient: 20% B for 1.5 min, 20% to 50% B for 1.5 min, 50% to 100% B for 1 min, 100% B for 1 min, 100% to 0% B for 3 min, 100% A for 3 min [55]. The diagram of the PAH analysis is shown in Figure 15.

The specific method for the determination of TPH and PAHs in soil samples was described elsewhere [55,56].




Figure 15. Scheme for determination of TPH and PAHs in soil samples.

4.2.4. Ecotoxicological Analyses

Soil toxicology analysis was performed using commercially available toxicology tests Phytotoxkit ™ (MicroBioTests Inc., Nazareth, Belgium), Ostracodtoxkit ™ (MicroBioTests Inc., Nazareth, Belgium), Microtox® Solid Phase Test (SDI, Newark, DE, USA). The use of living organisms as bioindicators belonging to various taxonomic groups (higher plants, crustaceans, bacteria) representing all trophic groups (producers, consumers and decomposers) allowed for a comprehensive assessment of the effectiveness of the process for cleaning soil contaminated with oil derivatives.

Toxicological tests were carried out in soil samples taken before process purification (initial soil—G6), after completion of biodegradation with biopreparations B1 (soil G6-3B1) and B2 (soil G6-3B2) and after completion of phytoremediation under the systems tested (soil G6-3B2(F1), G6-3B2(F2), G6-3B2(F3)). A detailed discussion of the toxicological tests used and the description of the procedures for performing them are provided in Supplementary Materials and elsewhere [55,56].

4.2.5. Data Analysis and Statistical Information

Statistical analysis was performed with Statistica 14.0. Standard deviation (SD), standard deviation (RSD) (%) and Pearson correlation coefficient were calculated. The obtained data were first tested for normal distribution. Then, they were analysed with one-way ANOVA, followed by a post hoc pairwise Tukey test (when the ANOVA produced significant results). Significance was set at $p < 0.05$ [55,56].

4.3. Experiment Description

4.3.1. Biodegradation by Inoculation

Biodegradation tests using inoculation with biopreparations made with native microorganisms were conducted using the ex situ prism method. The layer of gravel used for the placement of perforated pipes responsible for the supply of air to the site was covered with soil collected from the G6 mining pit in the form of two identical prisms with a weight of 60 kg each. The soil was tested for the presence of petroleum products, the concentration of heavy metals and the C:N:P ratio. To ensure the optimum conditions for the development of microorganisms in the biopreparation, the C:N:P ratio was adjusted via supplementation with mineral fertiliser Azofoska (Inco, Warszawa, Poland), and the pH of the soil was adjusted using fertiliser lime (Biovita, Tenczynek, Poland). The soil prepared in this way had a C:N:P ratio of 100:10:1 and a pH of 7.5. Then, each prism was inoculated with biopreparation B1 (prism No. 1) or biopreparation B2 (prism No. 2) in the amount of 6 L per prism. Biodegradation tests using inoculation with biopreparations made with native microorganisms were conducted for 3 months at a constant temperature of 25 °C and constant humidity (20% to 25%) [55,56]. A diagram of the stand for testing the process of biodegradation is shown in Supplementary Materials (Figure S1).

4.3.2. Phytoremediation

The phytoremediation tests were conducted using the pot method on a semi-technical scale. Pots with a volume of 10 L were filled with soil (15 kg), which had been pre-treated by inoculation with the biopreparation selected based on biodegradation tests of petroleum pollutants (biopreparation B2). The soil tested had a known concentration of petroleum pollutants (TPH = 2640.83 mg/kg ds and PAHs = 6.84 mg/kg ds), a suitable C:N:P ratio of 100:10:1 and humidity of 20%, which enabled optimum conditions for the remediation process. Then, 8 seeds of maize were sown in each pot, and, depending on the system tested, 1 litre of water, biopreparation B2 or biopreparation B2 with added γ-PGA was added to each pot.

- system 1—soil G6-3B2(F1)—(soil G6-3B2 + *Zea mays*);
- system 2—soil G6-3B2(F2)—(soil G6-3B2 + biopreparation B2 + *Zea mays*);
- system 3—soil G6-3B2(F3)—(soil G6-3B2 + biopreparation B2 with added γ-PGA + *Zea mays*).

Each of the systems tested was covered with foil until the plants germinated. The phytoremediation tests were carried out for 6 months in the actual conditions to test the possibility of using the proposed method in field conditions (waste pits). The layout of the site for phytoremediation tests is shown in Supplementary Materials (Figure S2).

5. Conclusions

Petroleum hydrocarbons are the leading group of pollutants generated by the oil and gas extraction industry. Due to their carcinogenic, neurotoxic and mutagenic properties, oil-derived pollutants pose a serious threat to the environment, as well as to the health and life of humans and animals. For this reason, it is extremely important to develop effective methods of cleaning up oil-contaminated areas. In our work, we indicated the enormous potential for bioremediation of soils contaminated with TPH and PAHs using bioaugmentation and bioaugmentation-assisted phytoremediation techniques. Biodegradation studies conducted using the ex situ piling method with the use of two types of biopreparations showed that the addition of fungi to the biopreparation significantly increases the biodegradation efficiency of both aliphatic and aromatic hydrocarbons. This confirms the assumption that the use of more strains of microorganisms in the biopreparation provides it with a broad spectrum of activity. Currently, more and more research is being conducted on accelerating the process of bioremediation of soils contaminated with toxic substances by combining several remediation techniques or their modifications. The combination of phytoremediation and bioaugmentation enabled an additional reduction in the con-

centration of petroleum hydrocarbons in the soil tested and an increase in the efficiency of its treatment of 49.08% (TPH) and 40.74% (PAH). The use of γ-PGA as an additive to the biopreparation allowed for increased efficiency of phytoremediation supported by bioaugmentation of 16.27% (TPH) and 20.06% (PAH). The assessment of the effectiveness of the applied bioremediation treatments included the results of toxicological tests carried out using the Phytotoxkit ™, Ostracodtoxkit ™, Microtox® Solid Phase test kits. Depending on the test used, soil toxicity varied, but it was correlated with the concentration of pollutants contained in the soil.

Promising results obtained during biodegradation tests with the use of biopreparations B1 and B2 and phytoremediation (using *Zea mays* as a phytoremediant) supported by bioaugmentation may in the future be a step forward in the methods of bioremediation of soils contaminated with TPH and PAHs and may be used in field conditions.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/molecules28166104/s1, Table S1—Total content of selected groups of TPH and PAHs in soil G6 and after the biodegradation process with the use of biopreparation B1 (soil G6-3B1) and B2 (soil G6-3B2), Table S2—Total content of selected groups of TPH and PAHs in soil after the phytoremediation process in the tested systems, Ecotoxicological analyses—description of the methodology, Figure S1—Stand for testing the process of biodegradation of petroleum hydrocarbons, Figure S2—Stand for testing the process of phytoremediation of soil G6-3B2.

Author Contributions: Conceptualisation, K.W. and T.S.; software, K.W.; validation, K.W. and P.K.; formal analysis, K.W., T.S., P.K. and J.B.; investigation, K.W., T.S., P.K. and J.B.; writing—original draft preparation, K.W. and T.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was financially supported by the Polish Ministry of Education and Science within statutory funding for the Oil and Gas Institute—National Research Institute (Kraków, Poland).

Institutional Review Board Statement: Not applicable. The studies used commercial toxicological tests not subject to ethical evaluation.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

Sample Availability: Not applicable.

References

1. Steliga, T. Bioremediacja odpadów wiertniczych zanieczyszczonych substancjami ropopochodnymi ze starych dołów urobkowych. *Pr. Inst. Naft. Gazu* **2009**, *163*, 1–331.
2. Kluk, D.; Steliga, T. Efektywna Metoda Identyfikacji Zanieczyszczeń Ropopochodnych (TPH) i Wielopierścieniowych Węglowodorów Aromatycznych (WWA) w Glebach. *Naft. Gaz Sci. Technol. Oil Gas Ind.* **2017**, *73*, 488–495. [CrossRef]
3. Jaekel, U.; Musat, N.; Adam, B.; Kuypers, M.; Grundmann, O.; Musat, F. Anaerobic Degradation of Propane and Butane by Sulfate-Reducing Bacteria Enriched from Marine Hydrocarbon Cold Seeps. *ISME J.* **2013**, *7*, 885–895. [CrossRef]
4. Kurth, E.G.; Doughty, D.M.; Bottomley, P.J.; Arp, D.J.; Sayavedra-Soto, L.A. Involvement of BmoR and BmoG in N-Alkane Metabolism in 'Pseudomonas Butanovora'. *Microbiology* **2008**, *154*, 139–147. [CrossRef] [PubMed]
5. Kotani, T.; Kawashima, Y.; Yurimoto, H.; Kato, N.; Sakai, Y. Gene Structure and Regulation of Alkane Monooxygenases in Propane-Utilizing *Mycobacterium* sp. TY-6 and *Pseudonocardia* sp. TY-7. *J. Biosci. Bioeng.* **2006**, *102*, 184–192. [CrossRef]
6. Mishra, S.; Jyot, J.; Kuhad, R.C.; Lal, B. Evaluation of Inoculum Addition to Stimulate In Situ Bioremediation of Oily-Sludge-Contaminated Soil. *Appl. Environ. Microbiol.* **2001**, *67*, 1675–1681. [CrossRef] [PubMed]
7. Sabirova, J.S.; Ferrer, M.; Regenhardt, D.; Timmis, K.N.; Golyshin, P.N. Proteomic Insights into Metabolic Adaptations in *Alcanivorax Borkumensis* Induced by Alkane Utilization. *J. Bacteriol.* **2006**, *188*, 3763–3773. [CrossRef]
8. Wang, X.-B.; Chi, C.-Q.; Nie, Y.; Tang, Y.-Q.; Tan, Y.; Wu, G.; Wu, X.-L. Degradation of Petroleum Hydrocarbons (C6–C40) and Crude Oil by a Novel Dietzia Strain. *Bioresour. Technol.* **2011**, *102*, 7755–7761. [CrossRef]
9. Kaplan, C.W.; Kitts, C.L. Bacterial Succession in a Petroleum Land Treatment Unit. *Appl. Environ. Microbiol.* **2004**, *70*, 1777–1786. [CrossRef]
10. Abbasian, F.; Lockington, R.; Mallavarapu, M.; Naidu, R. A Comprehensive Review of Aliphatic Hydrocarbon Biodegradation by Bacteria. *Appl. Biochem. Biotechnol.* **2015**, *176*, 670–699. [CrossRef]

11. Brown, L.M.; Gunasekera, T.S.; Striebich, R.C.; Ruiz, O.N. Draft Genome Sequence of Gordonia Sihwensis Strain 9, a Branched Alkane-Degrading Bacterium. *Genome Announc.* **2016**, *4*, e00622-16. [CrossRef] [PubMed]
12. Churchill, S.A.; Harper, J.P.; Churchill, P.F. Isolation and Characterization of a *Mycobacterium* Species Capable of Degrading Three- and Four-Ring Aromatic and Aliphatic Hydrocarbons. *Appl. Environ. Microbiol.* **1999**, *65*, 549–552. [CrossRef] [PubMed]
13. Brzeszcz, J.; Steliga, T.; Kapusta, P.; Turkiewicz, A.; Kaszycki, P. R-Strategist versus K-Strategist for the Application in Bioremediation of Hydrocarbon-Contaminated Soils. *Int. Biodeterior. Biodegrad.* **2016**, *106*, 41–52. [CrossRef]
14. Yakimov, M.M.; Giuliano, L.; Gentile, G.; Crisafi, E.; Chernikova, T.N.; Abraham, W.-R.; Lünsdorf, H.; Timmis, K.N.; Golyshin, P.N. *Oleispira antarctica* gen. nov., sp. nov., a Novel Hydrocarbonoclastic Marine Bacterium Isolated from Antarctic Coastal Sea Water. *Int. J. Syst. Evol. Microbiol.* **2003**, *53*, 779–785. [CrossRef]
15. Röling, W.F.M.; Milner, M.G.; Jones, D.M.; Fratepietro, F.; Swannell, R.P.J.; Daniel, F.; Head, I.M. Bacterial Community Dynamics and Hydrocarbon Degradation during a Field-Scale Evaluation of Bioremediation on a Mudflat Beach Contaminated with Buried Oil. *Appl. Environ. Microbiol.* **2004**, *70*, 2603–2613. [CrossRef]
16. Abalos, A.; Viñas, M.; Sabaté, J.; Manresa, M.A.; Solanas, A.M. Enhanced Biodegradation of Casablanca Crude Oil by A Microbial Consortium in Presence of a Rhamnolipid Produced by Pseudomonas Aeruginosa AT10. *Biodegradation* **2004**, *15*, 249–260. [CrossRef]
17. Al-Hadhrami, M.N.; Lappin-Scott, H.M.; Fisher, P.J. Studies on the Biodegradation of Three Groups of Pure N-Alkanes in the Presence of Molasses and Mineral Fertilizer by Pseudomonas Aeruginosa. *Mar. Pollut. Bull.* **1997**, *34*, 969–974. [CrossRef]
18. Karamalidis, A.K.; Evangelou, A.C.; Karabika, E.; Koukkou, A.I.; Drainas, C.; Voudrias, E.A. Laboratory Scale Bioremediation of Petroleum-Contaminated Soil by Indigenous Microorganisms and Added Pseudomonas Aeruginosa Strain Spet. *Bioresour. Technol.* **2010**, *101*, 6545–6552. [CrossRef]
19. Varjani, S.J.; Rana, D.P.; Jain, A.K.; Bateja, S.; Upasani, V.N. Synergistic Ex-Situ Biodegradation of Crude Oil by Halotolerant Bacterial Consortium of Indigenous Strains Isolated from on Shore Sites of Gujarat, India. *Int. Biodeterior. Biodegrad.* **2015**, *103*, 116–124. [CrossRef]
20. Zhang, Z.; Hou, Z.; Yang, C.; Ma, C.; Tao, F.; Xu, P. Degradation of N-Alkanes and Polycyclic Aromatic Hydrocarbons in Petroleum by a Newly Isolated Pseudomonas Aeruginosa DQ8. *Bioresour. Technol.* **2011**, *102*, 4111–4116. [CrossRef]
21. Popp, N.; Schlömann, M.; Mau, M. Bacterial Diversity in the Active Stage of a Bioremediation System for Mineral Oil Hydrocarbon-Contaminated Soils. *Microbiology* **2006**, *152*, 3291–3304. [CrossRef] [PubMed]
22. Bhattacharya, D.; Sarma, P.M.; Krishnan, S.; Mishra, S.; Lal, B. Evaluation of Genetic Diversity among *Pseudomonas citronellolis* Strains Isolated from Oily Sludge-Contaminated Sites. *Appl. Environ. Microbiol.* **2003**, *69*, 1435–1441. [CrossRef] [PubMed]
23. Lalucat, J.; Bennasar, A.; Bosch, R.; García-Valdés, E.; Palleroni, N.J. Biology of *Pseudomonas stutzeri*. *Microbiol. Mol. Biol. Rev.* **2006**, *70*, 510–547. [CrossRef]
24. Whyte, L.G.; Slagman, S.J.; Pietrantonio, F.; Bourbonnière, L.; Koval, S.F.; Lawrence, J.R.; Inniss, W.E.; Greer, C.W. Physiological Adaptations Involved in Alkane Assimilation at a Low Temperature by *Rhodococcus* sp. Strain Q15. *Appl. Environ. Microbiol.* **1999**, *65*, 2961–2968. [CrossRef]
25. Hamamura, N.; Olson, S.H.; Ward, D.M.; Inskeep, W.P. Microbial Population Dynamics Associated with Crude-Oil Biodegradation in Diverse Soils. *Appl. Environ. Microbiol.* **2006**, *72*, 6316–6324. [CrossRef] [PubMed]
26. Zhukov, D.V.; Murygina, V.P.; Kalyuzhnyi, S.V. Kinetics of the Degradation of Aliphatic Hydrocarbons by the Bacteria Rhodococcus Ruber and Rhodococcus Erythropolis. *Appl. Biochem. Microbiol.* **2007**, *43*, 587–592. [CrossRef]
27. Barnes, N.M.; Khodse, V.B.; Lotlikar, N.P.; Meena, R.M.; Damare, S.R. Bioremediation Potential of Hydrocarbon-Utilizing Fungi from Select Marine Niches of India. *3 Biotech* **2018**, *8*, 21. [CrossRef]
28. Daccò, C.; Girometta, C.; Asemoloye, M.D.; Carpani, G.; Picco, A.M.; Tosi, S. Key Fungal Degradation Patterns, Enzymes and Their Applications for the Removal of Aliphatic Hydrocarbons in Polluted Soils: A Review. *Int. Biodeterior. Biodegrad.* **2020**, *147*, 104866. [CrossRef]
29. Yang, S.; Zhang, J.; Liu, Y.; Feng, W. Biodegradation of Hydrocarbons by Purpureocillium Lilacinum and Penicillium Chrysogenum from Heavy Oil Sludge and Their Potential for Bioremediation of Contaminated Soils. *Int. Biodeterior. Biodegrad.* **2023**, *178*, 105566. [CrossRef]
30. Li, J.; Xu, Y.; Song, Q.; Zhang, S.; Xie, L.; Yang, J. Transmembrane Transport Mechanism of N-Hexadecane by Candida Tropicalis: Kinetic Study and Proteomic Analysis. *Ecotoxicol. Environ. Saf.* **2021**, *209*, 111789. [CrossRef]
31. Al-Otibi, F.; Al-Zahrani, R.M.; Marraiki, N. The Crude Oil Biodegradation Activity of Candida Strains Isolated from Oil-Reservoirs Soils in Saudi Arabia. *Sci. Rep.* **2022**, *12*, 10708. [CrossRef] [PubMed]
32. Ma, Y.-L.; Lu, W.; Wan, L.-L.; Luo, N. Elucidation of Fluoranthene Degradative Characteristics in a Newly Isolated Achromobacter Xylosoxidans DN002. *Appl. Biochem. Biotechnol.* **2015**, *175*, 1294–1305. [CrossRef] [PubMed]
33. Mnif, S.; Sayadi, S.; Chamkha, M. Biodegradative Potential and Characterization of a Novel Aromatic-Degrading Bacterium Isolated from a Geothermal Oil Field under Saline and Thermophilic Conditions. *Int. Biodeterior. Biodegrad.* **2014**, *86*, 258–264. [CrossRef]
34. Cerniglia, C.E. Biodegradation of Polycyclic Aromatic Hydrocarbons. *Biodegradation* **1992**, *3*, 351–368. [CrossRef]
35. Eskandari, S.; Hoodaji, M.; Tahmourespour, A.; Abdollahi, A.; Baghi, T.; Eslamian, S.; Ostad-Ali-Askari, K. Bioremediation of Polycyclic Aromatic Hydrocarbons by Bacillus Licheniformis ATHE9 and Bacillus Mojavensis ATHE13 as Newly Strains Isolated from Oil-Contaminated Soil. *J. Geogr. Environ. Earth Sci. Int.* **2017**, *11*, 1–11. [CrossRef]

36. Xiao, J.; Guo, L.; Wang, S.; Lu, Y. Comparative Impact of Cadmium on Two Phenanthrene-Degrading Bacteria Isolated from Cadmium and Phenanthrene Co-Contaminated Soil in China. *J. Hazard. Mater.* **2010**, *174*, 818–823. [CrossRef]
37. Kasai, Y.; Kishira, H.; Harayama, S. Bacteria Belonging to the Genus *Cycloclasticus* Play a Primary Role in the Degradation of Aromatic Hydrocarbons Released in a Marine Environment. *Appl. Environ. Microbiol.* **2002**, *68*, 5625–5633. [CrossRef]
38. Juhasz, A.L.; Naidu, R. Bioremediation of High Molecular Weight Polycyclic Aromatic Hydrocarbons: A Review of the Microbial Degradation of Benzo[a]Pyrene. *Int. Biodeterior. Biodegrad.* **2000**, *45*, 57–88. [CrossRef]
39. Kanaly, R.A.; Harayama, S. Biodegradation of High-Molecular-Weight Polycyclic Aromatic Hydrocarbons by Bacteria. *J. Bacteriol.* **2000**, *182*, 2059–2067. [CrossRef]
40. Zhang, Y.; Wang, F.; Zhu, X.; Zeng, J.; Zhao, Q.; Jiang, X. Extracellular Polymeric Substances Govern the Development of Biofilm and Mass Transfer of Polycyclic Aromatic Hydrocarbons for Improved Biodegradation. *Bioresour. Technol.* **2015**, *193*, 274–280. [CrossRef]
41. Rehmann, K.; Hertkorn, N.; Kettrup, A.A. Fluoranthene Metabolism in *Mycobacterium* sp. Strain KR20: Identity of Pathway Intermediates during Degradation and Growth. *Microbiology* **2001**, *147*, 2783–2794. [CrossRef] [PubMed]
42. Moody, J.D.; Freeman, J.P.; Doerge, D.R.; Cerniglia, C.E. Degradation of Phenanthrene and Anthracene by Cell Suspensions of *Mycobacterium* sp. Strain PYR-1. *Appl. Environ. Microbiol.* **2001**, *67*, 1476–1483. [CrossRef] [PubMed]
43. Kim, S.-J.; Kweon, O.; Freeman, J.P.; Jones, R.C.; Adjei, M.D.; Jhoo, J.-W.; Edmondson, R.D.; Cerniglia, C.E. Molecular Cloning and Expression of Genes Encoding a Novel Dioxygenase Involved in Low- and High-Molecular-Weight Polycyclic Aromatic Hydrocarbon Degradation in *Mycobacterium Vanbaalenii* PYR-1. *Appl. Environ. Microbiol.* **2006**, *72*, 1045–1054. [CrossRef] [PubMed]
44. Kim, S.-J.; Kweon, O.; Jones, R.C.; Freeman, J.P.; Edmondson, R.D.; Cerniglia, C.E. Complete and Integrated Pyrene Degradation Pathway in *Mycobacterium vanbaalenii* PYR-1 Based on Systems Biology. *J. Bacteriol.* **2007**, *189*, 464–472. [CrossRef]
45. Kweon, O.; Kim, S.-J.; Jones, R.C.; Freeman, J.P.; Adjei, M.D.; Edmondson, R.D.; Cerniglia, C.E. A Polyomic Approach to Elucidate the Fluoranthene-Degradative Pathway in *Mycobacterium vanbaalenii* PYR-1. *J. Bacteriol.* **2007**, *189*, 4635–4647. [CrossRef] [PubMed]
46. Hedlund, B.P.; Geiselbrecht, A.D.; Bair, T.J.; Staley, J.T. Polycyclic Aromatic Hydrocarbon Degradation by a New Marine Bacterium, *Neptunomonas naphthovorans* gen. nov., sp. nov. *Appl. Environ. Microbiol.* **1999**, *65*, 251–259. [CrossRef]
47. Saito, A.; Iwabuchi, T.; Harayama, S. A Novel Phenanthrene Dioxygenase from *Nocardioides* sp. Strain KP7: Expression in *Escherichia coli*. *J. Bacteriol.* **2000**, *182*, 2134–2141. [CrossRef]
48. Jiang, J.; Liu, H.; Li, Q.; Gao, N.; Yao, Y.; Xu, H. Combined Remediation of Cd–Phenanthrene Co-Contaminated Soil by Pleurotus Cornucopiae and Bacillus Thuringiensis FQ1 and the Antioxidant Responses in Pleurotus Cornucopiae. *Ecotoxicol. Environ. Saf.* **2015**, *120*, 386–393. [CrossRef]
49. Brito, E.M.S.; De La Cruz Barrón, M.; Caretta, C.A.; Goñi-Urriza, M.; Andrade, L.H.; Cuevas-Rodríguez, G.; Malm, O.; Torres, J.P.M.; Simon, M.; Guyoneaud, R. Impact of Hydrocarbons, PCBs and Heavy Metals on Bacterial Communities in Lerma River, Salamanca, Mexico: Investigation of Hydrocarbon Degradation Potential. *Sci. Total Environ.* **2015**, *521–522*, 1–10. [CrossRef]
50. Ghosh, I.; Jasmine, J.; Mukherji, S. Biodegradation of Pyrene by a Pseudomonas Aeruginosa Strain RS1 Isolated from Refinery Sludge. *Bioresour. Technol.* **2014**, *166*, 548–558. [CrossRef]
51. Yang, Y.; Chen, R.F.; Shiaris, M.P. Metabolism of Naphthalene, Fluorene, and Phenanthrene: Preliminary Characterization of a Cloned Gene Cluster from Pseudomonas Putida NCIB 9816. *J. Bacteriol.* **1994**, *176*, 2158–2164. [CrossRef] [PubMed]
52. Singh, P.; Tiwary, B.N. Optimization of Conditions for Polycyclic Aromatic Hydrocarbons (PAHs) Degradation by Pseudomonas Stutzeri P2 Isolated from Chirimiri Coal Mines. *Biocatal. Agric. Biotechnol.* **2017**, *10*, 20–29. [CrossRef]
53. Peng, T.; Kan, J.; Hu, J.; Hu, Z. Genes and Novel sRNAs Involved in PAHs Degradation in Marine Bacteria *Rhodococcus* sp. P14 Revealed by the Genome and Transcriptome Analysis. *3 Biotech* **2020**, *10*, 140. [CrossRef] [PubMed]
54. Lang, F.S.; Destain, J.; Delvigne, F.; Druart, P.; Ongena, M.; Thonart, P. Biodegradation of Polycyclic Aromatic Hydrocarbons in Mangrove Sediments Under Different Strategies: Natural Attenuation, Biostimulation, and Bioaugmentation with Rhodococcus Erythropolis T902.1. *Water Air Soil Pollut.* **2016**, *227*, 297. [CrossRef]
55. Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J.; Skalski, T. Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil. *Materials* **2022**, *15*, 400. [CrossRef]
56. Steliga, T.; Wojtowicz, K.; Kapusta, P.; Brzeszcz, J. Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture. *Molecules* **2020**, *25*, 709. [CrossRef]
57. Ghosal, D.; Ghosh, S.; Dutta, T.K.; Ahn, Y. Current State of Knowledge in Microbial Degradation of Polycyclic Aromatic Hydrocarbons (PAHs): A Review. *Front. Microbiol.* **2016**, *7*, 1369. [CrossRef]
58. Shuttleworth, K.L.; Sung, J.; Kim, E.; Cerniglia, C.E. Physiological and Genetic Comparison of Two Aromatic Hydrocarbon-Degrading Sphingomonas Strains. *Mol. Cells* **2000**, *10*, 199–205. [CrossRef]
59. Zylstra, G.J.; Kim, E. Aromatic Hydrocarbon Degradation by *Sphingomonas yanoikuyae* B1. *J. Ind. Microbiol. Biotechnol.* **1997**, *19*, 408–414. [CrossRef]
60. Moscoso, F.; Teijiz, I.; Deive, F.J.; Sanromán, M.A. Efficient PAHs Biodegradation by a Bacterial Consortium at Flask and Bioreactor Scale. *Bioresour. Technol.* **2012**, *119*, 270–276. [CrossRef]
61. Tiwari, B.; Manickam, N.; Kumari, S.; Tiwari, A. Biodegradation and Dissolution of Polyaromatic Hydrocarbons by *Stenotrophomonas* sp. *Bioresour. Technol.* **2016**, *216*, 1102–1105. [CrossRef] [PubMed]

62. Chen, S.; Yin, H.; Ye, J.; Peng, H.; Liu, Z.; Dang, Z.; Chang, J. Influence of Co-Existed Benzo[a]Pyrene and Copper on the Cellular Characteristics of *Stenotrophomonas maltophilia* during Biodegradation and Transformation. *Bioresour. Technol.* **2014**, *158*, 181–187. [CrossRef] [PubMed]
63. Balachandran, C.; Duraipandiyan, V.; Balakrishna, K.; Ignacimuthu, S. Petroleum and Polycyclic Aromatic Hydrocarbons (PAHs) Degradation and Naphthalene Metabolism in *Streptomyces* sp. (ERI-CPDA-1) Isolated from Oil Contaminated Soil. *Bioresour. Technol.* **2012**, *112*, 83–90. [CrossRef] [PubMed]
64. Mahajan, M.; Manek, D.; Vora, N.; Kothari, R.K.; Mootapally, C.; Nathani, N.M. Fungi with High Ability to Crunch Multiple Polycyclic Aromatic Hydrocarbons (PAHs) from the Pelagic Sediments of Gulfs of Gujarat. *Mar. Pollut. Bull.* **2021**, *167*, 112293. [CrossRef] [PubMed]
65. Peidro-Guzmán, H.; Pérez-Llano, Y.; González-Abradelo, D.; Fernández-López, M.G.; Dávila-Ramos, S.; Aranda, E.; Hernández, D.R.O.; García, A.O.; Lira-Ruan, V.; Pliego, O.R.; et al. Transcriptomic Analysis of Polyaromatic Hydrocarbon Degradation by the Halophilic Fungus *Aspergillus sydowii* at Hypersaline Conditions. *Environ. Microbiol.* **2021**, *23*, 3435–3459. [CrossRef]
66. Birolli, W.G.; De Santos, D.A.; Alvarenga, N.; Garcia, A.C.F.S.; Romão, L.P.C.; Porto, A.L.M. Biodegradation of Anthracene and Several PAHs by the Marine-Derived Fungus *Cladosporium* sp. CBMAI 1237. *Mar. Pollut. Bull.* **2018**, *129*, 525–533. [CrossRef]
67. Wu, Y.-R.; Luo, Z.-H.; Vrijmoed, L.L.P. Biodegradation of Anthracene and Benz[a]Anthracene by Two Fusarium Solani Strains Isolated from Mangrove Sediments. *Bioresour. Technol.* **2010**, *101*, 9666–9672. [CrossRef]
68. Medaura, M.C.; Guivernau, M.; Moreno-Ventas, X.; Prenafeta-Boldú, F.X.; Viñas, M. Bioaugmentation of Native Fungi, an Efficient Strategy for the Bioremediation of an Aged Industrially Polluted Soil With Heavy Hydrocarbons. *Front. Microbiol.* **2021**, *12*, 626436. [CrossRef]
69. Varjani, S.J. Microbial Degradation of Petroleum Hydrocarbons. *Bioresour. Technol.* **2017**, *223*, 277–286. [CrossRef]
70. Mrozik, A.; Piotrowska-Seget, Z. Bioaugmentation as a Strategy for Cleaning up of Soils Contaminated with Aromatic Compounds. *Microbiol. Res.* **2010**, *165*, 363–375. [CrossRef]
71. Oliver, J.D. Recent Findings on the Viable but Nonculturable State in Pathogenic Bacteria. *FEMS Microbiol. Rev.* **2010**, *34*, 415–425. [CrossRef] [PubMed]
72. Liu, Q.; Tang, J.; Gao, K.; Gurav, R.; Giesy, J.P. Aerobic Degradation of Crude Oil by Microorganisms in Soils from Four Geographic Regions of China. *Sci. Rep.* **2017**, *7*, 14856. [CrossRef] [PubMed]
73. Kothari, V.; Panchal, M.; Srivastava, N. Microbial Degradation of Hydrocarbons. *Microb. Degrad. Hydrocarb.* **2014**, 1–16.
74. Warczewska, Z.; Niewiadomska, A.; Grzyb, A. Wybrane metody bioremediacji in situ z wykorzystaniem mikroorganizmów. *Woda Sr. Obsz. Wiej.* **2018**, *26*, 65–78.
75. Barathi, S.; Vasudevan, N. Utilization of Petroleum Hydrocarbons by Pseudomonas Fluorescens Isolated from a Petroleum-Contaminated Soil. *Environ. Int.* **2001**, *26*, 413–416. [CrossRef] [PubMed]
76. Kamath, R.; Rentz, J.A.; Schnoor, J.L.; Alvarez, P.J.J. Chapter 16 Phytoremediation of Hydrocarbon-Contaminated Soils: Principles and Applications. In *Studies in Surface Science and Catalysis*; Elsevier: Amsterdam, The Netherlands, 2004; Volume 151, pp. 447–478, ISBN 9780444516992.
77. Wang, D.; Xie, Y.; Jaisi, D.P.; Jin, Y. Effects of Low-Molecular-Weight Organic Acids on the Dissolution of Hydroxyapatite Nanoparticles. *Environ. Sci. Nano* **2016**, *3*, 768–779. [CrossRef]
78. Khan, S.; Afzal, M.; Iqbal, S.; Khan, Q.M. Plant–Bacteria Partnerships for the Remediation of Hydrocarbon Contaminated Soils. *Chemosphere* **2013**, *90*, 1317–1332. [CrossRef]
79. Pawlik, M.; Piotrowska-Seget, Z. Endophytic Bacteria Associated with *Hieracium piloselloides*: Their Potential for Hydrocarbon-Utilizing and Plant Growth-Promotion. *J. Toxicol. Environ. Health Part A* **2015**, *78*, 860–870. [CrossRef]
80. Weyens, N.; Taghavi, S.; Barac, T.; Van Der Lelie, D.; Boulet, J.; Artois, T.; Carleer, R.; Vangronsveld, J. Bacteria Associated with Oak and Ash on a TCE-Contaminated Site: Characterization of Isolates with Potential to Avoid Evapotranspiration of TCE. *Environ. Sci. Pollut. Res.* **2009**, *16*, 830–843. [CrossRef]
81. Weyens, N.; Van Der Lelie, D.; Artois, T.; Smeets, K.; Taghavi, S.; Newman, L.; Carleer, R.; Vangronsveld, J. Bioaugmentation with Engineered Endophytic Bacteria Improves Contaminant Fate in Phytoremediation. *Environ. Sci. Technol.* **2009**, *43*, 9413–9418. [CrossRef]
82. Weyens, N.; Van Der Lelie, D.; Taghavi, S.; Newman, L.; Vangronsveld, J. Exploiting Plant–Microbe Partnerships to Improve Biomass Production and Remediation. *Trends Biotechnol.* **2009**, *27*, 591–598. [CrossRef] [PubMed]
83. Weyens, N.; Van Der Lelie, D.; Taghavi, S.; Vangronsveld, J. Phytoremediation: Plant–Endophyte Partnerships Take the Challenge. *Curr. Opin. Biotechnol.* **2009**, *20*, 248–254. [CrossRef] [PubMed]
84. Cooney, J.J.; Silver, S.A.; Beck, E.A. Factors Influencing Hydrocarbon Degradation in Three Freshwater Lakes. *Microb. Ecol.* **1985**, *11*, 127–137. [CrossRef]
85. Małachowska-Jutsz, A.; Janosz, W.; Rudek, J. Ochrona Środowiska. Toksyczność gleby zanieczyszczonej olejem silnikowym poddanej samooczyszczaniu oraz fitoremediacj. *Ochr. Sr.* **2012**, *34*, 15–20.
86. Ruley, J.A.; Tumuhairwe, J.B.; Amoding, A.; Opolot, E.; Oryem-Origa, H.; Basamba, T. Assessment of Plants for Phytoremediation of Hydrocarbon-Contaminated Soils in the Sudd Wetland of South Sudan. *Plant Soil Environ.* **2019**, *65*, 463–469. [CrossRef]
87. Rabêlo, F.H.S.; Vangronsveld, J.; Baker, A.J.M.; Van Der Ent, A.; Alleoni, L.R.F. Are Grasses Really Useful for the Phytoremediation of Potentially Toxic Trace Elements? A Review. *Front. Plant Sci.* **2021**, *12*, 778275. [CrossRef] [PubMed]

88. Xi, Y.; Song, Y.; Johnson, D.M.; Li, M.; Liu, H.; Huang, Y. Se Enhanced Phytoremediation of Diesel in Soil by Trifolium Repens. *Ecotoxicol. Environ. Saf.* **2018**, *154*, 137–144. [CrossRef]
89. Steliga, T.; Kluk, D. Assessment of the Suitability of Melilotus Officinalis for Phytoremediation of Soil Contaminated with Petroleum Hydrocarbons (TPH and PAH), Zn, Pb and Cd Based on Toxicological Tests. *Toxics* **2021**, *9*, 148. [CrossRef]
90. Steliga, T.; Kluk, D. Application of Festuca Arundinacea in Phytoremediation of Soils Contaminated with Pb, Ni, Cd and Petroleum Hydrocarbons. *Ecotoxicol. Environ. Saf.* **2020**, *194*, 110409. [CrossRef]
91. Peng, S.; Zhou, Q.; Cai, Z.; Zhang, Z. Phytoremediation of Petroleum Contaminated Soils by Mirabilis Jalapa L. in a Greenhouse Plot Experiment. *J. Hazard. Mater.* **2009**, *168*, 1490–1496. [CrossRef]
92. Li, L.; Zhu, P.; Wang, X.; Zhang, Z. Phytoremediation Effect of Medicago Sativa Colonized by Piriformospora Indica in the Phenanthrene and Cadmium Co-Contaminated Soil. *BMC Biotechnol.* **2020**, *20*, 20. [CrossRef] [PubMed]
93. Tang, J.; Wang, R.; Niu, X.; Zhou, Q. Enhancement of Soil Petroleum Remediation by Using a Combination of Ryegrass (*Lolium perenne*) and Different Microorganisms. *Soil Tillage Res.* **2010**, *110*, 87–93. [CrossRef]
94. Yousaf, S.; Andria, V.; Reichenauer, T.G.; Smalla, K.; Sessitsch, A. Phylogenetic and Functional Diversity of Alkane Degrading Bacteria Associated with Italian Ryegrass (*Lolium multiflorum*) and Birdsfoot Trefoil (*Lotus corniculatus*) in a Petroleum Oil-Contaminated Environment. *J. Hazard. Mater.* **2010**, *184*, 523–532. [CrossRef] [PubMed]
95. Pawlik, M.; Cania, B.; Thijs, S.; Vangronsveld, J.; Piotrowska-Seget, Z. Hydrocarbon Degradation Potential and Plant Growth-Promoting Activity of Culturable Endophytic Bacteria of Lotus Corniculatus and Oenothera Biennis from a Long-Term Polluted Site. *Environ. Sci. Pollut. Res.* **2017**, *24*, 19640–19652. [CrossRef]
96. Banks, M.K.; Kulakow, P.; Schwab, A.P.; Chen, Z.; Rathbone, K. Degradation of Crude Oil in the Rhizosphere of *Sorghum bicolor*. *Int. J. Phytoremediation* **2003**, *5*, 225–234. [CrossRef]
97. Baoune, H.; Aparicio, J.D.; Acuña, A.; El Hadj-khelil, A.O.; Sanchez, L.; Polti, M.A.; Alvarez, A. Effectiveness of the *Zea Mays*-Streptomyces Association for the Phytoremediation of Petroleum Hydrocarbons Impacted Soils. *Ecotoxicol. Environ. Saf.* **2019**, *184*, 109591. [CrossRef]
98. Liao, C.; Xu, W.; Lu, G.; Liang, X.; Guo, C.; Yang, C.; Dang, Z. Accumulation of Hydrocarbons by Maize (*Zea mays* L.) in Remediation of Soils Contaminated with Crude Oil. *Int. J. Phytoremediation* **2015**, *17*, 693–700. [CrossRef]
99. Anukwa, F.A.; Onuoha, E.M.; Nkang, A.; Nkereuwem, J. Phytoremediation Potential of *Zea mays* L. and *Panicum coloratum* L. on Hydrocarbon Polluted Soils. *Int. J. Bot.* **2020**, *17*, 1–7. [CrossRef]
100. Song, X.; Li, C.; Chen, W. Phytoremediation Potential of Bermuda Grass (*Cynodon dactylon* (L.) Pers.) in Soils Co-Contaminated with Polycyclic Aromatic Hydrocarbons and Cadmium. *Ecotoxicol. Environ. Saf.* **2022**, *234*, 113389. [CrossRef]
101. Gworek, B.; Klimczak, K.; Kijeńska, M. The Relation between Polyaromatic Hydrocarbon Concentration in Sewage Sludge and Its Uptake by Plants: Phragmites Communis, Polygonum Persicaria and Bidens Tripartita. *PLoS ONE* **2014**, *9*, e109548. [CrossRef]
102. Liu, R.; Jadeja, R.N.; Zhou, Q.; Liu, Z. Treatment and Remediation of Petroleum-Contaminated Soils Using Selective Ornamental Plants. *Environ. Eng. Sci.* **2012**, *29*, 494–501. [CrossRef] [PubMed]
103. Heidari, S.; Fotouhi Ghazvini, R.; Zavareh, M.; Kafi, M. Physiological Responses and Phytoremediation Ability of Eastern Coneflower (*Echinacea purpurea*) for Crude Oil Contaminated Soil. *Casp. J. Environ. Sci.* **2018**, *16*, 149–164. [CrossRef]
104. Kaimi, E.; Mukaidani, T.; Miyoshi, S.; Tamaki, M. Ryegrass Enhancement of Biodegradation in Diesel-Contaminated Soil. *Environ. Exp. Bot.* **2006**, *55*, 110–119. [CrossRef]
105. Shirdam, R.; Zand, A.D.; Bidhendi, G.N.; Mehrdadi, N. Phytoremediation of Hydrocarbon-Contaminated Soils with Emphasis on the Effect of Petroleum Hydrocarbons on the Growth of Plant Species. *Phytoprotection* **2009**, *89*, 21–29. [CrossRef]
106. Hall, J.; Soole, K.; Bentham, R. Hydrocarbon Phytoremediation in the Family Fabacea—A Review. *Int. J. Phytoremediation* **2011**, *13*, 317–332. [CrossRef]
107. Moore, F.P.; Barac, T.; Borremans, B.; Oeyen, L.; Vangronsveld, J.; Van Der Lelie, D.; Campbell, C.D.; Moore, E.R.B. Endophytic Bacterial Diversity in Poplar Trees Growing on a BTEX-Contaminated Site: The Characterisation of Isolates with Potential to Enhance Phytoremediation. *Syst. Appl. Microbiol.* **2006**, *29*, 539–556. [CrossRef]
108. Doty, S.L. Enhancing Phytoremediation through the Use of Transgenics and Endophytes. *New Phytol.* **2008**, *179*, 318–333. [CrossRef]
109. Wang, L.; Chen, S.; Yu, B. Poly-γ-Glutamic Acid: Recent Achievements, Diverse Applications and Future Perspectives. *Trends Food Sci. Technol.* **2022**, *119*, 1–12. [CrossRef]
110. Hara, T.; Ueda, S. Regulation of Poly Glutamate Production in *Bacillus subtilis* (*Natto*): Transformation of High PGA Productivity. *Agric. Biol. Chem.* **1982**, *46*, 2275–2281. [CrossRef]
111. Yanagibashi, T.; Kobayashi, M.; Omori, K. Application of Poly-γ-Glutamic Acid Flocculant to Flocculation–Sedimentation Treatment of Ultrafine Cement Suspension. *Water* **2019**, *11*, 1748. [CrossRef]
112. Li, M.; Zhu, X.; Yang, H.; Xie, X.; Zhu, Y.; Xu, G.; Hu, X.; Jin, Z.; Hu, Y.; Hai, Z.; et al. Treatment of Potato Starch Wastewater by Dual Natural Flocculants of Chitosan and Poly-Glutamic Acid. *J. Clean. Prod.* **2020**, *264*, 121641. [CrossRef]
113. Liang, J.; Shi, W. Poly-γ-Glutamic Acid Improves Water-Stable Aggregates, Nitrogen and Phosphorus Uptake Efficiency, Water-Fertilizer Productivity, and Economic Benefit in Barren Desertified Soils of Northwest China. *Agric. Water Manag.* **2021**, *245*, 106551. [CrossRef]
114. Wojtowicz, K.; Steliga, T.; Skalski, T. Badania Laboratoryjne Wpływu Dodatku γ-PGA Na Efektywność Biodegradacji Węglowodorów Ropopochodnych. *Nafta-Gaz* **2022**, *78*, 668–678. [CrossRef]

115. Zhao, G.; Sheng, Y.; Wang, C.; Yang, J.; Wang, Q.; Chen, L. In Situ Microbial Remediation of Crude Oil-Soaked Marine Sediments Using Zeolite Carrier with a Polymer Coating. *Mar. Pollut. Bull.* **2018**, *129*, 172–178. [CrossRef] [PubMed]
116. Luo, S.G.; Chen, S.C.; Cao, W.Z.; Lin, W.H.; Sheu, Y.T.; Kao, C.M. Application of γ-PGA as the Primary Carbon Source to Bioremediate a TCE-Polluted Aquifer: A Pilot-Scale Study. *Chemosphere* **2019**, *237*, 124449. [CrossRef]
117. Pang, X.; Lei, P.; Feng, X.; Xu, Z.; Xu, H.; Liu, K. Poly-γ-Glutamic Acid, a Bio-Chelator, Alleviates the Toxicity of Cd and Pb in the Soil and Promotes the Establishment of Healthy *Cucumis sativus* L. Seedling. *Environ. Sci. Pollut. Res.* **2018**, *25*, 19975–19988. [CrossRef]
118. Peng, Y.-P.; Chang, Y.-C.; Chen, K.-F.; Wang, C.-H. A Field Pilot-Scale Study on Heavy Metal-Contaminated Soil Washing by Using an Environmentally Friendly Agent—Poly-γ-Glutamic Acid (γ-PGA). *Environ. Sci. Pollut. Res.* **2020**, *27*, 34760–34769. [CrossRef] [PubMed]
119. Yang, Z.-H.; Dong, C.-D.; Chen, C.-W.; Sheu, Y.-T.; Kao, C.-M. Using Poly-Glutamic Acid as Soil-Washing Agent to Remediate Heavy Metal-Contaminated Soils. *Environ. Sci. Pollut. Res.* **2018**, *25*, 5231–5242. [CrossRef]
120. Wu, M.; Li, W.; Dick, W.A.; Ye, X.; Chen, K.; Kost, D.; Chen, L. Bioremediation of Hydrocarbon Degradation in a Petroleum-Contaminated Soil and Microbial Population and Activity Determination. *Chemosphere* **2017**, *169*, 124–130. [CrossRef]
121. Steliga, T.; Jakubowicz, P.; Kapusta, P. Changes in Toxicity during in Situ Bioremediation of Weathered Drill Wastes Contaminated with Petroleum Hydrocarbons. *Bioresour. Technol.* **2012**, *125*, 1–10. [CrossRef]
122. Nwankwegu, A.S.; Orji, M.U.; Onwosi, C.O. Studies on Organic and In-Organic Biostimulants in Bioremediation of Diesel-Contaminated Arable Soil. *Chemosphere* **2016**, *162*, 148–156. [CrossRef] [PubMed]
123. Abbasian, F.; Lockington, R.; Megharaj, M.; Naidu, R. A Review on the Genetics of Aliphatic and Aromatic Hydrocarbon Degradation. *Appl. Biochem. Biotechnol.* **2016**, *178*, 224–250. [CrossRef] [PubMed]
124. Tipu, M.I.; Ashraf, M.Y.; Sarwar, N.; Akhtar, M.; Shaheen, M.R.; Ali, S.; Damalas, C.A. Growth and Physiology of Maize (*Zea mays* L.) in a Nickel-Contaminated Soil and Phytoremediation Efficiency Using EDTA. *J. Plant Growth Regul.* **2021**, *40*, 774–786. [CrossRef]
125. Daryabeigi Zand, A.; Mühling, K.H. Phytoremediation Capability and Copper Uptake of Maize (*Zea mays* L.) in Copper Contaminated Soils. *Pollutants* **2022**, *2*, 53–65. [CrossRef]
126. Wu, L.; Li, Z.; Han, C.; Liu, L.; Teng, Y.; Sun, X.; Pan, C.; Huang, Y.; Luo, Y.; Christie, P. Phytoremediation of soil contaminated with cadmium, copper and polychlorinated biphenyls. *Int. J. Phytoremediation* **2012**, *14*, 570–584. [CrossRef]
127. Gobetti, S.; Kumor, A.; Prach, M.; Sinkkonen, A.; Pacwa-Płociniczak, M. The Multiway Support of Phytoremediation Efficiency of Soil That Has Been Historically Co-Contaminated with Petroleum Hydrocarbons and Heavy Metals Using *Zea mays*. *SSRN J.* **2022**. [CrossRef]
128. Ali, M.H.; Khan, M.I.; Bashir, S.; Azam, M.; Naveed, M.; Qadri, R.; Bashir, S.; Mehmood, F.; Shoukat, M.A.; Li, Y.; et al. Biochar and *Bacillus* sp. MN54 Assisted Phytoremediation of Diesel and Plant Growth Promotion of Maize in Hydrocarbons Contaminated Soil. *Agronomy* **2021**, *11*, 1795. [CrossRef]
129. Ijaz, A.; Imran, A.; Anwar Ul Haq, M.; Khan, Q.M.; Afzal, M. Phytoremediation: Recent Advances in Plant-Endophytic Synergistic Interactions. *Plant Soil* **2016**, *405*, 179–195. [CrossRef]
130. Zając, E.; Fabiańska, M.J.; Jędrszczyk, E.; Skalski, T. Hydrocarbon Degradation and Microbial Survival Improvement in Response to γ-Polyglutamic Acid Application. *Int. J. Environ. Res. Public Health* **2022**, *19*, 15066. [CrossRef]
131. Wrenn, B.A.; Venosa, A.D. Selective Enumeration of Aromatic and Aliphatic Hydrocarbon Degrading Bacteria by a Most-Probable-Number Procedure. *Can. J. Microbiol.* **1996**, *42*, 252–258. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

molecules

MDPI

Oil-contaminated soil remediation with the biodegradation by autochthonous microorganisms and phytoremediation by maize (*Zea mays*)

Katarzyna Wojtowicz [1]*, Teresa Steliga [1], Piotr Kapusta [1], Joanna Brzeszcz[1]

[1] Oil and Gas Institute – National Research Institute, 31-503 Krakow, ul. Lubicz 25 A, Poland
* Correspondence: wojtowiczk@inig.pl *

Table S1 Total content of selected groups of TPH and PAHs in soil G6 and after the biodegradation process with the use of biopreparation B1 (soil G6-3B1) and B2 (soil G6-3B2) , (repetition number n=7-10, p<0,05)

Hydrocarbons	Content ± SD (mg/kg dry mass soil)		
		After 90 Days	
	Initial Soil G6	Soil G6-3B1 soil G6 inoculated with biopreparation B1	Soil G6-3B2 soil G6 inoculated with biopreparation B2
TPH	4527,7 ± 316,94	3090,60 ± 216,34	2640,83 ± 184,94
Unidentified aliphatic hydrocarbons	2147,6± 150,33	1472,89 ± 103,10	1332,13 ± 93,33
$\sum nC_6–nC_9$	49,30 ± 4,93	19,79 ± 1,98	13,42 ± 1,34
$\sum nC_{10}–nC_{21}$	1163,60 ± 116,36	616,19 ± 61,62	455,35 ± 45,50
$\sum nC_{22}–nC_{30}$	831,30± 83,13	676,74 ±	557,34±
$\sum nC31–nC_{36}$	235,80± 23,58	211,41 ± 21,14	195,89± 19,59
isoprenoids	100,10 ± 10,01	94,07 ± 9,41	86,70 ± 8,67
∑PAHs	10,48 ± 1,05	7,61 ± 0,76	6,84 ± 0,68
∑two -ring PAHs	5,31 ± 0,53	3,48 ± 0,65	2,99 ± 0,03
∑ three - ring PAHs	2,23 ± 0,22	1,64 ± 0,15	1,50 ± 0,15
∑ four- ring PAHs	2,06 ± 0,21	1,71 ± 0,17	1,62 ± 0,16
∑ five- ring PAHs	0,66 ± 0,07	0,58 ± 0,06	0,55 ± 0,06
∑ six-ring PAHs	0,22 ±0,02	0,20 ± 0,02	0,19 ± 0,02

Table S2 Total content of selected groups of TPH and PAHs in soil after the phytoremediation process in the tested systems, (repetition number n=7-10, p<0,05)

Hydrocarbons	Content ± SD (mg/kg dry mass soil)		
	After 180 Days		
	System 1	System 2	System 3
TPH	2023,58± 141,65	1344,83± 94,14	915,16± 64,06
Unidentified aliphatic hydrocarbons	1043,21± 73,02	686,45± 48,05	523,97 ± 36,68
$\sum nC_6–nC_9$	7,11 ± 0,71	3,48 ± 0,35	1,82 ± 0,18
$\sum nC_{10}–nC_{21}$	284,98 ± 28,50	124,81 ± 12,48	48,59 ± 4,86
$\sum nC_{22}–nC_{30}$	448,64 ± 44,86	310,94 ± 31,07	166,71 ± 16,67
$\sum nC31–nC_{36}$	175,17 ± 17,52	147,96 ± 14,80	107,98 ± 10,80
isoprenoids	79,47 ± 7,95	71,19 ± 7,12	66,11 ± 6,61
∑PAHs	5,58 ± 0,56	4,05 ± 0,41	2,68 ± 0,27
∑two -ring PAHs	2,11 ± 0,21	1,36 ± 0,14	0,63 ± 0,06
∑ three - ring PAHs	1,29 ± 0,13	0,87 ± 0,09	0,55 ± 0,06
∑ four- ring PAHs	1,48 ± 0,15	1,23 ± 0,12	0,99 ± 0,01
∑ five- ring PAHs	0,52 ± 0,05	0,44 ± 0,04	0,38 ± 0,04
∑Six-ring PAHs	0,18 ± 0,02	0,15 ± 0,02	0,13 ± 0,01

Ecotoxicological analyses –description of the methodology

Phytotoxkit™ test (MicroBioTests Inc., Belgium)

Chronic toxicity assessment test *Phytotoxkit™* is based on the evaluation of germination and early growth of plants (root elongation inhibition measurement. Three types of plants selected according to germination rate and root growth rate are used in the test, which allows making a complete determination in 3 days of incubation: monocotyledonous - sorghum *(Sorghum saccharatum),* and dicotyledonous - cress *(Lepidium sativum)* and white mustard *(Sinapis alba).* The determination process was carried out using three repetitions for each test plant. The tests were performed on transparent polystyrene test plates. *Incubation conditions:* temperature T = 25°C in darkness, incubation time t = 72 h. *Test reaction:* inhibition of germination and early root growth.

The inhibition in seed germination and root growth was calculated according to a following equation:

$$I = \frac{(A - B)}{B} \cdot 100\%$$

where: ***I*** *– inhibition (%),* ***A*** *– seed germination or root length in control soil,* ***B*** *– seed germination or root length in test soil*

Ostracodtoxkit(F)™ (MicroBioTest Inc. Belgium)

Ostracodtoxkit(F) test belongs to direct contact tests of chronic toxicity estimation with the use of crustaceans *Heterocypris incongruens*. The direct contact test is done with the

application of young bottom crustaceans (*Heterocypris incongruens*), which hatched from cysts during 52 hours (according to the producer's procedure). The test is carried out on 6-hole polystyrene microplates. A standardised nutrient is an algae suspension, which is applied to the suspension in portions of 2 ml. Portions of 10 ostracods are added to each cell of the microplate. 0.5 g of a control soil is applied to the row A, whereas 0.5 g of the soil sample is added to the consecutive rows. Incubation lasts for 6 days in a temperature of 25°C. Results reading consists of estimation of life microforms amount in each hole of the microplate and measurement of their length. Growth inhibition of *H. incongruens* was calculated as

$$GI = 100 - \left(\frac{A}{B} \cdot 100\%\right)$$

where: A – ostracods length in tested soil, B – ostracods length in reference soil

Microtox® Solid Phase Test (SDI, USA)

Microtox Test created in the USA in 1979 as the first bioindication test, combines typical bioindication and analytic precision. Luminescence bacteria *Vibro fischeri*, which uses about 10% of its metabolism in order to emit light, was used as a bioindicator. In the presence of toxical substances, there is decrease in luminescence and increase in general toxicity of a sample. In an electron transportation system of the bacteria, lucipherasa enzyme (alcane oxygenaze) catalyses oxidation of a reduced substrate (reduced flavine mononucleotide, riboflavin phosphate or flavine adenin dinucleotide) and during this process luminescence, which can be measured with a photometer, takes place. The obtained substrates of this reaction are oxygen and long-chain aldehyde. In the presence of substances that have negative influence on cell metabolism, the decrease in luminescence of bacteria is immediate.

Test Microtox®, produced by SDI Company (USA), enables a direct contact of luminescence bacteria *Vibro fischeri* with a wastewater sample, which leads not only to determination of substances dissolved in water, but also to recognition of lipophilicity systems and poorly dissolved systems in water. Lyophilised bacteria *Vibro fischeri* can be stored in a period of a year in temperature of -20°C and then used for testing immediately after suspending in deionised water. A tests with serial dilutions of the wastewater were done according to a standard procedure and results were obtain in Delta Tox analyser. In this analysis, the concentration of wastewater, which reduces the luminescence by 50%, was determined (half maximal effective concentration, EC_{50}). In this presentation a lower of EC_{50} denotes a higher toxicity. To simplify this interpretation, EC_{50} values were converted to Toxicity Units (TU) in which a higher value denotes a higher toxicity according to the following formula.

$$TU = \frac{1}{EC_{50}} \cdot 100$$




Figure S1. Stand for testing the process of biodegradation of petroleum hydrocarbons.




Figure S2. Stand for testing the process of phytoremediation of soil G6-3B2

applied sciences

MDPI

Article

Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using *Echinacea purpurea*

Katarzyna Wojtowicz *, Teresa Steliga * and Piotr Kapusta

Oil and Gas Institute—National Research Institute, ul. Lubicz 25 A, 31-503 Krakow, Poland; piotr.kapusta@inig.pl
* Correspondence: wojtowiczk@inig.pl (K.W.); teresa.steliga@inig.pl (T.S.)

Featured Application: The results obtained during the study of phytoremediation assisted by bioaugmentation will in the future contribute to a better and faster process of cleaning areas contaminated with petroleum substances.

Abstract: Phytoremediation supported by bioaugmentation is a promising method considered for cleaning up areas polluted with petroleum hydrocarbons. In this study, phytoremediation was carried out using *Echinacea purpurea* as a phytoremediant on two types of soil: Soil DW—aged soil taken from an excavation pit, Soil OS—soil taken from an oil spill area. The tests for each soil were carried out in six test systems (non-inoculation, inoculation with the B1 microbial consortium, inoculation with the B2 microbial consortium, inoculation with the B1 microbial consortium with the addition of γ-PGA (γ-poly glutamic acid), inoculation with the B2 microbial consortium with the addition of γ-PGA and inoculation with the γ-PGA solution) for 6 months. The effectiveness of the remediation treatments used was assessed based on chromatographic analyses of soil and plant material (roots, shoots) and toxicological analyses using four types of toxicological tests (Phytotoxkit™ (MicroBioTests Inc., Gent, Belgium), Ostracodtoxkit™ (MicroBioTests Inc., Gent, Belgium), Microtox® Solid Phase Test (Modern Water Inc., New Castle, DE, USA), MARA (NCIMB Ltd., Aberdeen, UK)). The research conducted showed that the most effective method of bioremediation of soils contaminated with petroleum hydrocarbons was phytoremediation supported by bioaugmentation with the microbial consortium B2 with γ-PGA, which allowed for reducing the concentration of total petroleum hydrocarbons (TPHs) and polycyclic aromatic hydrocarbons (PAHs) in the tested soils by 53.98% and 49.54% (Soil DW-5) and 60.47% and 37.55% (Soil OS-5), respectively. In turn, the lowest bioremediation efficiency was recorded in non-inoculated systems, for which the concentration of TPHs and PAHs at the end of the study decreased by 18.40% and 16.14% (Soil DW-1) and 21.87% and 18.20% (Soil OS-1), respectively. The results of toxicological analyses confirmed the relationship between the concentration of TPHs and PAHs in the soil and its toxicity level.

Keywords: bioremediation; *Echinacea purpurea*; toxicological tests; microbial consortium (biopreparation); petroleum hydrocarbons; phytoextraction; phytoremediation

check for updates

Citation: Wojtowicz, K.; Steliga, T.; Kapusta, P. Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using *Echinacea purpurea*. *Appl. Sci.* **2023**, *13*, 13077. https://doi.org/10.3390/app132413077

Academic Editors: Fulvio Celico, Sara Remelli and Pietro Rizzo

Received: 30 October 2023
Revised: 6 December 2023
Accepted: 6 December 2023
Published: 7 December 2023

Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).

1. Introduction

The soil environment is prone to contamination by both organic and inorganic substances, which poses potential threats to the fauna and flora. Contamination of soils with petroleum hydrocarbons and heavy metals with carcinogenic, mutagenic and teratogenic properties seems particularly dangerous [1,2]. In view of the above, there is an urgent need to rehabilitate the areas contaminated with such xenobiotics and restore them [3]. One common method of land remediation is phytoremediation, which is a green chemistry technique, which uses plants capable of removing, metabolizing, assimilating and adsorbing harmful substances. Plants used in phytoremediation are classified as so-called hyperaccumulators. Hyperaccumulators are plants characterized by their ability to accumulate

large amounts of xenobiotics without the harm caused by phytotoxic effects [4–6]. For sites contaminated with petroleum hydrocarbons, plants belonging to the so-called naphthophytes are used in the phytoremediation process. Naphthophytes show an evolutionarily acquired ability to grow on oil lands [7–9]. Identifying potential plants, which can be used in the remediation of land contaminated with both organic and inorganic substances, is an interesting area of research into the phytoremediation process. One promising plant in this regard appears to be purple coneflower (*Echinacea purpurea*).

Echinacea purpurea, commonly known as purple coneflower, is a plant species from the Asteraceae family, originating from North America [10]. Adaptability to different climatic conditions, resistance to environmental stress and the ability to grow in different soil conditions have made *Echinacea purpurea* suitable for cultivation in most climate zones. In addition, *Echinacea purpurea* is characterized by high biomass production and a well-developed root system (fibrous root). The large surface area and high activity of the root system (production of root exudates) of purple coneflower allow for the stimulation of microbial activity in the root zone, thus enhancing mineralization at the root–soil interface, which enables rhizodegradation [11]. Among other mechanisms occurring during phytoremediation are biodegradation, phytoextraction, phytovolatilization, phytostabilization and abiotic loss, which can act individually or co-occur [12]. Phytoremediation studies conducted using *Echinacea purpurea* have shown that the plant has a high tolerance for petroleum and is suitable for remediating soils with petroleum concentrations exceeding 10,000 mg/kg [11]. In addition, purple coneflower has been proven to be effective in cleaning up soils polluted with PAHs and alkyl-PAHs [13–16]. A few studies also suggest that *Echinacea purpurea* can be used to clean up areas tainted with heavy metals [13,14]. However, it should be noted that when planning remediation strategies for land contaminated with various types of xenobiotics, the concentration, age and type of contamination should be taken into account. It has been proven that aged soils containing a greater variety of contaminants, with higher concentrations of xenobiotics, undergo the treatment process more slowly than freshly contaminated soils with a single type of contaminant. For real soils with a wide spectrum of contaminants, the efficiency of phytoremediation alone may be too low [17]. Therefore, it is critical to find a new, efficient and economically beneficial approach to the phytoremediation process.

In recent years, so-called combined methods using micro-organisms have become increasingly popular for remediating land contaminated with petroleum substances [18]. The most commonly used combination methods include phytoremediation in combination with bioaugmentation. It is based on the interaction of plants and microbes to increase the efficiency of pollutant degradation. The bioaugmentation-assisted phytoremediation method exploits the potential of micro-organisms present in the inoculant to biodegrade petroleum pollutants and their influence on the phytoremediation process. Increasing the concentration of bacteria, fungi and yeast in the root zone has been proven to promote root growth, allowing it to extract more nutrients from the soil, resulting in better growth. Moreover, the symbiotic interaction of micro-organisms with the root system allows for a better degradation of pollutants, transformation of complex organic molecules into simpler and less toxic compounds and their transfer to the plant [18–21]. It should be remembered that the inoculant used in bioaugmentation treatments should contain micro-organisms capable of degrading a broad spectrum of contaminants. The bacteria most commonly described in the literature for use in remediating land contaminated with petroleum hydrocarbons include *Achromobacter* sp. [22,23], *Bacillus* sp. [23–29], *Dietzia* sp. [30,31], *Gordonia* sp. [32,33], *Mycolicibacterium* sp. [34–39], *Pseudomonas* sp. [40–45], *Rhodococcus* sp. [46–51]. Among the fungi capable of metabolizing oil-derived substances are *Aspergillus* sp. [52,53], *Candida* sp. [54,55], *Cladosporium* sp. [56,57], *Fusarium* sp. [58,59] and *Penicillium* sp. [60,61]. It should also be remembered that in order to ensure the best possible efficiency of the bioaugmentation process, it is advisable to use inoculants developed on the basis of selected micro-organisms isolated from the natural environment. This avoids antagonistic interactions with naturally occurring micro-organisms in the tested

soil [17]. In addition, substances can be added to the inoculant to ensure better microbial growth by providing an easily assimilable carbon source or using a natural biosurfactant, such as γ-PGA [3,17,62,63].

Bioaugmentation and phytoremediation are technologies, which have been studied individually many times, but there are few publications describing their interaction in the bioremediation of soils polluted with TPHs and PAHs. It was assumed that the correct choice of inoculant in the bioaugmentation-assisted phytoremediation method would help ensure high efficiency of the treatment process of contaminated land. In the present study, it was decided to experimentally test the effectiveness of the combination of phytoremediation and bioaugmentation in several variants of conducting the process on real samples of land contaminated with TPHs and PAHs, and to evaluate the change in toxicity of soils after treatment.

2. Materials and Methods

2.1. Soil, Microbial Consortia, γ-PGA and Plant Description

Two types of soil contaminated with petroleum hydrocarbons were used in the study. Soils were taken from two locations, which were the G-70 pit area—which was used in the 1970s to store drilling waste mixed with soil (Soil DW)—and an oil spill site (Soil OS). Forty soil samples weighing 2 kg were collected from each site. Samples were taken from the surface layer (0–0.25 m below ground level) and the deep layer (0.25–1.5 m below ground level). Each type of soil taken was mixed and homogenized for averaging, followed by physical and chemical analyses. The composition of the soils used in the bioaugmentation-assisted phytoremediation study is shown in Table 1.

Table 1. Results of physical and chemical analyses of soils used in studies of bioaugmentation-assisted phytoremediation process.

Parameter		Soil DW	Soil OS
pH_{H2O}		7.06	6.99
Initial water moisture (%)		21.9	23.01
BTEX (benzene, toluene, p-, m-ksylen)		0.59	0.97
TPH		5176.03	4025.65
WWA		17.71	10.30
Cl^{-} [a]		168.5	258.6
$S—SO_4^{2-}$ [a]		781.5	826.3
$N—NH_4^{+}$ [a]		70.1	79.4
$N—NO_3^{-}$ [a]		241.2	230.4
$P—PO_4^{3-}$ [a]		27.26	32.1
Al_2O_3 [b]		688.5	426.9
SiO_2 [b]		236.5	315.6
Fe_2O_3 [b]		5.3	6.2
MgO [b]		3.9	3.0
CaO [b]		6.9	6.5
Sand (%)		54.01	75.21
Silt (%)		10.37	6.92
Clay (%)		35.62	17.87
Heavy metal content [a]	As	0.9	0.7
	Ba	28.9	23.5
	Cd	1.0	0.7
	Cr	11.1	9.1
	Co	3.1	3.0
	Cu	22.2	24.5
	Hg	0.5	0.2
	Pb	22.9	15.4
	Mo	1.8	2.1
	Sn	5.4	6.7
	Zn	20.6	17.1
	Ni	11.8	9.3

COD—Chemical oxygen demand, [a] Content expressed as ((mg kg^{-1} dry mass), [b] Content expressed as (g kg^{-1} dry mass).

Two microbial consortia developed from indigenous micro-organisms isolated from contaminated sites were used to carry out the bioaugmentation procedure. Microbial consortium B1 contained the bacterial strains *Dietzia* sp. IN118, *Gordonia* sp. IN101, *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN119, *Rhodococcus glopberulus* IN113 and *Raoultella* sp. IN109. Microbial consortium B2 consisted of bacteria present in microbial consortium B1 and was additionally enriched with fungal strains *Aspergillus sydowii*, *Aspergillus versicolor*, *Candida* sp., *Cladosporium halotolerans*, *Penicillium chrysogenum*. The density of cells in both consortia was in the range of 5×10^8–1×10^9 CFU/mL. The bacterial and fungal strains used in the microbial consortia had proven abilities to metabolize petroleum hydrocarbons and did not exhibit pathogenic properties. A detailed description of the design of microbial consortia B1 and B2, along with the method of identification of the strains of bacteria and fungi present in them, is provided in a previous article [17]. More information about the microbial consortia used in the research is available in Supplementary Materials. A commercially available formulation of Ambiogel® (Ambioteco, Staszów, Poland) containing 5% pure poly-γ-glutamic acid [64], whose impact on the efficiency of bioremediation has been confirmed in earlier studies, was used as an additive for microbial consortia. The phytoremediation process was carried out using seeds of *Echinacea purpurea* (Legutko, Poland), which were sown in each of the tested systems.

2.2. Course of the Experiment

Phytoremediation studies using *Echinacea purpurea* as a phytoremediator and bioaugmentation-assisted phytoremediation were conducted under laboratory conditions using the vase method. Experiments were conducted on two types of soil with known contaminant concentrations. Prior to soil treatment, the soil was fertilized with the mineral fertilizer "Azofoska" (Inco, Warszawa, Poland) composed of 13.6% N-total nitrogen; 5.5% N-nitrogen; 8.1% N-ammonium nitrogen; 6.4% P_2O_5 soluble in neutral ammonium citrate solution and water, 4.2% P_2O_5 soluble in water; 19.1%K_2O soluble in water; 4.5% MgO; 23.0% SO_3; 0.045% B; 0.18% Cu; 0.17% Fe; 0.27% Mn; 0.04% Mo; 0.045% Zn. This was to ensure that the C:N:P ratio in the tested soils was 100:10:1 [1,17,35], which is optimal for the proper development and activity of micro-organisms responsible for hydrocarbon degradation. The soil reaction was brought to a value of 7.5–7.6 through supplementation with fertilizer lime of natural origin (Biovita, Tenczynek, Poland) [65–67]. Soils prepared in this way were placed in 10 L pots, and *Echinacea purpurea* was sown in them. The process of treating contaminated soils was carried out in 12 systems. Depending on the system studied, the soil was bioaugmented with the appropriate inoculant. Systems 1 and 7—soils not inoculated, systems 2 and 8—soils inoculated with microbial consortium B1, systems 3 and 9—soils inoculated with microbial consortium B2, systems 4 and 10—soils inoculated with microbial consortium B1 with the addition of γ-PGA, systems 5 and 11—soils inoculated with microbial consortium B2 with the addition of γ-PGA, systems 6 and 12—soils inoculated with γ-PGA. The study was carried out in two series of repetitions in a greenhouse, ensuring constant conditions for the experiment, i.e., temperature of 22–25 °C, aeration, humidity (soil sprinkling) and sunlight (system of LED lighting lamps with a light intensity of 3000–5000 lumens ensuring the maintenance of the day and night cycle). The first series of studies was completed after 3 months, while the second series was completed after 6 months. This allowed us to control changes in the amount of TPHs and PAHs in soil and plant material during the treatment process carried out in the systems studied. A schematic of the test bed for bioaugmentation-assisted phytoremediation is shown in Figure 1.




Figure 1. Schematic of the stand for conducting bioaugmentation-assisted phytoremediation using *Echinacea purpurea* (laboratory conditions).

After the experiment, the test plants were removed from the pots in which phytoremediation tests were carried out and then washed with distilled water to remove any soil residues. Plants were separated into shoots and roots. The plant material was dried and homogenized, and its biomass was determined. After testing, the soil was dried to an air-dry state, ground and sieved through a 1 mm sieve. Soil samples and plant material prepared in this way were analyzed for TPH and PAH content. In addition, the soil was tested for changes in toxicity using commercially available toxicological tests.

2.2.1. Analysis of TPH in Soil and Plant Material

Extraction of TPH from the soil/plant material was carried out by sonication. For this purpose, the dried test material (5 g) was placed in a 200 mL Erlenmajer flask, followed by the addition of 20 mL of dichloromethane (Chempur, Piekary Śląskie, Poland). The analyte extraction process was carried out in three repetition series of 15 min each. The resulting extracts were filtered and then subjected to purification of polar substances by dSPE solid-phase extraction. Purification of the extracts was carried out on pre-conditioned Florisil Chromabond columns No. 730081 (Macharey-Nagel, Dueren, Germany). The purified extracts were concentrated on a rotary evaporator (Chemland, Piekary ŚląskiePoland) and subjected to chromatographic analysis using a Clarus 500 gas chromatograph from Perkin Elmer (Waltham, MA, USA), according to the procedure for determining TPH developed at the Department of Complex Fluids Exploitation Technology at INiG—PIB [65,66]. The most important operating parameters of the chromatograph are listed in Table 2.

Table 2. Operating parameters of chromatographs during the determination of TPHs and PAHs.

Chromatographic Analysis of Aliphatic Hydrocarbons—TPH, nC_6–nC_{44}, Pristane, Phytane (Clarus 500 GC Perkin Elmer) Parameter of Chromatography					
Temperature (°C)			Column	Detector	Calibration Standards
Injector	Carrier Gas	Temperature Program			
290 °C	He 20 mL min^{-1}	30 °C—isothermal run for 2 min 30–105 °C—temp. increase rate 10 °C min^{-1} 105–285 °C—temp. increase rate 5 °C min^{-1} 285 °C—isothermal run for 5 min^{-1}	RTX-1 30 m × 0.53 mm (Restek, Bellefonte, PA, USA)	FID 300 °C	Reference soil: BAM—K010 (Tusnovic Instruments, Poland) Standard mixture hydrocarbons (nC_6–nC_{44}) ASTM® No. D2807 (Supelco, Saint Louis, MO, USA) Fuel Oil Degradation Mix nC_{17}, pristane, nC_{18}, phytane No. A029668 (Restek, Bellefonte, PA, USA)
Chromatographic Analysis of Polycyclic Aromatic Hydrocarbons—PAH (Vanquish Core Thermo Scientific) Parameter of Chromatography					
Eluents	**Flow**	**Gradient**	**Column**	**Detector**	**Calibration Standards**
A—methanol B—acetonitrile (Chempur, Piekary śląskie, Poland)	1.5 mL min^{-1}	20% B—for 1.5 min 20–50% B—for 1.5 min 50–100% B—for 1 min 100% B—for 1 min 100–0% B—for 3 min 100% A—for 3 min	NUCLEODUR C_{18} PAH column 125 mm × 4 mm, 3 µm (Marcherey-Nagel, Dueren, Germany)	UV-ViS FLD	Reference soil: PAHs by HPLC No. SQC017-40G (Sigma-Aldrich, Saint Louis, MO, USA) Certified PAH-Mix solution No. 722393 (Marcherey-Nagel, Dueren, Germany)

2.2.2. Analysis of PAHs in Soil and Plant Material

Isolation of PAHs from soil/plant material was carried out using the QuEChERS method developed at the Department of Reservoir Fluid Exploitation Technology at INiG—PIB (oil–gas). The dried test material was placed in a 50 mL vial, followed by the addition of 5 mL of distilled water, 10 mL of acetonitrile (Chempur, Piekary Śląskie, Poland) and $MgSO_4$+ NaCl extraction mixture No. SST640 (Interchim, Montluçon, France). The vials were tightly sealed and shaken for 5 min. After this time, the samples were transferred to an MPW-223e centrifuge (MPW, Warszawa, Poland) and centrifuged at 3500 rpm for 10 min. The decanted extract was cleaned of interfering substances in special Clean-up Kit Fruit and Vegetables AOAC vials No. JO3937 (Interchim, Montluçon, France). Purified extracts were filtered through a syringe filter with a 0.2 µm nylon (PA) membrane and subjected to chromatographic analysis. PAH determination was performed on a Vanquish Core Series liquid chromatograph from Thermo Fisher Scientific (Waltham, MA, USA) [17,68]. The most important operating parameters of the chromatograph are listed in Table 2.

2.2.3. Soil Toxicity Analysis

The toxicity of soils at baseline and after completion of the bioaugmentation-assisted phytoremediation process carried out in the studied systems was evaluated using 4 commercially available microbiotests: Phytotoxkit™ (MicroBioTests Inc., Gent Belgium), Ostracodtoxkit™ (MicroBioTests Inc., Gent, Belgium), Microtox® Solid Phase Test (Modern Water Inc., New Castle, DE, USA) and MARA (NCIMB Ltd., Aberdeen, UK) [65]. Phytotoxkit™ and Ostracodtoxkit™ tests are among the chronic toxicity assessment tests. The Phytotoxkit™ test is based on evaluating the germination and early growth of test plants, which include *Sorghum saccharatum*, *Lepidium sativum* and *Sinapis alba*. The result of the test is inhibition of germination and early root growth after three days of incubation at 25 °C. The Ostracodtoxkit™ test is based on assessing the percentage mortality and growth inhibition of crustaceans of the species *Heterocypris incongruens* after 6 days of contact with the test soil. The Microtox® Solid Phase Test is an acute toxicity test and involves evaluating the decrease in fluorescence of *Vibrio fischeri* bacteria following con-

tact with the test sample. The result of the Microtox® Solid Phase test is EC_{50}, which is the concentration of the test sample, which produces 50% of the test reaction. To assess environmental risk, the MARA test was used, based on evaluating the degree of growth inhibition of 18 test organisms (*Microbacterium spaciec, Brevundimonas diminuta, Citrobacter freudii, Comamonas testosteroni, Entrococcus casseliflavus, Delftia acidovorans, Kurthia gibsoni, Staphylococcus warneri, Pseudomonas aurantiaca, Serriatia rudidaea* and *Pichia anomala*) after 18 h of incubation [17,62,65,69]. All toxicological tests were conducted in accordance with the manufacturers' procedures, while the results obtained were converted to TU values.

3. Results

3.1. Analysis of Petroleum Hydrocarbons in Soil, Root and Shoot

Chromatographic analyses of the soil for TPH content showed that bioaugmentation with microbial consortia developed on the basis of indigenous micro-organisms significantly affects the efficiency of its treatment process using *Echinacea purpurea* as a phytoremediator. At the end of the 6-month experiment, in non-inoculated systems, TPH biodegradation efficiency was low, at 18.40% (Soil DW-1) and 21.87% (Soil OS-1). The application of inoculation with bacterial microbial consortium B1 increased the biodegradation rate of TPH to 35.56% (Soil DW-2) and 40.85% (Soil OS-2). In systems inoculated with microbial consortium B2 enriched with fungal species capable of degrading petroleum hydrocarbons, TPH biodegradation rates were 44.16% (Soil DW-3) and 49.78% (Soil OS-3). In addition, it was observed that the use of γ-PGA significantly increases the efficiency of cleaning up soils polluted with petroleum substances. In soil taken from the site of an oil spill pit (Soil DW), the application of inoculation with microbial consortia B1 and B2 with the addition of γ-PGA in the phytoremediation process resulted in a reduction in TPH concentrations by 46.71% in system 4 and 53.98% in system 5. In addition, in the system inoculated with γ-PGA alone, phytoremediation efficiency using *Echinacea purpurea* was 3.4% higher than in the non-inoculated system. Similarly, in soil taken from an oil spill site (Soil OS), the phytoremediation efficiency was highest in systems inoculated with microbial consortia containing γ-PGA, achieving 51.19% (Soil OS-4) and 60.47% (Soil OS-5). Inoculation with pure γ-PGA compared to the non-inoculated system led to an increase of 5.02%. In addition, based on chromatographic analyses using two types of soil with different ages of contamination, it was found that aged soils (Soil DW) were less amenable to treatment than freshly contaminated soils (Soil OS). The changes in TPH concentrations in DW and OS soils subjected to bioaugmentation-assisted phytoremediation in the systems studied are shown in Figure 2.

For a complete evaluation of the effectiveness of the bioaugmentation-assisted soil phytoremediation process, the distribution of alkanes in the studied systems before and after the experiment was determined. It was determined that hydrocarbons with a carbon chain length of nC_{10}–nC_{21} were best removed from the soil matrix. Slightly worse values of biodegradation degrees were obtained for light hydrocarbons with carbon chain lengths of nC_6–nC_9. Heavy hydrocarbons with carbon chain lengths of nC_{22}–nC_{30} and nC_{31}–nC_{36} degraded at a much lower rate, consistent with previous studies. Moreover, it was observed that in soils vaccinated with a microbial consortium enriched with selected fungal species, a greater reduction in the content of heavy hydrocarbons was obtained than in systems inoculated with a bacterial biopreparation. A summary of the concentrations of selected groups of aliphatic hydrocarbons in DW and OS soils before and after phytoremediation studies in different inoculation systems is provided in Table 3.

Chromatographic analyses of the roots and shoots showed that aliphatic hydrocarbons can be extracted from the soil into *Echinacea purpurea* in all systems tested. However, it was observed that in systems not inoculated and inoculated with γ-PGA alone, the concentration of TPH in the roots and shoots was very low, which directly affected the efficiency of phytoremediation. In addition, poor growth of *Echinacea purpurea* was observed in systems 1 and 7, and 6 and 12 due to a poorly developed root system. Low TPH concentrations are also due to the fact that the diffusion of petroleum substances into the

plant is greatly impeded due to the hydrophobic nature of hydrocarbons and their large molecular weights. Therefore, the uptake of lipophilic substances by plants is positively correlated with the lipid content of the root epidermis. Bioaugmentation of the soil with microbial consortia B1 and B2 allowed for the concentration of micro-organisms to be increased in the root zone of the phytoremediant used, which had a beneficial effect on the development of roots, and thus, the growth of *Echinacea purpurea*. In addition, it is likely that soil bioaugmentation increased the plant's production of surface-active compounds, such as small-molecule organic acids or polypeptides, capable of lowering surface tension and breaking down or converting complex organic compounds into simpler compounds with greater bioavailability [8,9,70–74]. Therefore, higher concentrations of TPH were determined in the analyzed plant material taken from systems inoculated with B1 and B2 microbial consortia. Moreover, the use of γ-PGA as an additive to microbial consortia made it possible to provide the plant with a readily available source of carbon and energy, which further influenced biomass production. It is also worth noting that some literature sources report that γ-PGA can be considered a biosurfactant, which affects the diffusion of xenobiotics into the root [17,62,64,75]. Chromatographic analyses of the roots and shoots showed that extraction of aliphatic hydrocarbons from the soil to the root occurred most readily in systems 4 and 5, and 10 and 11. It should also be noted that in all systems tested, higher concentrations of TPH were obtained in the roots of *Echinacea purpurea* than in the shoots. A summary of TPH concentrations in the roots and shoots of *Echinacea purpurea* as a function of time and the system tested is provided in Figures 3 and 4.




Figure 2. TPH content in soils after 3 and 6 months of phytoremediation process aided by bioaugmentation in the tested systems (System 1—soil DW + *E. purpurea*, System 2—soil DW + *E. purpurea* + B1 microbial consortium, System 3—soil DW + *E. purpurea* + B2 microbial consortium, System 4—soil DW + *E. purpurea* + B1 microbial consortium with the addition of γ-PGA, System 5—soil DW + *E. purpurea* + B2 microbial consortium with the addition of γ-PGA, System 6—soil DW + *E. purpurea* + γ-PGA, System 7—soil OS + *E. purpurea*, System 8—soil OS + *E. purpurea* + B1 microbial consortium, System 9—soil OS + *E. purpurea* + B2 microbial consortium, System 10—soil OS + *E. purpurea* + B1 microbial consortium with the addition of γ-PGA, System 11—soil OS + *E. purpurea* + B2 microbial consortium with the addition of γ-PGA, System 12—soil OS + *E. purpurea* + γ-PGA).

Table 3. Contents of selected groups of aliphatic hydrocarbons in the studied soils (Soil DW and Soil OS) before and after treatment with phytoremediation assisted by inoculation.

Parameter	Content ± SD (mg/kg Dry Mass Soil)						
	Initial Soil DW	After 180 Days					
		Soil DW-1	Soil DW-2	Soil DW-3	Soil DW-4	Soil DW-5	Soil DW-6
TPH	5176.03	4223.45	3335.33	2890.1	2758.11	2381.96	4047.68
$\sum nC_6–nC_9$	26.31	20.36	7.58	3.44	2.52	1.98	18.75
$\sum nC_{10}–nC_{21}$	892.48	683.89	407.46	300.37	271.65	196.37	627.57
$\sum nC_{22}–nC_{30}$	1092.38	919.85	845.72	845.72	777.35	661.4	896.15
$\sum nC31–nC_{36}$	383.94	347.88	333.86	316.11	317.99	290.3	342.39
isoprenoids	169.93	159.37	147.65	142.73	141.63	135.23	156.52
Unidentified aliphatic hydrocarbons	2610.98	2092.1	1593.07	1347.24	1246.98	1096.69	2006.31
Parameter	**Initial Soil OS**	**After 180 Days**					
		Soil OS-1	**Soil OS-2**	**Soil OS-3**	**Soil OS-4**	**Soil OS-5**	**Soil OS-6**
TPH	4025.65	3145.08	2381.17	2021.52	1965.03	1591.26	2943.24
$\sum nC_6–nC_9$	4.09	2.99	0.8	0.46	0.39	0.28	2.75
$\sum nC_{10}–nC_{21}$	1013.77	735.64	422.35	352.87	258.55	191.87	658.24
$\sum nC_{22}–nC_{30}$	358.69	290.79	263.5	263.5	231.71	193.99	285.91
$\sum nC31–nC_{36}$	200.59	179.49	172.95	153.23	164.76	144.65	176.54
isoprenoids	384.97	356.53	325.83	319.22	309.93	300.32	348.62
Unidentified aliphatic hydrocarbons	2063.54	1579.64	1195.74	963.57	999.69	760.15	1471.18




Figure 3. TPH content of *Echinacea purpurea* roots after 3 and 6 months of phytoremediation process aided by bioaugmentation in the tested systems.

Chromatographic analyses of the soils for polycyclic aromatic hydrocarbons confirmed the observations made during TPH analyses. The PAH concentration in systems not inoculated and inoculated with γ-PGA alone was the lowest, at 16.14% in Soil DW-1, 14.19% in Soil DW-6, 16.28% in Soil OS-1 and 18.20% in Soil OS-6. Phytoremediation assisted by bioaugmentation with B1 and B2 microbial consortia reduced PAH concentrations in the tested soils by 30.82% (Soil DW-2), 36.73% (Soil DW-3), 27.10% (Soil OS-2) and 31.07% (Soil OS-3). The use of γ-PGA as an additive to microbial consortia enabled an additional reduction in PAH concentrations in the tested soils, with biodegradation rates of 39.22% (Soil DW-4), 49.54% (Soil DW-5), 32.89% (Soil OS-4) and 37.55% (Soil OS-5). The changes in ΣPAH concentrations in SD and OS soils subjected to bioaugmentation-assisted phytoremediation are shown in Figure 5.




Figure 4. TPH content of *Echinacea purpurea* shoots after 3 and 6 months of phytoremediation process aided by bioaugmentation in the tested systems.




Figure 5. PAH content in soil after 3 and 6 months of bioaugmentation-assisted phytoremediation process in the studied systems.

The tested soils significantly differed in the qualitative and quantitative composition of individual groups of polycyclic aromatic hydrocarbons. Soil taken from the pit area (Soil DW) had higher concentrations of naphthalene relative to the other PAHs. In contrast, in the soil taken from the oil spill site, the distribution of the different groups of PAHs was more even, with a slight predominance of three- and four-ring PAHs. In such case, it is extremely important to study the distribution of individual groups of polycyclic aromatic hydrocarbons, taking into account the number of hydrocarbon rings. Indeed, the efficiency of biodegradation and phytoremediation depends on the number of rings in the hydrocarbon molecule [17]. Qualitative and quantitative chromatographic analyses of PAHs showed that naphthalene was most readily degraded in both tested soils. Lower

values of biodegradation coefficients were recorded for three-ring PAHs. Four-ring PAHs were definitely less biodegradable. Six-ring PAHs had by far the worst distribution. It can therefore be assumed that during phytoremediation, bioaugmentation-assisted PAHs are degraded in the following order: PAH-2-ring → PAH-3-ring → PAH-4-ring → PAH-5-ring → PAH-6-ring [17,62,76]. Analyzing the qualitative and quantitative composition of the treated soils, it was observed that despite the higher degrees of biodegradation of the sum content of polycyclic aromatic hydrocarbons in SD-1-6 soils compared to the similarly inoculated OS-1-6 soils, the degrees of biodegradation of individual PAH groups were higher in OS-1 soils. This is influenced by the qualitative and quantitative composition of PAHs and the age of the pollutants. Considering the method of bioaugmentation-assisted phytoremediation, it can be seen that the use of fungi in the microbial consortia and the addition of γ-PGA positively affect the biodegradation efficiency of PAHs in soils. A summary of the concentrations of selected groups of polycyclic aromatic hydrocarbons in DW and OS soils before and after phytoremediation studies in different inoculation systems is provided in Table 4.

Table 4. Contents of selected groups of PAHs in the studied soils (Soil DW and Soil OS) before and after treatment with phytoremediation assisted by inoculation.

Parameter	Content ± SD (mg/kg Dry Mass Soil)						
	Initial Soil DW	After 180 Days					
		Soil DW-1	Soil DW-2	Soil DW-3	Soil DW-4	Soil DW-5	Soil DW-6
∑PAH	17.71	14.85	12.25	11.204	10.76	8.95	14.19
∑two-ring PAHs	5.985	4.667	3.309	2.795	2.646	2	4.361
∑ three-ring PAHs	4.815	3.961	3.216	2.813	2.825	1.639	3.763
∑ four-ring PAHs	5.196	4.655	4.263	4.16	3.918	3.502	4.529
∑ five-ring PAHs	1.278	1.168	1.087	1.065	1.017	0.952	1.143
∑ six-ring PAHs	0.434	0.4	0.377	0.37	0.358	0.34	0.392
Parameter	**Initial Soil OS**	**After 180 Days**					
		Soil OS-1	**Soil OS-2**	**Soil OS-3**	**Soil OS-4**	**Soil OS-5**	**Soil OS-6**
∑PAHs	10.3	8.626	7.51	7.1	6.92	6.34	8.43
∑two-ring PAHs	0.495	0.335	0.195	0.153	0.15	0.129	0.316
∑ three-ring PAHs	1.325	0.979	0.728	0.612	0.639	0.381	0.938
∑ four-ring PAHs	5.026	4.275	3.808	3.704	3.508	2.468	4.187
∑ five-ring PAHs	2.322	2.03	1.85	1.736	1.727	1.651	1.995
∑ six-ring PAHs	1.136	1.008	0.93	0.898	0.891	0.859	0.993

Chromatographic analyses of *Echinacea purpurea* roots and shoots in the tested systems showed insignificant concentrations of the analyzed xenobiotics in plant tissues. As with TPH, the highest PAH concentrations were determined in plants growing in systems inoculated with microbial consortium B1 and B2 with γ-PGA. Slightly lower PAH values were determined in the roots and stems of *Echinacea purpurea* taken from systems inoculated with microbial consortium only. In plants taken from non-inoculated systems and those inoculated with γ-PGA alone, the PAH concentrations were trace. Regardless of the system tested, higher amounts of PAHs were determined in root tissues than in shoot tissues. A summary of PAH concentrations in the roots and shoots of *Echinacea purpurea* as a function of time and the system tested is provided in Figures 6 and 7.




Figure 6. PAH content of *Echinacea purpurea* roots after 3 and 6 months of phytoremediation process aided by bioaugmentation in the tested systems.




Figure 7. PAH content of *Echinacea purpurea* shoots after 3 and 6 months of phytoremediation process aided by bioaugmentation in the tested systems.

3.2. *Toxicological Analysis*

Toxicological tests carried out on initial soil samples and after remediation treatments in the studied systems allowed for the assessment of the impact of changes in the concentrations of petroleum substances on soil biocenosis. Each of the toxicological tests performed showed the greatest decrease in soil toxicity in systems in which phytoremediation (*Echinacea purpurea*) was carried out supported by bioaugmentation with the B2 microbial consortium with the addition of γ-PGA. Phytoremediation supported by vaccina-

tion with the B1 microbial consortium with γ-PGA turned out to be slightly less effective. By far the smallest decrease in soil toxicity was observed in non-inoculated systems and in systems inoculated with γ-PGA alone. The obtained results of toxicological tests were consistent with the results of chromatographic analyses, which confirms the assumption that soil toxicity largely depends on the amount of TPHs and PAHs in the soil. A summary of the results of all toxicological tests performed for Soil DW and Soil OS subjected to the bioremediation process in the tested systems is presented in Figures 8 and 9.




Figure 8. Summary of the results of toxicological tests of drilling waste soil (Soil DW) subjected to the purification process in the analyzed systems ($n = 3$, $p < 0.05$).




Figure 9. Summary of the results of toxicological tests of oil spill soil (Soil OS) subjected to the purification process in the analyzed systems ($n = 3$, $p < 0.05$).

Manufacturer-level toxicity tests carried out using the Phytotoxkit test took into account two toxicity assessment criteria: inhibition of root growth and inhibition of germi-

nation. Taking into account the criterion of root growth inhibition, the toxicity expressed in the TU toxicity unit was determined for Soil DW and Soil OS initial soils at the levels of 12.2 and 9.8 (*Lapidum sativum*), 10.5 and 8.5 (*Sorgum saccharatum*), and 12.3 and 9.2 (*Sinapis alba*), respectively. The degrees of toxicity of Soil DW after bioremediation in the tested systems ranged from 10.5 to 4 (*Lapidum sativum*), 9.8 to 3.6 (*Sorgum saccharatum*) and from 10.3 to 3.3 (*Sinapis alba*). The toxicity of Soil OS after 6 months of purification, depending on the tested system, was in the range of 9.8–3.7 (*Lapidum sativum*), 7.8–3.3 (*Sorgum saccharatum*) and 9.5–2.9 (*Sinapis alba*). For the second parameter tested (germination inhibition), the TU values calculated for the tested plants in the initial soil samples were in the range of 9.3–7.6 (Soil DW) and 6.8–5.3 (Soil OS). As a result of the remediation procedures, a reduction in the degree of soil toxicity was achieved, with values ranging from 8.8 to 2.6 (Soil DW) and from 6.8 to 2.6 (Soil OS).

Soil toxicity at the consumer level was determined on the basis of the Ostracodtoxkit toxicological test, which showed that the use of phytoremediation supported by bioaugmentation with the B2 microbial consortium with γ-PGA caused the greatest decrease in the toxicity of soils polluted with petroleum hydrocarbons compared to *Heterocypris incongruens*. In the soil taken from the excavated pit area (Soil DW), the TU value decreased from 11.2 to 4.2 (Soil DW-5), and in the soil from the oil spill area (Soil OS), the TU value decreased from 10.9 to 3.6. The values of soil toxicity degrees after completion of phytoremediation supported by bioaugmentation with microbial consortium B1 with γ-PGA were 5.2 (Soil DW-4) and 4.2 (Soil OS-4). The highest toxicity values after remediation treatments were recorded in non-inoculated soils, at 10.2 (Soil DW-1) and 9.9 (Soil OS-1).

The use of *Vibrio fischeri* luminescent bacteria in the Microtox test made it possible to determine the toxicity of the tested soils at the level of decomposers. The values of the toxicity degrees of the initial Soil DW and Soil OS soils were 14.1 and 11.4, respectively. The remediation procedures carried out made it possible to reduce the toxicity levels of Soil DW to 12.3 (Soil DW-1), 8.5 (Soil DW-2), 6 (Soil DW-3), 5.4 (Soil DW-4), 4.4 (Soil DW-5) and 11.5 (Soil DW-6). In turn, the toxicity of Soil OS after 6 months of remediation treatments decreased to the levels of 10.5 (Soil OS-1), 7.8 (Soil OS-2), 5.4 (Soil OS-3), 4.6 (Soil OS -4), 3.9 (Soil OS-5) and 10.9 (Soil OS-6).

In addition, the MARA environmental risk assessment test was performed for each soil, which used 11 test strains with different sensitivity to xenobiotics present in the soil. The toxicities calculated based on the mean toxic concentration (MTC), expressed in toxicity units, were 14.3 (Soil DW) and 11.8 (Soil OS). As a result of the remediation treatments, the toxicity of soil DW decreased from 12.2 (in non-inoculated Soil DW-1) to 4.0 (in Soil DW-5). However, the values of Soil OS toxicity degrees after completion of remediation treatments ranged from 10.8 (Soil OS-1) to 3.4 (Soil OS-5).

4. Discussion

Phytoremediation is now a widely used method for remediating land contaminated with petroleum hydrocarbons, but for aged drilling waste or soils heavily contaminated with various types of xenobiotics, its effectiveness is usually too low. Therefore, so-called combined methods using micro-organisms are increasingly being used [18]. The study showed that the effectiveness of phytoremediation using *Echinacea purpurea* as a test plant largely depends on the concentration, type and age of the contaminant, as well as the conditions under which the treatment process is carried out [17]. The chromatographic analyses performed proved that the use of phytoremediation supported by inoculation with microbial consortia produced on the basis of indigenous micro-organisms allows for a significant reduction in the amount of TPHs and PAHs in the studied soils. This is probably due to the fact that the effectiveness of the soil remediation process using plants is affected by the number of bacteria, fungi and yeasts in the surrounding root zone [77]. Thus, the introduction of micro-organisms isolated from contaminated sites (biopreparation) into the soil makes it possible to improve the efficiency of contaminant elimination while avoiding antagonistic interactions of the soil's indigenous microflora with foreign

microbial cultures [9,78,79]. Moreover, inoculating the soil with the microbial consortia makes it possible to increase the concentration of micro-organisms near the roots, which influences their better growth [80]. The penetration of roots into the deeper layers of the rhizosphere causes the soil to expand, resulting in improved ventilation and permeability of the test soil, as well as increased interaction between micro-organisms and contamination. On the other hand, the plant's roots secrete organic compounds, such as organic acids, sugars, amino acids, etc., which provide a source of carbon and energy for the growth of rhizosphere micro-organisms. It should also be remembered that dead root cells are treated by soil micro-organisms as an excellent source of nutrients, which further increases their metabolic activity [9,81–83]. Therefore, the synergistic interactions occurring among the micro-organisms present in the *Echinacea purpurea* root inoculant allow for an improvement in bioavailability to pollutants, thereby increasing their degradation efficiency.

Based on the chromatographic analyses of the soils, roots and shoots, it can be concluded that during phytoremediation studies using *Echinacea purpurea*, the petroleum-based contaminants underwent mechanisms of degradation, extraction, inhibition or a combination thereof [12,18]. Chromatographic analyses of the soils showed a slight reduction in the concentration of TPHs and PAHs in non-inoculated systems, while inoculation with microbial consortia and microbial consortia with γ-PGA allowed for a significant increase in the treatment efficiency of the studied soils. This proves the validity of the phytoremediation method assisted by microbial consortium inoculation. It should also be noted that among the inoculants tested, the B2 microbial consortium with γ-PGA was the most effective. This confirms the hypothesis that the use of an inoculant, which contains in its composition micro-organisms from different species (bacteria, fungi) and additives promoting the degradation of xenobiotics (γ-PGA), allows for an improvement in the efficiency of phytoremediation of soils, as well as biodegradation of oil pollutants by micro-organisms. Purification of the soils from toxic substances occurs due to the ability of the micro-organisms present in microbial consortia, natural soil microflora and *Echinacea purpurea* to produce enzymes and other chemicals, which break down or convert organic pollutants into simpler compounds with lower toxicity and better assimilation by plants [81,82]. Enzymes such as nitroreductase, dehalogenase, lactase, peroxidase and others are responsible for the degradation of petroleum hydrocarbons [84]. Moreover, the decrease in the concentration of petroleum hydrocarbons in the soil may be due to the transfer of contaminants from the soil to the plant [85–87]. This assumption is confirmed by the results of chromatographic analyses of the root and aboveground parts of *Echinacea purpurea* (stem + leaves). The experiments conducted showed the presence of petroleum hydrocarbons (TPHs and PAHs) in the roots and shoots of purple coneflower after the experiments. It is likely that during the treatment process of the studied soils, some of the analyzed contaminants were adsorbed on the root surface [20,88], and some were accumulated in the above-surface part of the plant via phytoextraction [18,87,89]. It should also be noted that some plants—for the proper conduct of the process of photosynthesis—use the ability to retain hydrocarbons in the tissues of the roots. This helps provide the energy needed for survival and is considered a self-defense strategy [87]. In addition, some contaminants can undergo phytoevaporation, which involves moving the contaminants into a volatile state [12,90]. It is also extremely important that γ-PGA was used as an additive to the microbial consortia in the study. Systems inoculated with γ-PGA-enhanced microbial consortia showed the highest % biodegradation of TPHs and PAHs, as well as the highest biomass production. γ-PGA provides an easily digestible source of carbon and energy for micro-organisms and *Echinacea purpurea*, which significantly affects the growth of the plant and the development of root zone micro-organisms, providing greater efficiency in the treatment of contaminated land.

Analyzing the effectiveness of the phytoremediation processes carried out in the systems studied, one cannot but mention the effect of the concentration and age of the contaminant on the efficiency of the treatment process. Remediation of aged soils (drilling waste) occurs much more slowly than soils contaminated by oil spills. This is related to

the stronger adsorption of contaminants on the surface and in the porous space of aged soil [17,91]. Moreover, the study showed that the process of cleaning the soil from the tested contaminants proceeded at different rates depending on the type and dose of the toxicant. It was found that in soil containing a higher diversity of contaminants, the cleaning process was slower compared to soil from an oil spill site.

Toxicological tests carried out using bioindicators belonging to three taxonomic groups (bacteria, crustaceans, plants) confirmed the effectiveness of ongoing phytoremediation of soils contaminated with petroleum hydrocarbons in systems supported by inoculation with microbial consortia and microbial consortia with γ-PGA.

5. Conclusions

The purpose of this study was to evaluate the effectiveness of a phytoremediation method for soils contaminated with petroleum substances (TPHs and PAHs) supported by bioaugmentation with microbial consortia and microbial consortia with the addition of γ-PGA. For this study, we used *Echinacea purpurea*, characterized by an extensive fibrous root system, high tolerance for various soil conditions and environmental stresses, and proven ability to phytoremediate hydrocarbon-contaminated soils. The tests were conducted under laboratory conditions to ensure that the conditions were the same for all systems tested. On the basis of chromatographic analyses of the soils, roots and shoots, it was proved that combined methods (phytoremediation assisted by bioaugmentation) of remediation of land contaminated with petroleum hydrocarbons allow using the potential of both the process of biodegradation of contaminants by micro-organisms present in the inoculant and phytoremediation of land with the help of plant organisms. This makes it possible to significantly increase the efficiency of the remediation process of contaminated land and restore its usable properties, as confirmed by the results of the toxicological tests conducted.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/app132413077/s1, Construction of B1 and B2 microbial consortia, Table S1: Identification of bacterial strains used in B1 and B2 microbial consortia.

Author Contributions: Conceptualization, K.W. and T.S.; software, K.W.; validation, K.W. and P.K.; formal analysis, K.W., T.S. and P.K.; investigation, K.W., T.S. and P.K.; writing—original draft preparation, K.W. and T.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was financially supported by the Polish Ministry of Education and Science within statutory funding for Oil and Gas Institute—National Research Institute.

Institutional Review Board Statement: The studies used commercial toxicological tests not subject to ethical evaluation.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author. The data are not publicly available due to privacy.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Steliga, T. *Bioremediacja Odpadów Wiertniczych Zanieczyszczonych Substancjami Ropopochodnymi ze Starych Dołów Urobkowych*; Instytut Nafty i Gazu: Krakow, Poland, 2009.
2. Kluk, D.; Steliga, T. Efektywna Metoda Identyfikacji Zanieczyszczeń Ropopochodnych (TPH) i Wielopierścieniowych Węglowodorów Aromatycznych (WWA) w Glebach. *Nafta-Gaz* **2017**, *73*, 488–495. [CrossRef]
3. Wojtowicz, K.; Steliga, T.; Skalski, T. Badania Laboratoryjne Wpływu Dodatku γ-PGA Na Efektywność Biodegradacji Węglowodorów Ropopochodnych. *Nafta-Gaz* **2022**, *78*, 668–678. [CrossRef]
4. Tang, K.H.D. Phytoremediation: Where Do We Go from Here? *Biocatal. Agric. Biotechnol.* **2023**, *50*, 102721. [CrossRef]
5. Rai, P.K.; Kim, K.-H.; Lee, S.S.; Lee, J.-H. Molecular Mechanisms in Phytoremediation of Environmental Contaminants and Prospects of Engineered Transgenic Plants/Microbes. *Sci. Total Environ.* **2020**, *705*, 135858. [CrossRef] [PubMed]
6. Liu, J.; Xin, X.; Zhou, Q. Phytoremediation of Contaminated Soils Using Ornamental Plants. *Environ. Rev.* **2018**, *26*, 43–54. [CrossRef]

7. Małachowska-Jutsz, A.; Janosz, W.; Rudek, J. Toksyczność gleby zanieczyszczonej olejem silnikowym poddanej samooczyszczaniu oraz fitoremediacji. *Ochr. Sr.* **2012**, *34*, 15–20.
8. Khan, S.; Afzal, M.; Iqbal, S.; Khan, Q.M. Plant–Bacteria Partnerships for the Remediation of Hydrocarbon Contaminated Soils. *Chemosphere* **2013**, *90*, 1317–1332. [CrossRef]
9. Pawlik, M.; Piotrowska-Seget, Z. Endophytic Bacteria Associated with *Hieracium piloselloides*: Their Potential for Hydrocarbon-Utilizing and Plant Growth-Promotion. *J. Toxicol. Environ. Health Part A* **2015**, *78*, 860–870. [CrossRef]
10. Saeidnia, S.; Manayi, A.; Vazirian, M. *Echinacea purpurea*: Pharmacology, Phytochemistry and Analysis Methods. *Pharmacogn. Rev.* **2015**, *9*, 63. [CrossRef]
11. Heidari, S.; Fotouhi Ghazvini, R.; Zavareh, M.; Kafi, M. Physiological Responses and Phytoremediation Ability of Eastern Coneflower (*Echinacea purpurea*) for Crude Oil Contaminated Soil. *CJES* **2018**, *16*, 149–164. [CrossRef]
12. Kubińska, N. Phytoremediation as an Approach to Clean up Contaminated Soil, Including Petroleum Product Contamination. *Nafta-Gaz* **2020**, *76*, 322–330. [CrossRef]
13. Pretorius, T.R.; Charest, C.; Kimpe, L.E.; Blais, J.M. The Accumulation of Metals, PAHs and Alkyl PAHs in the Roots of *Echinacea purpurea*. *PLoS ONE* **2018**, *13*, e0208325. [CrossRef]
14. Hou, L.; Liu, R.; Li, N.; Dai, Y.; Yan, J. Study on the Efficiency of Phytoremediation of Soils Heavily Polluted with PAHs in Petroleum-Contaminated Sites by Microorganism. *Environ. Sci. Pollut. Res.* **2019**, *26*, 31401–31413. [CrossRef]
15. Sun, C.; Shen, X.; Zhang, Y.; Song, T.; Xu, L.; Xiao, J. Molecular Defensive Mechanism of *Echinacea purpurea* (L.) Moench against PAH Contaminations. *Int. J. Mol. Sci.* **2023**, *24*, 11020. [CrossRef]
16. Sun, C.; Xiao, J.; Bai, L.; Bai, J.; Liu, J.; Geng, L.; Zhang, Y. Defined and Natural PAH Contaminations Shift PAH-Degrading Bacterial Community in Rhizosphere of Ornamental Plant Species *Echinacea purpurea* L. *Environ. Technol. Innov.* **2023**, *31*, 103189. [CrossRef]
17. Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J. Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (*Zea mays*). *Molecules* **2023**, *28*, 6104. [CrossRef] [PubMed]
18. Sui, X.; Wang, X.; Li, Y.; Ji, H. Remediation of Petroleum-Contaminated Soils with Microbial and Microbial Combined Methods: Advances, Mechanisms, and Challenges. *Sustainability* **2021**, *13*, 9267. [CrossRef]
19. Ptaszek, N.; Pacwa-Płociniczak, M.; Noszczyńska, M.; Płociniczak, T. Comparative Study on Multiway Enhanced Bio- and Phytoremediation of Aged Petroleum-Contaminated Soil. *Agronomy* **2020**, *10*, 947. [CrossRef]
20. Abdullah, S.R.S.; Al-Baldawi, I.A.; Almansoory, A.F.; Purwanti, I.F.; Al-Sbani, N.H.; Sharuddin, S.S.N. Plant-Assisted Remediation of Hydrocarbons in Water and Soil: Application, Mechanisms, Challenges and Opportunities. *Chemosphere* **2020**, *247*, 125932. [CrossRef]
21. Martínez-Rabelo, F.; Gómez-Guzmán, L.A.; Garcia-Segura, D.R.; Villegas-García, E.; Rodriguez-Campos, J.; Velázquez-Fernández, J.B.; Hernández-Castellanos, B.; Barois, I.; Contreras-Ramos, S.M. Hydrocarbon Bioremediation in a Pilot-Scale: A Combination of Bioaugmentation, Phytoremediation, and Vermiremediation. *Environ. Technol. Innov.* **2023**, *31*, 103210. [CrossRef]
22. Cutright, T.J.; Lee, S. Remediation of PAH-Contaminated Soil Using *Achromobacter* sp. *Energy Sources* **1994**, *16*, 279–287. [CrossRef]
23. Hong, Y.-H.; Ye, C.-C.; Zhou, Q.-Z.; Wu, X.-Y.; Yuan, J.-P.; Peng, J.; Deng, H.; Wang, J.-H. Genome Sequencing Reveals the Potential of *Achromobacter* sp. HZ01 for Bioremediation. *Front. Microbiol.* **2017**, *8*, 1507. [CrossRef]
24. Cerqueira, V.S.; Hollenbach, E.B.; Maboni, F.; Vainstein, M.H.; Camargo, F.A.O.; Peralba, M.D.C.R.; Bento, F.M. Biodegradation Potential of Oily Sludge by Pure and Mixed Bacterial Cultures. *Bioresour. Technol.* **2011**, *102*, 11003–11010. [CrossRef]
25. Janbandhu, A.; Fulekar, M.H. Biodegradation of Phenanthrene Using Adapted Microbial Consortium Isolated from Petrochemical Contaminated Environment. *J. Hazard. Mater.* **2011**, *187*, 333–340. [CrossRef] [PubMed]
26. Zheng, X.; Oba, B.T.; Shen, C.; Rong, L.; Zhang, B.; Huang, L.; Feng, L.; Liu, J.; Du, T.; Deng, Y. Effect of the Bacterial Community Assembly Process on the Microbial Remediation of Petroleum Hydrocarbon-Contaminated Soil. *Front. Microbiol.* **2023**, *14*, 1196610. [CrossRef] [PubMed]
27. Wang, D.; Lin, J.; Lin, J.; Wang, W.; Li, S. Biodegradation of Petroleum Hydrocarbons by Bacillus Subtilis BL-27, a Strain with Weak Hydrophobicity. *Molecules* **2019**, *24*, 3021. [CrossRef] [PubMed]
28. Baburam, C.; Mitema, A.; Tsekoa, T.; Feto, N.A. Bacillus Species and Their Invaluable Roles in Petroleum Hydrocarbon Bioremediation. In *Bacilli in Agrobiotechnology*; Islam, M.T., Rahman, M., Pandey, P., Eds.; Springer International Publishing: Cham, Switzerland, 2022; pp. 101–126. ISBN 9783030854645.
29. Rong, L.; Zheng, X.; Oba, B.T.; Shen, C.; Wang, X.; Wang, H.; Luo, Q.; Sun, L. Activating Soil Microbial Community Using Bacillus and Rhamnolipid to Remediate TPH Contaminated Soil. *Chemosphere* **2021**, *275*, 130062. [CrossRef]
30. Wang, X.-B.; Chi, C.-Q.; Nie, Y.; Tang, Y.-Q.; Tan, Y.; Wu, G.; Wu, X.-L. Degradation of Petroleum Hydrocarbons (C6–C40) and Crude Oil by a Novel Dietzia Strain. *Bioresour. Technol.* **2011**, *102*, 7755–7761. [CrossRef] [PubMed]
31. Chen, W.; Li, J.; Sun, X.; Min, J.; Hu, X. High Efficiency Degradation of Alkanes and Crude Oil by a Salt-Tolerant Bacterium Dietzia Species CN-3. *Int. Biodeterior. Biodegrad.* **2017**, *118*, 110–118. [CrossRef]
32. Liu, Y.; Wu, J.; Liu, Y.; Wu, X. Biological Process of Alkane Degradation by *Gordonia sihwaniensis*. *ACS Omega* **2022**, *7*, 55–63. [CrossRef]
33. Kim, H.; Dong, K.; Kim, J.; Lee, S. Characteristics of Crude Oil-degrading Bacteria *Gordonia iterans* Isolated from Marine Coastal in Taean Sediment. *MicrobiologyOpen* **2019**, *8*, e00754. [CrossRef] [PubMed]

34. Deng, Y.; Mou, T.; Wang, J.; Su, J.; Yan, Y.; Zhang, Y.-Q. Characterization of Three Rapidly Growing Novel *Mycobacterium* Species with Significant Polycyclic Aromatic Hydrocarbon Bioremediation Potential. *Front. Microbiol.* **2023**, *14*, 1225746. [CrossRef] [PubMed]
35. Brzeszcz, J.; Steliga, T.; Kapusta, P.; Turkiewicz, A.; Kaszycki, P. R-Strategist versus K-Strategist for the Application in Bioremediation of Hydrocarbon-Contaminated Soils. *Int. Biodeterior. Biodegrad.* **2016**, *106*, 41–52. [CrossRef]
36. Zhang, L.; Zhang, C.; Cheng, Z.; Yao, Y.; Chen, J. Biodegradation of Benzene, Toluene, Ethylbenzene, and o-Xylene by the Bacterium Mycobacterium Cosmeticum Byf-4. *Chemosphere* **2013**, *90*, 1340–1347. [CrossRef] [PubMed]
37. Rehmann, K.; Hertkorn, N.; Kettrup, A.A. Fluoranthene Metabolism in *Mycobacterium* sp. Strain KR20: Identity of Pathway Intermediates during Degradation and Growth. *Microbiology* **2001**, *147*, 2783–2794. [CrossRef] [PubMed]
38. Das, S.; Pettersson, B.M.F.; Behra, P.R.K.; Ramesh, M.; Dasgupta, S.; Bhattacharya, A.; Kirsebom, L.A. Characterization of Three *Mycobacterium* spp. with Potential Use in Bioremediation by Genome Sequencing and Comparative Genomics. *Genome Biol. Evol.* **2015**, *7*, 1871–1886. [CrossRef]
39. Solano-Serena, F.; Marchal, R.; Casarégola, S.; Vasnier, C.; Lebeault, J.-M.; Vandecasteele, J.-P. A Mycobacterium Strain with Extended Capacities for Degradation of Gasoline Hydrocarbons. *Appl. Environ. Microbiol.* **2000**, *66*, 2392–2399. [CrossRef]
40. Karamalidis, A.K.; Evangelou, A.C.; Karabika, E.; Koukkou, A.I.; Drainas, C.; Voudrias, E.A. Laboratory Scale Bioremediation of Petroleum-Contaminated Soil by Indigenous Microorganisms and Added *Pseudomonas aeruginosa* Strain Spet. *Bioresour. Technol.* **2010**, *101*, 6545–6552. [CrossRef]
41. Zhang, Z.; Hou, Z.; Yang, C.; Ma, C.; Tao, F.; Xu, P. Degradation of N-Alkanes and Polycyclic Aromatic Hydrocarbons in Petroleum by a Newly Isolated *Pseudomonas aeruginosa* DQ8. *Bioresour. Technol.* **2011**, *102*, 4111–4116. [CrossRef]
42. Bhattacharya, D.; Sarma, P.M.; Krishnan, S.; Mishra, S.; Lal, B. Evaluation of Genetic Diversity among *Pseudomonas citronellolis* Strains Isolated from Oily Sludge-Contaminated Sites. *Appl. Environ. Microbiol.* **2003**, *69*, 1435–1441. [CrossRef]
43. Ghosh, I.; Jasmine, J.; Mukherji, S. Biodegradation of Pyrene by a *Pseudomonas aeruginosa* Strain RS1 Isolated from Refinery Sludge. *Bioresour. Technol.* **2014**, *166*, 548–558. [CrossRef] [PubMed]
44. Palleroni, N.J.; Pieper, D.H.; Moore, E.R.B. Microbiology of Hydrocarbon-Degrading Pseudomonas. In *Handbook of Hydrocarbon and Lipid Microbiology*; Timmis, K.N., Ed.; Springer: Berlin/Heidelberg, Germany, 2010; pp. 1787–1798. ISBN 9783540775843.
45. Varjani, S.; Upasani, V.N.; Pandey, A. Bioremediation of Oily Sludge Polluted Soil Employing a Novel Strain of *Pseudomonas aeruginosa* and Phytotoxicity of Petroleum Hydrocarbons for Seed Germination. *Sci. Total Environ.* **2020**, *737*, 139766. [CrossRef]
46. Nazari, M.T.; Simon, V.; Machado, B.S.; Crestani, L.; Marchezi, G.; Concolato, G.; Ferrari, V.; Colla, L.M.; Piccin, J.S. *Rhodococcus*: A Promising Genus of Actinomycetes for the Bioremediation of Organic and Inorganic Contaminants. *J. Environ. Manag.* **2022**, *323*, 116220. [CrossRef] [PubMed]
47. Wei, Y.; Chen, J.; Wang, Y.; Meng, T.; Li, M. Bioremediation of the Petroleum Contaminated Desert Steppe Soil with *Rhodococcus erythropolis* KB1 and Its Effect on the Bacterial Communities of the Soils. *Geomicrobiol. J.* **2021**, *38*, 842–849. [CrossRef]
48. Pham, V.H.T.; Chaudhary, D.K.; Jeong, S.-W.; Kim, J. Van Hong Thi Pham; Chaudhary, D.K.; Jeong, S.-W.; Kim, J. Oil-Degrading Properties of a Psychrotolerant Bacterial Strain, *Rhodococcus* sp. Y2-2, in Liquid and Soil Media. *World J. Microbiol. Biotechnol.* **2018**, *34*, 33. [CrossRef]
49. Pacwa-Płociniczak, M.; Czapla, J.; Płociniczak, T.; Piotrowska-Seget, Z. The Effect of Bioaugmentation of Petroleum-Contaminated Soil with *Rhodococcus erythropolis* Strains on Removal of Petroleum from Soil. *Ecotoxicol. Environ. Saf.* **2019**, *169*, 615–622. [CrossRef]
50. Chen, X.; Shan, G.; Shen, J.; Zhang, F.; Liu, Y.; Cui, C. In Situ Bioremediation of Petroleum Hydrocarbon–Contaminated Soil: Isolation and Application of a *Rhodococcus* Strain. *Int. Microbiol.* **2022**, *26*, 411–421. [CrossRef]
51. Płociniczak, T.; Fic, E.; Pacwa-Płociniczak, M.; Pawlik, M.; Piotrowska-Seget, Z. Improvement of Phytoremediation of an Aged Petroleum Hydrocarbon-Contaminated Soil by *Rhodococcus erythropolis* CD 106 Strain. *Int. J. Phytoremediation* **2017**, *19*, 614–620. [CrossRef]
52. Mukherjee, A. Role of Aspergillus in Bioremediation Process. In *New and Future Developments in Microbial Biotechnology and Bioengineering*; Elsevier: Amsterdam, The Netherlands, 2016; pp. 209–214. ISBN 9780444635051.
53. Peidro-Guzmán, H.; Pérez-Llano, Y.; González-Abradelo, D.; Fernández-López, M.G.; Dávila-Ramos, S.; Aranda, E.; Hernández, D.R.O.; García, A.O.; Lira-Ruan, V.; Pliego, O.R.; et al. Transcriptomic Analysis of Polyaromatic Hydrocarbon Degradation by the Halophilic Fungus *Aspergillus sydowii* at Hypersaline Conditions. *Environ. Microbiol.* **2021**, *23*, 3435–3459. [CrossRef]
54. Li, J.; Xu, Y.; Song, Q.; Zhang, S.; Xie, L.; Yang, J. Transmembrane Transport Mechanism of N-Hexadecane by *Candida tropicalis*: Kinetic Study and Proteomic Analysis. *Ecotoxicol. Environ. Saf.* **2021**, *209*, 111789. [CrossRef]
55. Rana, S.; Handa, S.; Aggarwal, Y.; Puri, S.; Chatterjee, M. Role of *Candida* in the Bioremediation of Pollutants: A Review. *Lett. Appl. Microbiol.* **2023**, *76*, ovad103. [CrossRef] [PubMed]
56. Birolli, W.G.; Santos, D.A.; Alvarenga, N.; Garcia, A.C.F.S.; Romão, L.P.C.; Porto, A.L.M. Biodegradation of Anthracene and Several PAHs by the Marine-Derived Fungus *Cladosporium* sp. CBMAI 1237. *Mar. Pollut. Bull.* **2018**, *129*, 525–533. [CrossRef] [PubMed]
57. Bakri, M. Assessing Some Cladosporium Species in the Biodegradation of Petroleum Hydrocarbon for Treating Oil Contamination. *J. Appl. Microbiol.* **2022**, *133*, 3296–3306. [CrossRef] [PubMed]
58. Romauld, S.I.; Venkataraghavan, R.; Yuvaraj, D.; Devi, V.I.; Hashika, S. Mycoremediation of Hydrocarbon and Its Products Using *Fusarium oxysporum*. *Res. J. Pharm. Technol.* **2019**, *12*, 4216. [CrossRef]

59. Wu, Y.-R.; Luo, Z.-H.; Vrijmoed, L.L.P. Biodegradation of Anthracene and Benz[a]Anthracene by Two Fusarium Solani Strains Isolated from Mangrove Sediments. *Bioresour. Technol.* **2010**, *101*, 9666–9672. [CrossRef] [PubMed]
60. Zehra, A.; Dubey, M.K.; Meena, M.; Aamir, M.; Patel, C.B.; Upadhyay, R.S. Role of Penicillium Species in Bioremediation Processes. In *New and Future Developments in Microbial Biotechnology and Bioengineering*; Elsevier: Amsterdam, The Netherlands, 2018; pp. 247–260. ISBN 9780444635013.
61. Vanishree, M.; Thatheyus, A.J.; Ramya, D. Biodegradation of Petrol Using the Fungus *Penicillium* sp. *Sci. Int.* **2014**, *2*, 26–31. [CrossRef]
62. Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J.; Skalski, T. Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil. *Materials* **2022**, *15*, 400. [CrossRef] [PubMed]
63. Liang, J.; Shi, W. Poly-γ-Glutamic Acid Improves Water-Stable Aggregates, Nitrogen and Phosphorus Uptake Efficiency, Water-Fertilizer Productivity, and Economic Benefit in Barren Desertified Soils of Northwest China. *Agric. Water Manag.* **2021**, *245*, 106551. [CrossRef]
64. Zając, E.; Fabiańska, M.J.; Jedrszczyk, E.; Skalski, T. Hydrocarbon Degradation and Microbial Survival Improvement in Response to γ-Polyglutamic Acid Application. *Int. J. Environ. Res. Public Health* **2022**, *19*, 15066. [CrossRef]
65. Steliga, T.; Wojtowicz, K.; Kapusta, P.; Brzeszcz, J. Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture. *Molecules* **2020**, *25*, 709. [CrossRef]
66. Steliga, T.; Jakubowicz, P.; Kapusta, P. Changes in Toxicity during in Situ Bioremediation of Weathered Drill Wastes Contaminated with Petroleum Hydrocarbons. *Bioresour. Technol.* **2012**, *125*, 1–10. [CrossRef]
67. Nwankwegu, A.S.; Orji, M.U.; Onwosi, C.O. Studies on Organic and In-Organic Biostimulants in Bioremediation of Diesel-Contaminated Arable Soil. *Chemosphere* **2016**, *162*, 148–156. [CrossRef] [PubMed]
68. Wojtowicz, K. Opracowanie Metodyki Oznaczania WWA w Próbkach Gleb z Wykorzystaniem Chromatografii Cieczowej HPLC. *Nafte-Gaz* **2022**, *78*, 141–153. [CrossRef]
69. Steliga, T.; Jakubowicz, P.; Wojtowicz, K.; Kluk, D. Zastosowanie Testów Toksykologicznych w Przemyśle Naftowym. *Nafte-Gaz* **2018**, *74*, 684–689. [CrossRef]
70. Kamath, R.; Rentz, J.A.; Schnoor, J.L.; Alvarez, P.J.J. Chapter 16 Phytoremediation of Hydrocarbon-Contaminated Soils: Principles and Applications. In *Studies in Surface Science and Catalysis*; Elsevier: Amsterdam, The Netherlands, 2004; Volume 151, pp. 447–478. ISBN 9780444516992.
71. Wang, D.; Xie, Y.; Jaisi, D.P.; Jin, Y. Effects of Low-Molecular-Weight Organic Acids on the Dissolution of Hydroxyapatite Nanoparticles. *Environ. Sci. Nano* **2016**, *3*, 768–779. [CrossRef]
72. Weyens, N.; Taghavi, S.; Barac, T.; Van Der Lelie, D.; Boulet, J.; Artois, T.; Carleer, R.; Vangronsveld, J. Bacteria Associated with Oak and Ash on a TCE-Contaminated Site: Characterization of Isolates with Potential to Avoid Evapotranspiration of TCE. *Environ. Sci. Pollut. Res.* **2009**, *16*, 830–843. [CrossRef] [PubMed]
73. Weyens, N.; Van Der Lelie, D.; Artois, T.; Smeets, K.; Taghavi, S.; Newman, L.; Carleer, R.; Vangronsveld, J. Bioaugmentation with Engineered Endophytic Bacteria Improves Contaminant Fate in Phytoremediation. *Environ. Sci. Technol.* **2009**, *43*, 9413–9418. [CrossRef] [PubMed]
74. Weyens, N.; Van Der Lelie, D.; Taghavi, S.; Vangronsveld, J. Phytoremediation: Plant–Endophyte Partnerships Take the Challenge. *Curr. Opin. Biotechnol.* **2009**, *20*, 248–254. [CrossRef]
75. Li, D.; Hou, L.; Gao, Y.; Tian, Z.; Fan, B.; Wang, F.; Li, S. Recent Advances in Microbial Synthesis of Poly-γ-Glutamic Acid: A Review. *Foods* **2022**, *11*, 739. [CrossRef]
76. Abbasian, F.; Lockington, R.; Megharaj, M.; Naidu, R. A Review on the Genetics of Aliphatic and Aromatic Hydrocarbon Degradation. *Appl. Biochem. Biotechnol.* **2016**, *178*, 224–250. [CrossRef]
77. Rehman, K.; Imran, A.; Amin, I.; Afzal, M. Inoculation with Bacteria in Floating Treatment Wetlands Positively Modulates the Phytoremediation of Oil Field Wastewater. *J. Hazard. Mater.* **2018**, *349*, 242–251. [CrossRef] [PubMed]
78. Mani, D.; Kumar, C.; Patel, N.K.; Sivakumar, D. Enhanced Clean-up of Lead-Contaminated Alluvial Soil through *Chrysanthemum indicum* L. *Int. J. Environ. Sci. Technol.* **2015**, *12*, 1211–1222. [CrossRef]
79. Ojuederie, O.; Babalola, O. Microbial and Plant-Assisted Bioremediation of Heavy Metal Polluted Environments: A Review. *Int. J. Environ. Res. Public Health* **2017**, *14*, 1504. [CrossRef]
80. Kadam, S.K.; Watharkar, A.D.; Chandanshive, V.V.; Khandare, R.V.; Jeon, B.-H.; Jadhav, J.P.; Govindwar, S.P. Co-Planted Floating Phyto-Bed along with Microbial Fuel Cell for Enhanced Textile Effluent Treatment. *J. Clean. Prod.* **2018**, *203*, 788–798. [CrossRef]
81. Weyens, N.; Van Der Lelie, D.; Taghavi, S.; Newman, L.; Vangronsveld, J. Exploiting Plant–Microbe Partnerships to Improve Biomass Production and Remediation. *Trends Biotechnol.* **2009**, *27*, 591–598. [CrossRef] [PubMed]
82. Afzal, M.; Yousaf, S.; Reichenauer, T.G.; Sessitsch, A. The Inoculation Method Affects Colonization and Performance of Bacterial Inoculant Strains in the Phytoremediation of Soil Contaminated with Diesel Oil. *Int. J. Phytoremediation* **2012**, *14*, 35–47. [CrossRef]
83. Xu, X.; Liu, W.; Tian, S.; Wang, W.; Qi, Q.; Jiang, P.; Gao, X.; Li, F.; Li, H.; Yu, H. Petroleum Hydrocarbon-Degrading Bacteria for the Remediation of Oil Pollution Under Aerobic Conditions: A Perspective Analysis. *Front. Microbiol.* **2018**, *9*, 2885. [CrossRef]
84. Cherian, S.; Oliveira, M.M. Transgenic Plants in Phytoremediation: Recent Advances and New Possibilities. *Environ. Sci. Technol.* **2005**, *39*, 9377–9390. [CrossRef]

85. Lee, J.H. An Overview of Phytoremediation as a Potentially Promising Technology for Environmental Pollution Control. *Biotechnol Bioproc E* **2013**, *18*, 431–439. [CrossRef]
86. Mani, D.; Kumar, C. Biotechnological Advances in Bioremediation of Heavy Metals Contaminated Ecosystems: An Overview with Special Reference to Phytoremediation. *Int. J. Environ. Sci. Technol.* **2014**, *11*, 843–872. [CrossRef]
87. Kiamarsi, Z.; Kafi, M.; Soleimani, M.; Nezami, A.; Lutts, S. Conjunction of Vetiveria Zizanioides L. and Oil-Degrading Bacteria as a Promising Technique for Remediation of Crude Oil-Contaminated Soils. *J. Clean. Prod.* **2020**, *253*, 119719. [CrossRef]
88. Rohrbacher, F.; St-Arnaud, M. Root Exudation: The Ecological Driver of Hydrocarbon Rhizoremediation. *Agronomy* **2016**, *6*, 19. [CrossRef]
89. Aliku, C.B.; Madu, C.N.; Aliku, O. Accelerating Phytoextraction of Petroleum Hydrocarbon with Organic Stimulant. *MethodsX* **2021**, *8*, 101509. [CrossRef] [PubMed]
90. Fernández Rodríguez, M.D.; García Gómez, M.C.; Alonso Blazquez, N.; Tarazona, J.V. Soil Pollution Remediation. In *Encyclopedia of Toxicology*; Elsevier: Amsterdam, The Netherlands, 2014; pp. 344–355. ISBN 9780123864550.
91. Helmy, Q.; Laksmono, R.; Kardena, E. Bioremediation of Aged Petroleum Oil Contaminated Soil: From Laboratory Scale to Full Scale Application. *Procedia Chem.* **2015**, *14*, 326–333. [CrossRef]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

applied sciences

MDPI

Supplementary Materials

Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using *Echinacea purpurea*

Katarzyna Wojtowicz *, Teresa Steliga * and Piotr Kapusta

Oil and Gas Institute—National Research Institute, ul. Lubicz 25 A, 31-503 Krakow, Poland; piotr.kapusta@inig.pl
* Correspondence: wojtowiczk@inig.pl (K.W.); teresa.steliga@inig.pl (T.S.)

Construction of B1 and B2 microbial consortia

The bacterial and fungal strains used in study came from the hydrocarbon-degrading microbial collection of the Department of Microbiology (at the Oil and Gas Institute—National Research Institute, Poland). Strain diagnostic features were determined based on microscopic observations, morphology, growth on the selective agar media and biochemical profile (API tests, bioMerieux). The hydrocarbon-metabolising capabilities of all microbial strains were examined based on growth on the mineral medium supplemented with the compounds tested, as described in our previous work and according to the methods of Wrenn and Venosa. Each bacterial strain was grown individually in a nutrient broth supplemented with sodium acetate (0.2%, *w*/*v*), while each fungal strain was grown in a Czapek–Dox broth (both BD Difco™) and incubated at room temperature with shaking at 150 rpm for 24–72 h to obtain density $5*10^8$–$1*10^9$ cfu/mL. The biopreparations used in this study were constructed by mixing equal volumes of each strain. Then, the obtained mixed culture was kept in a nutrient broth supplemented with sodium acetate (0.2%, *w*/*v*) and sucrose and occasionally examined for the presence of all introduced strains. Table S1 presents the putative genes involved in petroleum hydrocarbon degradation found in genomes of the closest relatives of all bacterial strains used in this study

Table S1 Identification of bacterial strains used in B1 and B2 microbial consortia

Strain	The Closest Relative Based on 16S rRNA Accession Number (% of Identity) *	The Closest Relative for which the Genome Sequence is Available in NCBI GenBank, Accession Number (% of Identity) *	Putative Gene Encoding for the Enzymes Catalysing the Breakdown of Hydrocarbons
Dietzia sp. IN118	Dietzia sp. LJ3 MG049763 99.93%	Dietzia kunjamensis 313 CP099712 99.78%	Alkane 1-monooxygenase, aromatic ring-hydroxylating dioxygenase subunit alpha
Gordonia sp.IN101	Gordonia sp. Tm-B24 MT533993 99.63%	Gordonia terrae RL-JC02 CP049836 99.63%	Alkane 1-monooxygenase, pentachlorophenol monooxygenase, naphthalene 1,2-dioxygenase subunit alpha (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Mycolicibacterium frederiksbergense IN53	Mycolicibacterium frederiksbergense DSM 44346 (typical strain) NR_025393.1 99.58%	Mycolicibacterium frederiksbergense LB 501T 99.58%	Alkane 1-monooxygenase (2 copies), pentachlorophenol monooxygenase, naphthalene 1,2-dioxygenase subunit alpha (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Rhodococcus erythropolis IN119	Rhodococcus erythropolis KD-1 CP050124 99.42%	Rhodococcus erythropolis KD-1 CP050124 99.42%	Alkane 1-monooxygenase (5 copies), pentachlorophenol monooxygenase, cyclohexanone monooxygenase (2 copies), biphenyl 2,3-dioxygenase (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Rhodococcus globerulus IN113	Rhodococcus globerulus D757 CP079698 99.86%	Rhodococcus globerulus D757 CP079698 99.86%	Alkane 1-monooxygenase (3 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies), aromatic ring-hydroxylating dioxygenase subunit alpha (7 copies)
Raoultella sp. IN109	Raoultella planticola SCLZS62 CP082168 99.08%	Raoultella planticola SCLZS62 CP082168 99.08%	aromatic ring-hydroxylating dioxygenase subunit alpha (2 copies)

Identification of Fungal Strains by MALDI-TOF MS

Matrix-assisted laser desorption/ionisation time-of-flight mass spectrometry (MALDI-TOF MS) was used to identify specific fungal strains and was performed at the Jagiellonian Centre of Innovation (Kraków, Poland). A colony from a fresh overnight culture was used for sample preparation to obtain the mass spectrometer measurement. The material was suspended in 300 µL double-distilled water, following which, 900 µL ethanol was added, and the components were mixed well; thereafter, the mixture was centrifuged (2 min, 9000× *g*), and the supernatant was removed. For sample extraction, 40 µL of formic acid (70% in water) was added to the fungal pellet, following which, the components were mixed thoroughly, and 40 µL of acetonitrile was added. After performing centrifugation at 9000× *g* for 2 min, 1 µL of the supernatant containing the fungal extract was transferred to a well of the 96-well MALDI Biotarget plate (Bruker Daltonics, Bremen, Germany) and was allowed to dry at room temperature. Subsequently, the sample was overlaid with 2 µL of the MALDI matrix solution (α-cyano-4-hydroxy-cinnamic acid) and air-dried again. The measurements were performed using the MALDI BioTyper 2.0 Microflex LT system (Bruker), using the manufacturer's recommended settings. In order to perform microorganism identification, the raw protein spectral data were imported into the MALDI BioTyper 2.0 software (Bruker) and were analysed via standard pattern matching (with default parameter settings) against the spectral data in the BioTyper database (Bruker). Parameters such as the mass-to-charge ratio and peak intensity were considered to produce a matching score, which was subsequently used to rank the results. In these studies, the identification criteria were as follows: a score of 2.000 and above was considered sufficient

for correct species identification; score values of 1.700 to 1.999 confirmed genus identification; and scores lower than 1.700 indicated no identification.

International Biodeterioration & Biodegradation 206 (2026) 106217

Contents lists available at ScienceDirect

International Biodeterioration & Biodegradation

journal homepage: www.elsevier.com/locate/ibiod

Phytoremediation of soil contaminated with petroleum hydrocarbons using the wild plants *Scirpus sylvaticus* and *Cirsium oleraceum*, supported by bioaugmentation

Katarzyna Wojtowicz [a,*], Teresa Steliga [a], Joanna Brzeszcz [b], Janusz Fyda [c], Tomasz Skalski [d], Piotr Kapusta [b]

[a] Department of Reservoir Fluid Production Technology, Oil and Gas Institute – National Research Institute, Lubicz 25 A, Kraków, 31-503, Poland
[b] Department of Microbiology, Oil and Gas Institute – National Research Institute, Lubicz 25 A, Kraków, 31-503, Poland
[c] Institute of Environmental Sciences, Jagiellonian University in Kraków, Gronostajowa 7, Kraków, 30-387, Poland
[d] Biotechnology Centre, Silesian University of Technology, Krzywoustego 8, 44-100, Gliwice, Poland

ARTICLE INFO

Keywords:
Phytoremediation supported by bioaugmentation
Scirpus sylvaticus
Cirsium oleraceum
Soil contaminated with petroleum hydrocarbons
Soil ciliates

ABSTRACT

The research was aimed at determining the efficiency of phytoremediation supported by bioaugmentation using a microbial consortium based on autochthonous bacteria and wild plants naturally growing in an area of natural petroleum seeps (*Scirpus sylvaticus and Cirsium oleraceum)*,transplanted for the cleanup of hydrocarbon-contaminated soil. A six-month phytoremediation process resulted in a decrease in concentrations: TPH—from 4320 to 455–261 mg/kg dry mass, and PAHs—from 7.19 to 1.47–1.76 mg kg^{-1} dry mass. The greatest reductions were observed for n-C_{10}–n-C_{21} (91.9–99.8 %) and naphthalene (85.3–87.4 %), and the lowest for n-C_{30}–n-C_{36} (62.0–87.3 %) and 4–6-ring PAHs (44.0–76.3 %). Intense growth of roots and shoots was found in plants from bioaugmented soils. Toxicological studies performed using biotests with varying sensitivity (Ostracodtoxkit < Microtox < MARA) and phytotoxicity test (Phytotoxkit) indicated a decrease in toxicity levels after phytoremediation supported by bioaugmentation, which correlated with the drop in TPH and PAHs contents in the treated soil. The study also evaluated the structure of the microbial community in the remediated soil. The presence of bacteria in the rhizosphere accelerated the degradation of contaminants and increased plant tolerance to adverse environmental conditions. Additionally, changes in the community of soil ciliates were assessed, showingthat functional group responses, rather than species richness alone, may serve as sensitive indicators of soil recovery and ecological stability.This study highlights the novelty and ecological relevance of combining wild-adapted plants and microbial consortia for sustainable remediation, while also demonstrating the potential of ciliates as sensitive bioindicators of soil recovery.

1. Introduction

The intense progress of the petrochemical industry has considerably deteriorated the natural environment. It is estimated that 0.6 Mt of crude oil are introduced into the environment each year (Kvenvolden and Cooper, 2003). The primary sources of hydrocarbon contamination are leakages and spillages of crude oil generated during prospecting, extraction, transport, and storage (Zaltauskaitė et al., 2024). Another significant source of contamination is historical pollution, such as waste pits where drilling cuttings, bentonite-based drilling fluids, and petroleum-derived substances from oil-bearing formations were deposited. (Steliga et al., 2012). The presence of these pollutants in soil and water poses a serious threat to humans and animals (Macci et al., 2016; Guarino et al., 2017) due to their toxic, carcinogenic and mutagenic properties (Fatima et al., 2015; Aydin et al., 2017), which highlights the urgent need to develop effective remediation methods. Considering the limitations and environmental risks associated with conventional remediation techniques, biological methods have become increasingly favoured for restoring areas contaminated with petroleum hydrocarbons, with phytoremediation emerging as one of the most promising approaches.

Phytoremediation, involving symbiosis between plants and related

* Corresponding author.
E-mail address: katarzyna.wojtowicz@inig.pl (K. Wojtowicz).

https://doi.org/10.1016/j.ibiod.2025.106217
Received 5 May 2025; Received in revised form 28 August 2025; Accepted 15 September 2025
Available online 18 September 2025
0964-8305/© 2025 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

soil microorganisms in the root zone, has become a highly popular method for restoring contaminated soils. However, its efficiency depends on the contaminant type, the soil type, the applied plant species, and the presence of hydrocarbon degraders (Afzal et al., 2011, 2013; Khan et al., 2013; Zheng et al., 2018). The adaptation of plants to the contaminated environment plays an important role. Several studies have shown that wild plant species capable of growing in petroleum-contaminated areas, exhibit greater potential for rhizodegradation and phytostabilisation (Nemati et al., 2024; Melzi et al., 2024). In spite of this, the presence of pollutants may considerably constrain plant growth, and thus reduce the phytoremediation efficiency (Escalante-Espinosa et al., 2005; Kirk et al., 2005; Peng et al., 2009; Sun et al., 2010; Thion et al., 2013). Bioaugmentation may complement the catabolic activity of rhizosphere microorganisms, thus increasing a plant's tolerance to the contamination, enhancing plant growth, and biodegradation of contaminants (Sheng et al., 2008; Weyens et al., 2010; Glick and Stearns, 2011; Afzal et al., 2014). The root growth during phytoremediation leads to loosening of the soil and improves soil permeability and aeration (Weyens et al., 2009; Wojtowicz et al., 2023), thus contributing to an increase in hydrocarbon bioavailability. Certain microorganisms used in soil inoculation may affect the growth of phytoremediators by fixing nitrogen and increasing the solubility of phosphates and potassium (Eze et al., 2022). On the other hand, plant root exudates provide microorganisms with organic acids, sugars, and amino acids (Eze et al., 2022). However, it is important to note that indigenous microorganisms may compete with the introduced microorganisms for nutrients, which could potentially limit the effectiveness of bioaugmentation strategies, unless the microbial consortia are well adapted to the specific soil and plant conditions (Kuzyakov and Xu, 2013; Li et al., 2024a). To prevent this, bioaugmentation often uses consortia containing autochthonous hydrocarbon degraders. *Pseudomonas*, *Rhodococcus*, and *Mycobacterium*/*Mycolicibacterium* are well known for their diverse hydrocarbon degradation potential (Brzeszcz and Kaszycki, 2018). The metabolic capabilities towards both aliphatic and aromatic hydrocarbons were also found in other bacterial genera (for more detailed information, please see: Ehiosun et al., 2022, Zhao et al., 2021; Li et al., 2023, Brzeszcz et al., 2024).

The transformation of hydrocarbons largely depends on a network of trophic interactions between several types of soil microorganisms (Lara et al., 2007). Soil protozoa, and ciliates in particular, represent a distinct group of organisms in the rhizosphere, and constitute a key trophic level. They contribute to the nutrient cycling in the ecosystem by consuming bacteria and accelerating the turnover of bacterial biomass and organic matter, thus releasing nutrients accumulated in the microbial biomass, previously unavailable for primary producers (Acosta-Mercado et al., 2004; Wei et al., 2014; Mondragón-Camarilloet al., 2020; Jia et al., 2021). This essential role of ciliates in the soil environment makes them valuable bioindicators of environmental stress (Lara and Acosta-Mercado, 2012). Their abundant coexistence with bacteria or fungi suggests shared environmental preferences or potential biointeractions between them. It is necessary to study the ecological role of these protozoa as important soil microorganisms (Mondragón-Camarillo et al., 2020; Xu et al., 2022).

Phytoremediation-supported bioaugmentation presents great potential as a remediation method for hydrocarbon-contaminated areas. Previously, we prepared a microbial consortium exhibiting high potential to support phytoremediation (Wojtowicz et al., 2023a, 2023b). The objective of this research was to employ the mentioned approach through the use of plants tolerant to elevated hydrocarbon loads to clean up historically polluted soil. Therefore, we hypothesized that the combination of our consortium with wild plants (*Scirpus sylvaticus* and *Cirsium oleraceum*), naturally occurring in hydrocarbon-rich soils (natural oil seeps), might enhance the remediation of polluted soil. Our research aimed to: i) determine the efficiency of the applied remediation strategy, ii) evaluate soil quality toxicologically after treatments at multiple trophic levels, and iii) characterize the bacterial and ciliate communities in the treated soil.

Until now, research on phytoremediation supported by bioaugmentation has primarily focused on cultivated plants such as *Zea mays* and *Helianthus annuus* and their interactions with hydrocarbon-degrading microorganisms. Conversely, knowledge regarding the application of wild plants that naturally occur in petroleum-contaminated areas and their impact on soil biological communities remains limited. Furthermore, this study represents a preliminary investigation into the ciliate community in soil following phytoremediation, highlighting their roles as bioindicators of environmental stress and in soil formation and biogeochemical processes. It is imperative to comprehend these relationships in order to evaluate the efficacy and stability of soil reclamation processes.

A notable novelty of this study is the analysis of interactions between wild plants and soil microbiota, which facilitates a more profound comprehension of the mechanisms that support contaminant biodegradation and ecosystem restoration. This study constitutes the first attempt to determine the effectiveness of phytoremediation-supported bioaugmentation with *Scirpus sylvaticus* and *Cirsium oleraceum*. These plants grow in the areas of natural oil seeps and would therefore easily adapt to high hydrocarbon loads. The novelty also lies in the interactions occurring in the studied plant-soil microbiota system.

2. Materials and method

2.1. Soil, microbial consortium, and plants

The soil was collected from the area of a G6 waste pit located in south-eastern Poland (N: 49°40′16.22″, E: 22°04′42.99″). Fifty soil samples, weighing 2 kg each, were collected from each of two depth intervals: 0–0.25 and 0.25–1.00 m bgl. The samples were placed in tightly sealed containers, labelled, and delivered to a laboratory, where they were thoroughly mixed and homogenised. The physical and chemical parameters of the G6 soil are presented in Table S1 in Supplementary Materials. It was characterised by high TPH (4320.7 mg/kg dry mass, hereafter referred to as dm) and PAHs content (7.19 mg/kg dm). The content of individual hydrocarbon groups is presented in Fig. S1 in Supplementary Materials.

The microbial consortium (MC) consisted of autochthonous bacterial strains isolated from waste pit areas. Its composition and construction were described in our previous papers (Wojtowicz et al., 2023a, 2023b). It consisted of seven strains belonging to the following genera: *Dietzia*, *Gordonia*, *Mycolicibacterium*, *Pseudomonas*, *Raoultella*, and *Rhodococcus*. All were capable of degrading both aliphatic and aromatic hydrocarbons (Brzeszcz et al., 2024) (Table S2 in Supplementary Materials).

Two wild plant species were used: *Scirpus sylvaticus* and *Cirsium oleraceum*. They were collected from the area of a natural petroleum seep (N: 49°40′13.28″, E: 22°05′10.51″), near the soil sampling site. The plants were collected in early spring as young seedlings approximately 10 cm tall.

2.2. Description of the experiment

To determine the level of natural attenuation, the G6 soil (initial) was left for the duration of the experiment (6 months) - G6X. Then, the nitrogen and phosphorus contents were analysed, which showed that there were no favourable conditions for the development of microorganisms (N:P = 1:7) (Table S1). Based on literature data (Wu et al., 2017) and our previous research (Steliga and Kluk, 2020), the N:P ratio was adjusted to 10:1 — a proportion shown to enhance microbial activity by promoting balanced nutrient availability, which stimulates microbial growth and accelerates hydrocarbon biodegradation (Wu et al., 2017). This adjustment, along with the optimization of soil pH (7.6–7.8), was achieved by the application of 'Azofoska' mineral fertiliser and microelements (for nutrient supplementation), and fertiliser lime (to correct pH).The phytoremediation experiment was conducted

for a period of 6 months under real conditions at a prepared test station in boxes, where seedlings were planted. Twenty-kilogram soil samples were placed in experimental boxes (30 l), and planted with either *Scirpus sylvaticus* or *Cirsium oleraceum*. Half of the planted soils were inoculated with the MC consortium (10^8 CFU/cm^3). The whole study was conducted as follows:

a) natural attenuation, unplanted, unbioaugmented soil (G6X),
b) phytoremediation using *Scirpus sylvaticus* and supported by bioaugmentation (soil SS-G6-A),
c) phytoremediation using *Scirpus sylvaticus* (soil SS-G6-B),
d) phytoremediation using *Cirsium oleraceum* and supported by bioaugmentation (soil CO-G6-A),
e) phytoremediation using *Cirsium oleraceum* (soil CO-G6-B).

Soil moisture was maintained at 35–40 %. At the end of the experiment, the hydrocarbon content in soil and plant tissues (TPH, PAHs), plant growth, and soil quality (toxicity) were determined. Moreover, the characterisation of bacterial and ciliate communities was performed.

2.3. Analysis of TPH and PAHs in soil and plant tissues

Identification and quantitative determination of petroleum hydrocarbons in soil and plant material (roots, shoots) were carried out using gas chromatography (TPH analysis) and high-performance liquid chromatography (PAHs analysis) according to the procedures developed at the Department of Reservoir Fluids Exploitation Technology, INiG - PIB, as discussed in previous papers (Wojtowicz et al., 2022, 2023a). More details were provided in the Supplementary Materials.

2.4. Ecotoxicological analyses

Ecotoxicological analyses were performed by means of a set of tests, including the following: Microtox® Solid Phase, Ostracodtoxkit F, Phytotoxkit™, as well as MARA (Microbial Assay for Risk Assessment). The tests were performed following the standard procedures described previously (Kluk and Steliga, 2019; Wojtowicz et al., 2023a). More details were provided in the Supplementary Materials. The results were presented as EC_{50} and converted into TU. The test material constituted the reference G6 soil and soil after phytoremediation processes from test soils developed using wild plants supported by inoculation with an MC bacterial consortium.

2.5. Microbiological analysis

2.5.1. DNA extraction, bacterial 16S rRNA gene Illumina MiSeq-based sequencing, and sequence processing

The DNA isolation, 16S sequencing, and analysis were performed as described previously (Brzeszcz et al., 2020). Total DNA was extracted using Genomic Mini AX Bacteria Kit according to the manufacturer's protocol. The DNA quality and concentration were verified by electrophoresis and fluorometry (Qubit 4 Fluorometer, Thermo Fisher Scientific, USA). Library preparation, sequencing and sequence processing were done as previously (Brzeszcz et al., 2023). More details were provided in Supplementary Materials. The libraries were deposited in the NCBI Short Read Archive under BioProject accession number PRJNA1224491.

2.5.2. Determination and estimation of the abundance of ciliated protozoa

To estimate the abundance and diversity of ciliates in the treated and untreated soils, 10 g of each type of experimental soil was added separately to a 10 cm diameter Petri dish. Each sample was then saturated with 10 mL of deionised water. The samples were kept in a SANYO MLR-350 climate chamber at 20 ± 1 °C in the dark. Microscopic estimates of total diversity and abundance were made 3, 6, 9 and 12 days. We used the method (Finlay et al., 2000) with slight modifications. Soil interstitial water was collected by gently pressing the soil surface with a finger. A glass pipette was used to transfer 1 mL of the squeezed soil water into a Sedgewick-Rafter counting chamber S52 (Graticules Optics, UK). The abundance of ciliates and other organisms was then examined under a Nikon Eclipse E–80 microscope at 200 × magnification. Each time 200–400 μL of soil water was examined per sample. The number of ciliates per g dry soil was calculated taking into account the total volume of soil water in the incubated soil. The cumulative species diversity curve in terms of days counted represents an estimate of the total species richness of soil ciliates. Since statistically robust estimates of total diversity could be obtained using diversity estimates obtained on a few consecutive days, we counted ciliates on days 3, 6, 9 and 12. During the study, the protozoa were determined to the lowest possible taxonomic level using the Foissner keys (Foissner and Berger, 1996; Foissner et al., 1991, 1992, 1999).

2.6. Data analysis and statistical information

Statistical analysis of data from chemical and chromatographic studies was performed using Statistica 14.0. Standard deviation (SD) and the Pearson correlation coefficient were calculated. The obtained data were first tested for normal distribution. Then, they were analysed using a one-way analysis of variance (ANOVA), followed by Tukey's post hoc test (when the analysis of variance gave significant results). Significance was set at $p < 0.05$.

In microbiological studies of phytoremediation processes, an individual-based dilution method was used to indicate the effects of different treatments on community richness (Colwell et al., 2012). By estimating average species richness while accounting for variability in sampling effort, we elucidated patterns in taxa distribution along the treatments on different organisational levels, from Phylum to Genus. To create the rarefaction curve, random subsets or permutations of the collected data were generated. The calculations were made in PAST software, using a log-Gamma function for computing combinatorial terms (Hammer et al., 2001).

To test which environmental parameter are most responsible for microbial communities, redundancy analysis (RDA) was applied. In this method, a linear response between taxa and environmental variables should be obtained. Thus, we first chose detrended correspondence analysis (DCA) to measure the criterion where the longest gradient length of the taxa did not exceed the value of 3.0 (Lepš and Smilauer, 2003), and then a forward selection procedure was applied (Ter Braak and Smilauer, 2002).

3. Results

3.1. Biodegradation of petroleum hydrocarbons

A six-month bioremediation resulted in a TPH concentration drop from 4320.73–455.05 mg/kg dm in soil SS-G6-A and to 261.13 mg/kg dm in soil CO-G6-A. The respective values of TPH biodegradation levels in the planted and inoculated soils amounted to 89.47 % (SS-G6-A) and 93.96 % (CO-G6-A). In contrast, soils SS-G6-B and CO-G6-B, which were subjected to phytoremediation alone, reached 36.24 % and 43.61 % TPH reduction, respectively. In the self-purifying soil (G6X), the reduction was only 3.76 %. Corresponding TPH concentrations are presented in Table 1.

Individual n-alkane fractions (n-C_6–n-C_9, n-C_{10}–n-C_{21}, n-C_{22}–n-C_{30}, and n-C_{31}–n-C_{36}) were degraded to varying extents. The highest biodegradation levels were observed for the n-C_{10}–n-C_{21} range, with removal levels reaching up to 95.66 % in SS-G6-A and 97.2 % in CO-G6-A. In phytoremediated-only soils, the removal reached up to 55.08 % (SS-G6-B) and 61.5 % (CO-G6-B), while in G6X it did not exceed 6.11 %. Slightly lower efficiency was recorded for the n-C_6–n-C_9 and n-C_{22}–n-C_{30} fractions. The most recalcitrant were the long-chain n-C_{31}-n-C_{36} alkanes, with their content reduced by up to 77.97 % in SS-G6-A and 87.34 % in

Table 1
Concentrations of TPH and PAHs, and biodegradation ratios (n-C17/Pr, n-C18/Ph) in tested soils (n = 3, p < 0.05).

Soils	TPH [mg/kg dm]	nC_{17}/Pr	nC_{18}/Ph	PAHs [mg/kg dm]
Soil G6	4320.73 ± 258	2.64 ± 0.13	2.65 ± 0.14	7.19 ± 0.35
Soil G6X	4158.17 ± 228	2.54 ± 0.12	2.54 ± 0.12	7.00 ± 0.31
Soil SS-G6-A	455.05 ± 25	0.13 ± 0.01	0.15 ± 0.02	1.76 ± 0.08
Soil SS-G6-B	2755.10 ± 108	1.53 ± 0.09	1.71 ± 0.11	4.89 ± 0.25
Soil CO-G6-A	261.13 ± 13	0.05 ± 0.01	0.07 ± 0.01	1.47 ± 0.06
Soil CO-G6-B	2436.57 ± 95	1.39 ± 0.11	1.49 ± 0.07	4.31 ± 0.21

Soil PAHs were also removed as a result of the employed remediation procedures. Our data indicated that the combination of phytoremediation and bioaugmentation allowed for a reduction in PAHs concentrations by 75.52 % in soil SS-G6-A and 79.55 % in soil CO-G6-A. Phytoremediation alone caused a reduction of 31.99 % in soil SS-G6-B and 40.07 % in soil CO-G6-B, while in soil G6X, PAHs content decreased by 2.67 %. Thus, phytoremediation supported by bioaugmentation caused greater hydrocarbon degradation compared to phytoremediation alone. Corresponding PAHs concentrations are presented in Table 1.

CO-G6-A. In phytoremediated only soils, the decrease ranged from 17.73 % to 39.52 %, and did not exceed 2.70 % in G6X.The content of unidentified hydrocarbons decreased from 2049.43 to 1977.91 mg/kg dm (3.49 %) in soil G6X, and to 48.87–1382.77 mg/kg dm (corresponding to a 32.53–97.62 % reduction) in the remediated soils (see Table S3 in Supplementary Materials for concentration values of individual n-alkane fractions). The results obtained for the G6X sample did not show significant differences compared to the initial G6 soil, and therefore the G6 soil was treated as a control soil. Fig. 1 presents the concentration levels of individual n-alkanes in soil samples after bioremediation.

The biodegradation indices nC_{17}/Pr and nC_{18}/Ph in the soil G6 were, respectively, 2.64 and 2.65. For soil G6X, a slight drop in these parameters was observed. Among the remediated soils, higher values of biodegradation ratios were measured for phytoremediation-only soils (for SS-G6-B: nC_{17}/Pr = 1.53; nC_{18}/Ph = 1.71; for CO-G6-B: nC_{17}/Pr = 1.39; nC_{18}/Ph = 1.49). In turn, the lowest values were noted for the soils additionally bioaugmented (for SS-G6-A: nC_{17}/Pr = 0.13; nC_{18}/Ph = 0.15; CO-G6-A: nC_{17}/Pr = 0.05; nC_{18}/Ph = 0.07) (Table 1).

A detailed analysis of individual PAHs indicated that, as the number of aromatic rings in a molecule increases, its susceptibility to biodegradation decreases. The highest biodegradation levels were recorded for naphthalene, reaching 85.25 % in soil SS-G6-A, 87.40 % in soil CO-G6-A, up to 51.67 % in phytoremediated-only soils, and 3.40 % in G6X. The decrease of 3-ring PAHs content ranged from 79.25 to 84.94 % in soils




Fig. 1. Distribution of n-alkanes and PAHs content: initial and control soils (A and D); soils undergoing phytoremediation with inoculation of the MC consortium (B and E) and soils undergoing phytoremediation without inoculation (C and F) (n = 3, p < 0.05)
P- pristane, Ph-phytane, N – naphthalene, Ac – acenaphthene, Fluo – fluorene, Phen – phenanthrene, A-anthracene, F – Fluoranthene, Pyr – pyrene, BaA – benzo(a) anthracene, CH – chrysene, BbF – benzo(b)fluoranthene, BkF – benzo(k)fluoranthene, BaP – benzo(a)pyrene, DahA – dibenzo(a,h)anthracene, BghiP – benzo(g,h,i) perylene, Ind123P – indeno(1,2,3-c,d)pyrene.

subjected to phytoremediation with inoculation, up to 45.81 % in phytoremediated-only soils, and not exceeding 3.05 % in G6X. Biodegradation of 4- and 5-ring PAHs reached up to 69.49 % in SS-G6-A and 76.33 % in CO-G6-A. In phytoremediated-only soils, the removal ranged from 15.75 % to 32.35 %, while in G6X it did not exceed 2.49 %. The lowest biodegradation levels were observed for 6-ring PAHs, with values reaching up to 49.93 % (SS-G6-A) and 53.05 % (CO-G6-A). In phytoremediated-only soils, the degradation ranged from 10.72 % to 17.10 %, and did not exceed 0.99 % in G6X. (See Table S3 for concentration values of PAHs fractions). Fig. 1 presents the concentration of individual PAHs in soil after bioremediation.

Our results (Table 1) revealed that the most effective approach to remove hydrocarbons is a combined method, i.e., phytoremediation supported by bioaugmentation.

3.2. Analysis of plant material

3.2.1. Assessment of plant growth and biomass production

Soil bioaugmentation caused an increase in biomass production and the growth rate of *Scirpus sylvaticus* and *Cirsium oleraceum*, used as phytoremediators. Biomass produced by *Scirpus sylvaticus* and *Cirsium oleraceum* in bioaugmented soils amounted to 4.86 and 15.35 g/kg dm, respectively. In contrast, biomass production in uninoculated soils was much lower: 3.34 for *Scirpus sylvaticus* and 10.53 g/kg dm for *Cirsium oleraceum*, respectively.

Total dry biomass refers to the dry biomass of the plant roots and shoots. In the planted and inoculated soils, the dry root biomass was 1.97 for *Scirpus sylvaticus* and 7.87 g/kg dm for *Cirsium oleraceum*, while, in the uninoculated soils it was 1.30 and 5.83 g/kg dm, respectively. A similar trend was obtained for dry shoot biomass, which amounted to 2.89 in soil SS-G6-A, 2.04 in SS-G6-B, 7.48 in CO-G6-A, and 4.70 g/kg dm in CO-G6-B (Fig. 2A and B).

A morphological analysis of *Scirpus sylvaticus* collected from bioaugmented soil indicated intense plant growth over 6 months, including development of the root system and aboveground organs. Additionally, the formation of numerous young specimens was observed on stolons. Mature plants reached heights of 88–111 cm and were characterised by a short, triangular stem with flat, intensely green leaves measuring 89–109 cm in length. At the end of the experiment, the average shoot length in the inoculated soil was 103 cm. Its rapid growth, and substantial biomass increase in the inoculated soil contributed to the high hydrocarbon removal efficiencies (TPH = 89.47 %, PAHs = 75.52 %). In the uninoculated soil, *Scirpus sylvaticus* reached heights of 68–77 cm, and an average shoot length of 67 cm at the end of the experiment. Both its roots and leaves were less developed, which corresponded to lower phytoremediation efficiency (Fig. 2C).

A morphological analysis of *Cirsium oleraceum* after the end of the experiment indicated vigorous root growth (numerous filamentous roots branching out from a thick, cylindrical rhizome) in both inoculated and uninoculated soils. However, the roots collected from soil CO-G6-A were thicker and more intensely branched. The aboveground organs included a straight hollow stem and large leaves with variously shaped blades. The size and development of the aboveground organs were the main distinguishing features between plants from soils CO-G6-A and CO-G6-B. Mature specimens from the inoculated soil (CO-G6-A) reached heights from 155 to 170 cm, with an average shoot length of 150 cm. During summer, flower formation (cylindrical and oval capitula) was observed in two *Cirsium oleraceum* specimens. Intense growth of roots, stems, and leaves contributed to the efficiency of hydrocarbon biodegradation (TPH = 93.96 %, PAHs = 79.55 %). In the uninoculated soil, plant heights ranged from 92 to 105 cm, with an average shoot length of 96 cm at the end of the experiment. The roots and the leaves were less developed, and no inflorescence formation was observed (Fig. 2D).

3.2.2. TPH and PAHs content in plant tissues

The assessment of hydrocarbon biodegradation efficiency in the tested soils was supplemented by an analysis of hydrocarbon content in plant tissues. TPH concentrations in the roots of *Scirpus sylvaticus* grown in soils SS-G6-A and SS-G6-B were 150.4 and 136.7 mg/kg dm respectively, while in *Cirsium oleraceum* from soils CO-G6-A and CO-G6-B they were 529.7and 465.9 mg/kg dm, respectively. At the end of the experiment, TPH concentrations in shoots were as follows: *Scirpus sylvaticus* –




Fig. 2. Biomass increase of roots and shoots of: A) *Scirpus sylvaticus*, B) *Cirsium oleraceum* after the completion of the phytoremediation processes in soils, and average length of: C) *Scirpus sylvaticus*, D) *Cirsium oleraceum* during the phytoremediation experiment.




Fig. 3. TPH concentrations measured in the roots and shoots of A) *Scirpus sylvaticus*, B) *Cirsium oleraceum*, and PAHs concentrations measured in the roots and shoots of C) *Scirpus sylvaticus* and D) *Cirsium oleraceum* after the completion of phytoremediation experiment.

108.8 mg/kg dm (SS-G6-A) and 91.6 mg/kg dm (SS-G6-B); *Cirsium oleraceum* – 292.7 mg/kg dm (CO-G6-A) and 225.3 mg/kg dm (CO-G6-B) (Fig. 3A and B). The higher concentrations of aliphatic hydrocarbons in plants grown in soils SS-G6-A and CO-G6-A indicate that bio-augmentation considerably enhanced the phytoremediation process. Similar results were obtained for PAHs. In the roots of *Scirpus sylvaticus* grown in soils SS-G6-A and SS-G6-B, PAHs levels were 102.7 and 73.1 µg/kg dm, respectively. In *Cirsium oleraceum* from CO-G6-A and CO-G6-B, concentrations reached 583.3 and 463.9 µg/kg dm, respectively. In shoots, PAHs concentrations were: 85.8 (SS-G6-A), 41.3 (SS-G6-B), 429.5 (CO-G6-A), and 226.6 µg/kg dm (CO-G6-B) (Fig. 3C and D).

When comparing TPH and PAHs content in plant tissues, greater hydrocarbon accumulation was observed in the roots than in the shoots. Moreover, slightly higher concentrations were found in the tissues of *Cirsium oleraceum*. This might be the result of a much better-developed root system, which allows for more efficient functioning of mechanisms and interactions occurring in the plant-soil microbiota system.

3.3. Toxicological analysis

The toxicological tests indicated that the G6 reference soil exhibited noticeable toxic properties towards bacteria, crustaceans, and embryophytes, as evidenced by EC_{50} concentration values. In the Microtox, Ostracodtoxkit and MARA tests, the EC_{50} values were 11.3, 12.6, and 10.3 mg soil/dm^3, respectively, corresponding to toxic units (TU) of 10.3, 9.2, and 11.3 (Fig. 4). In the Phytotoxkit test, which assessed *Sorghum saccharatum*, *Lepidium sativum*, and *Sinapis alba*, the resulting EC_{50} values ranged from 13.9 to 16.8 mg soil/dm^3 (germination), and from 11.3 to 18.6 mg soil/dm^3 (root growth inhibition). No change in EC_{50} values was observed in soil G6X. Phytoremediation led to an increase in EC_{50} values across all toxicological tests. In the Microtox test, EC_{50} values for soils SS-G6-B and CO-G6-B were 15.2 and 16.4 mg soil/dm^3, respectively, in the Ostracodtoxkit test they were 17.9 and 18.7 mg soil/dm^3, and in MARA they were 14.6 and 15.6 mg soil/dm^3. In the Phytotoxkit test, EC_{50} values for soil remediated with *Scirpus sylvaticus*




Fig. 4. Toxicological test summary for soil samples subjected to phytoremediation by: A) *Scirpus sylvaticus*, B) *Cirsium oleraceum* ($n = 3$, $p < 0.05$).

reached up to 18.8 mg soil/dm^3 for germination and 19.9 mg soil/dm^3 for root growth inhibition. In the case of *Cirsium oleraceum*, the values reached up to 20.1 mg soil/dm^3 for germination and 22.1 mg soil/dm^3 for root growth inhibition. Soil inoculation with the MC consortium to support phytoremediation contributed to a further increase in EC_{50} values. In SS-G6-A soil, the EC_{50} values were 40.2 mg soil/dm^3 (Microtox), 42.6 mg soil/dm^3 (Ostracodtoxkit test), up to 48.5 mg soil/dm^3 for germination, and up to 54.3 mg soil/dm^3 for root growth inhibition in the Phytotoxkit test. In CO-G6-A soil, the values were 68.9 mg soil/dm^3 (Microtox), 78.4 mg soil/dm^3 (Ostracodtoxkit), up to 87.9 mg soil/dm^3 (germination), and up to 89.5 mg soil/dm^3 (root growth inhibition). The higher EC_{50} values observed in remediated soils indicate improved soil quality resulting from the applied remediation techniques (Fig. 4A and B). The consistently elevated EC_{50} values in soils subjected to phytoremediation supported by bioaugmentation confirm that the combined approach achieved the highest remediation efficiency.

3.4. Soil microbial community

3.4.1. Assessment of the bacterial community after phytoremediation processes

16S amplicon sequencing revealed that all treatments led to significant changes in the microbial diversity. The contaminated soil was characterised by the presence of Archaea (almost 2 % of total reads), represented almost exclusively by methanogens with predominance of *Methanosaeta* genus (1.54 %). Phytoremediated soils were almost devoid of Archaea (0.01–0.1 %). This clearly suggests that all remediated soils represent an aerobic environment, not appropriate for strict anaerobes as methanogens are. For Bacteria significant differences were observed even at phylum level (Fig. 5A). While Actinobacteriota and Proteobacteria occurred in the highest numbers in all analysed soils, relative abundance of the former was higher in contaminated soil (37.57 %) than in all remediated soils (10.06–28.23 %). The same was observed for Firmicutes; their abundance was higher in contaminated soil (2.86 %) than in remediated soils (0.24–1.08 %). The last phylum whose members prevailed in contaminated soil was Desulfobacterota (1.34 % vs. 0.23–0.72 %). On the other hand, relative abundance of Gemmatimonadota and Verrucomicrobiota was an order of magnitude lower in polluted soil (0.28 % and 0.30 % respectively) than in remediated soils (2.47–5.16 % and 3.02–5.74 %). Members of Gemmatimonadota are common in soils and are involved in diverse metabolic processes (Mujakić et al., 2022). Therefore their increased numbers in remediated soils can be an indirect indicator of contaminant removal. The same tendency was observed for representatives of Planctomycetota phylum; their relative abundance was lower in contaminated soil (0.3 %) in comparison with remediated soils (1.63–3.42 %). ASVs specific for Methylomirabilota were not present in contaminated soil, and their numbers were higher in both soils remediated with *Scirpus sylvaticus* (over 1 %) than in soils planted with *Cirsium oleraceum*.

Further evident differences were seen on family and genus levels (Fig. 5B and C). *Microbacteriaceae* (21.99 %) with *Agromyces* (11.21 %) and *Galbitalea* (8.04 %), *Rhodocyclaceae*, represented almost solely by *Sulfuritalea* (9.62 %), *Cellulomonadaceae* (4.59 %), with *Cellulomonas* (4.19 %), potential hydrocarbon degraders of *Pseudoxanthomonas* genus (Patel et al., 2012) (4.78 %), and *Rhodoferax* (4.53 %), dominated in contaminated soil. Their corresponding relative abundances in remediated soils were 0.66–5.57 % for *Microbacteriaceae*, 0.26–3.11 % for *Agromyces*, 0.51–2.27 % for *Galbitalea*, 0.32–2.77 % for *Sulfuritalea*, 0.20–1.50 % for *Cellulomonadaceae*, 0.20–1,24 % for *Cellulomonas*, 0.02–0.26 % for *Pseudoxanthomonas* and 0.26–1.18 % for *Rhodoferax*.

The remediated soils were characterised by the dominance of the hydrocarbon-degrading *Sphingomonas* genus (Cunliffe and Kertesz, 2006; Chaudhary and Kim, 2017) (5.33–10.99 %), *Porticoccaceae* (2.51–6.28 %) with *C1-B045* (2.08–5.92 %), *Gemmatimonadaceae* (3.05–5.91 %) with *Gemmatimonas* (1.82–3.64 %), *Acidithiobacillaceae* (1.23–4.65 %) with *KCM-B-112* (1.00–4.56 %), *Moraxellaceae* (0.97–5.08 %) with *Cavicella* (0.2–4.17 %), *Chthoniobacteraceae* represented almost exclusively by *Candidatus* Udaeobacter (1.22–3.39 %), *Chitinophagaceae* (1.81–2.95 %), *Nitrosomonadaceae* (0.94–3.54 %) with *Ellin6067* (0.61–3.54 %), *Arenimonas* (0.75–3.21 %), *Iamia* (0.74–2.81 %), and *Rhizobacter* (1.19–1.97 %). In contaminated soil, the relative abundance of the above-mentioned taxa was significantly lower; 3.14 % for *Sphingomonas*, 0.20 % for *Gemmatimonadaceae* (0.11 % for *Gemmatimonas*), 0.06 % for *Acidithiobacillaceae* with *KCM-B-112*, 0.16 % for *Candidatus* Udaeobacter, 0.70 % for *Chitinophagaceae* and 0.15 % for *Iamia*. No specific ASVs were found for *Porticoccaceae* (*C1-B045*), *Moraxellaceae* (*Cavicella*), *Nitrosomonadaceae* (*Ellin6067*), *Arenimonas* and *Rhizobacter*.

Potential hydrocarbon degraders other than *Sphingomonas*, such as: *Pseudomonas*, *Nocardioides* and *Pseudarthrobacter* (Zhang et al., 2016) were also present among 20 most abundant bacterial genera. No differences were observed between contaminated and remediated soils for *Nocardioides* (0.53–2.82 %) and *Pseudarthrobacter* (0.82–6.59 %), whereas there was a decrease in relative abundance of *Pseudomonas* in both soils remediated with *Scirpus sylvaticus* (0.91 %) when compared to contaminated soil (3.80 %) and those remediated with *Cirsium oleraceum* (2.44–2.72 %).

Bacteria that were introduced to remediated soils as members of the microbial consortium did not remain in numbers higher than those observed in contaminated soil, with the single exception of *Mycobacterium*. Its relative abundance was higher in remediated soils (2.98 % in SS–G6–A, 1.26 % in CO–G6–A) than in contaminated soil (0.6 %). Due to short sequences it was impossible to distinguish between *Mycobacterium* and *Mycolicibacterium* genera, they may also correspond to the introduced strain *Mycolicibacterium frederiksbergense* IN53.




Fig. 5. Effect of the phytoremediation and bioaugmentation on the relative abundance of bacterial taxa A) phylum, B) family (20 most abundant), C) genus (20 most abundant).

3.4.2. Statistical analysis of microbial communities

Rarefaction analysis for all levels of community organisation showed a positive effect of treatments on microbial diversity compared to the reference. The most significant gradient of diversity was observed at the genus level, where in contaminated soil G6 approximately 350 genera were predicted, while in SS-G6-A,more than 500 genera were detected. This indicates that the application of *Scirpus sylvaticus* and *Cirsium oleraceum* together with the MC consortium enriches the microbial communities and enhances soil enzymatic functionality (Fig. 6A).

Redundancy analysis described 90.6 % of species-environment variation. The first redundancy axis (Axis 1) revealed the gradient of plant-driven remediation. Some microbial phyla (Halobacterota, Desulfobacterota, Caldatribacteriota, Spirochaetota) were mostly associated with G6 soil, which had higher concentrations of all hydrocarbon parameters. The second axis, which describes 13.1 % of the variance, is mostly related to contamination. Along this gradient, the MC consortium was positioned opposite to the hydrocarbon vectors, indicating a positive effect of the consortium on hydrocarbon removal. Bacterial phyla positively associated with contamination included Actinobacteriota, Hydrogenedentota, and Proteobacteria, whereas groups avoiding contaminated areas were primarily WS2, NB1-j, Elusimicrobiota, and RCP2-54. Forward selection of environmental variables showed that the most influential parameter on microbial phyla was contaminated soil G6 ($F = 3.69$, $p < 0.03$), explaining 21 % of taxa variation. The second most influential parameter was the MC consortium ($F = 2.99$, $p < 0.05$), accounting for 18 % of taxa variation. The remaining environmental factors did not significantly improve overall taxa variation (Fig. 6B).

3.4.3. Diversity and abundance of ciliated protozoa

In total, 26 species of ciliated protozoa were identified. This analysis revealed the presence of 9–21 species of ciliated protozoa in the investigated soils. (Table S5 in Supplementary Materials). The highest diversity was observed in the soil SS-G6-B, whereas few species were found in the soil G6. Moderate diversity, reaching 17 species over all consecutive days of the study, was observed in soil CO-G6-B.

The number of ciliate species and their densities varied according to the sample and the day of analysis (Fig. 7). In G6 soil, the most abundant and dominant species on day 3 was *Colpoda inflata*, which had a density exceeding 2800 individuals/cm^3 (ind./cm^3). The remaining four species showed densities of approximately 160 ind./cm^3. On the following days, ciliate abundance decreased sharply, reaching a density of 170 ind./cm^3 for four species on day 12 (Fig. 7A). A similar pattern of change in abundance and species number was observed in soil SS-G6-A (Fig. 7B). The total density on day 3 reached 930 ind./cm^3, composed of four species, of which the dominant species was *Cyrtolophosis muscicola*, with a density of 815 ind./cm^3. The other three species had lower densities, ranging from 5 ind./cm^3 (*Perisincirra* sp.) to 95 ind./cm^3 (*Tachysoma pellionellum*). On successive days, the abundance of ciliates decreased, finally reaching three species with a total density of 35 ind./cm^3.

A different pattern was observed in soil SS-G6-B (Fig. 7C). Biodiversity was relatively high, reaching 11 species on Days 3 and 6. At the same time, the total density of these species was low, ranging from 375




Fig. 6. A) Individual rarefaction curves for the applied treatments and different levels of microbial variation from phylla to genus. soil G6 (red), soil CO-G6-B (green) soil CO G6-A (blue), soil SS-G6-B (turquoise), soil SS-G6-A (navy). For detailed sample description: see Section 3.4.1.1. B) Redundancy analysis of environmental factors (arrows) and microbial phyla (circles) in two dimensional space:P- pristane, Ph-phytane, for the rest of abbreviations see Fig. 1 and Table S4 in Supplementary materials.




Fig. 7. Changes in number of species (bars) and total abundance of ciliates (line) over time in A) G6, and treatments soils B) SS-G6-A, C) SS-G6-B, D) CO-G6-A, E) CO-G6-B.




Fig. 8. Changes in main ecological biodiversity indices values over time in the soil G6 (dark blue line), and treatments: SS-G6-A (orange), SS-G6-B (grey), CO-G6-A (yellow) and CO-G6-B (green). For detailed sample description see Section 2.2.

to 620 ind./cm^3. After 9 days, the number of species slightly decreased, while ciliate abundance increased, reaching 3770 ind./cm^3, mainly composed of *Cinetochilum margaritaceum*, a cosmopolitan bacteriophagous ciliate, with a density of 3700 ind./cm^3. The remaining six species had low abundance in the sample. On the last day, ciliate abundance decreased significantly to 1410 ind./cm^3, but biodiversity remained relatively high, represented by 9 species (Fig. 7C).

In the case of soils CO-G6-A and CO-G6-B, the differences in abundance and biodiversity were less pronounced (Fig. 7D and E). In both soils, the number of species ranged from a minimum of 4 to a maximum of 10, depending on the day of the experiment. In the CO-G6-A, the number of species reached 10 on day 3, was lowest on day 6, and remained stable at 8 species after days 9 and 12. Abundance followed the species richness pattern, ranging from 545 ind./cm^3 on day 3, to 205 ind./cm^3 on day 6, and to 675 ind./cm^3 on the last day. The addition of microbial consortium did not significantly affect ciliate species richness and abundance (Fig. 7E).

The biodiversity indices for the ciliate community are shown in Fig. 8A–F (or in Table S6 in Supplementary Materials). Simpson's dominance index in G6 reached 0.9, indicating strong dominance of a single species; on day 3, in the following days, it decreased to 0.29, on day 12. In soils SS-G6-A and SS-G6-B, the dominance index changed from 0.78 on day 3 to 0.54 on day 12, and from 0.18 to 0.85, respectively. The Shannon index (Fig. 8E) reached 2 at the beginning of the study for soil SS-G6-B and decreased to 0.12 on day 9. The most stable Shannon index values during the study were observed for soil CO-G6-A, fluctuating between 1.27 and 1.70 depending on the day of measurement. Similar values of H' were found for soil CO-G6-B, but it decreased to 0.76 on the last day of the study.

4. Discussion

Soil contamination with petroleum hydrocarbons (TPH and PAHs) affects plant growth by causing inhibition, and alters soil chemical properties and fertility (i.e., it reduces plant biomass, chlorophyll content, pH, and electrical conductivity). Moreover, the presence of TPH in the soil increases stress indicators (malondialdehyde and glutathione) and the activity of antioxidant enzymes (polyphenol oxidase and catalase) (Yuan et al., 2023). These compounds constitute a serious threat to soil quality, and are degraded via a number of biochemical processes involving microorganisms present both in the soil and in plants. Interactions in the soil-plant- microbiota system are critical for the efficient removal of petroleum hydrocarbons. Soil microorganisms are mostly responsible for the decomposition of contaminants via biodegradation, through which petroleum hydrocarbons are transformed into less toxic products, such as carbon dioxide or intermediate products (Kang et al., 2017; Eze et al., 2022).

As demonstrated in the extant literature, plants have the capacity to remove petroleum hydrocarbons through a variety of mechanisms, including phytoaccumulation, phytostabilization, phytodegradation, phytovolatilization, and rhizodegradation (Maliszewska-Kordybach, 2005; Hou et al., 2015; Xie et al., 2017). Phytoaccumulation and phytodegradation are defined as the absorption and accumulation of contaminants by the root system, as well as the adsorption of compounds on the root surface and their aggregation in the rhizosphere (Hou et al., 2021; Pourbabaee et al., 2021). The process of phytoaccumulation has been shown to result in the long-term immobilization of contaminants in roots or aboveground tissues (Cherian and Oliveira, 2005). The presence of TPH and PAHs in the roots and shoots of *Scirpus sylvaticus* and *Cirsium oleraceum* suggests that the aforementioned mechanisms may have been triggered by the bioaugmentation-assisted phytoremediation approach. In the phytostabilization mechanism, the mobility of contaminants is reduced through the promotion of soil particle aggregation by the root system and root secretions, which in turn promotes their immobilization in the soil (Shang et al., 2024). This phenomenon also supports the activity of microorganisms present in the bacterial consortium by enhancing their ability to degrade contaminants. Plants used in bioaugmentation-assisted phytoremediation can contribute to reducing the concentration of petroleum hydrocarbons in soil by degrading them through a process known as phytodegradation. Contaminants are broken down by plant-derived enzymes, which are secreted into the root zone and participate indirectly in degradation processes (Kang et al., 2013; Kumar et al., 2019). The most commonly identified plant enzymes involved in degrading petroleum hydrocarbons are peroxidases (Gan et al., 2022) and laccases (Diefenbach et al., 2024).

Furthermore, the interaction between plants and microorganisms in the rhizosphere has been demonstrated to significantly accelerate the degradation of petroleum hydrocarbons (Germida et al., 2002). Root secretions play a pivotal role in this process by increasing the bioavailability of pollutants and enhancing the degradation of compounds that are particularly resistant, such as polycyclic aromatic hydrocarbons (PAHs) (Liu et al., 2014). These substances act as both chemical signals and sources of carbon and energy, thereby stimulating the growth and activity of microorganisms capable of biodegrading hydrocarbons (Kuiper et al., 2004; Gerhardt et al., 2009).

The purpose of our research was to use two wild plants, *Scirpus sylvaticus and Cirsium oleraceum*, originating from a natural petroleum seep, for use in phytoremediation of soil contaminated with petroleum hydrocarbons (TPH and PAHs). This study proposes a novel approach to the remediation of such environments, as it is the first report on the use of these two species, which naturally colonize oil-contaminated habitats. The combination of wild plant species with a selected microbial consortium (MC), composed exclusively of hydrocarbon-degrading bacteria, represents a significant innovation. This approach integrates the adaptive potential of plants to survive in contaminated environments with the targeted degradation activity of hydrocarbon-degrading microorganisms.

There are currently no reports on phytoremediation using *Scirpus sylvaticus*. Pilot studies demonstrating phytoremediation of hydrocarbon-contaminated water using the closely related plant *Actinoscirpus grossus* (formerly a member of the *Scirpus* genus) revealed high degradation effectiveness of C_8–C_{34} hydrocarbons, as the TPH content dropped from 121.8 to 64.4 mg/kg over 72 days. Other results demonstrated that, due to interactions between *Actinoscirpus grossus* and rhizobacteria, this plant is capable of removing petroleum contaminants from water, offering an alternative, environmentally friendly solution (Al-Baldawi et al., 2015; Hunt et al., 2019). Al-Baldawi et al. (2014) further demonstrated that *Actinoscirpus grossus* grown in a horizontal subsurface flow constructed wetland system (HSFCW) removed up to 72.5 % of TPH from water and 59 % from sand within 63 days, even in the absence of aeration. Similarly, in a pilot-scale study, a 0.1 % diesel contamination was reduced by 82 % (Al-Baldawi et al., 2013). Studies using *Scirpus triqueter* in microcosms simulating the phytoremediation of diesel-contaminated wetland soils, conducted by Wei et al. (2014), demonstrated that contaminant removal in the rhizosphere zone reached 57.6–65.5 %, significantly higher than in non-vegetated soils (approximately 27.8–36.3 %). Similarly, Purwanti et al. (2023) reported that *Scirpus mucronatus* exhibited the ability to degrade hydrocarbons in diesel-contaminated sand, with removal efficiencies reaching approximately 42 %. This efficiency increased to about 52 % when rhizobacterial strains such as *Bacillus subtilis* and *B. licheniformis* were introduced.

Melzi et al. (2024) indicated that the phytoremediation of contaminated soils planted with *Zea mays* and *Helianthus annuus*, and inoculated with a soil-based suspension originating from a spill area covered with *Scirpus sylvaticus*, led to hydrocarbon removal rates of 80–82 % for *Z. mays* and 71–72 % for *H. annuus*. The bacterial strains isolated from the suspension were able to grow on specific hydrocarbons, suggesting that the rhizosphere of *Scirpus sylvaticus* harbors a microbial community with strong biodegradation capabilities.

In the present study, a six-month phytoremediation process using *Scirpus sylvaticus* resulted in TPH and PAHs removal of 36.24 and 31.99

%, respectively (soil SS-G6-B). A more pronounced effect was observed when *Cirsium oleraceum* was used as the phytoremediator, with TPH and PAHs levels reduced by 43.61 and 40.07 %, respectively (soil CO-G6-B). The efficiency of phytoremediation can be further enhanced by: identifying optimal combinations of plant species and microbial strains for specific contaminants, and supporting the process not only by stimulating the natural activity of plants and soil microbiota, but also by introducing a high number of hydrocarbon-degrading bacteria (Graj et al., 2013; Pacwa-Płociniczak et al., 2020; Hoang et al., 2021). As demonstrated in our previous studies, phytoremediation using *Echinacea purpurea* combined with microbial consortium B1 (comprising *Dietzia* sp. IN118, *Gordonia* sp. IN101, *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN119, *Rhodococcus globerulus* IN113, and *Raoultella* sp. IN109) and consortium B2 (B1 enriched with fungal strains *Aspergillus sydowii*, *Aspergillus versicolor*, *Candida* sp., *Cladosporium halotolerans*, and *Penicillium chrysogenum*) resulted in TPH removal of 35.56 % and 44.16 % for soil DW (from a waste pit), and 40.85 % and 49.78 % for soil OS (from an oil spill), respectively (Wojtowicz et al., 2023a). Similarly, in the case of *Zea mays*, the application of biopreparation B2 led to a 49.08 % reduction in TPH and a 40.74 % reduction in PAHs after six months of remediation (Wojtowicz et al., 2023b). These findings provide strong evidence for the effectiveness of phytoremediation when supported by targeted bioaugmentation.

In comparison with the findings of previous studies, the results obtained in the present research, employing the wild plant species *Cirsium oleraceum* and *Scirpus sylvaticus*, are clearly encouraging. When combined with the MC microbial consortium, TPH and PAHs removal reached 89.47 % and 75.52 % for *Scirpus sylvaticus* (SS-G6-A), and as much as 93.69 % and 79.55 % for *Cirsium oleraceum* (CO-G6-A), respectively. These values are considerably higher than those obtained with cultivated species such as *Zea mays* or *Echinacea purpurea*, thus highlighting the enhanced potential of wild plants adapted to oil-contaminated habitats. The superiority of wild plant species over cultivated ones in the phytoremediation of petroleum hydrocarbons is further supported by the findings of Muratova et al. (2008), who investigated a former oil-sludge pit using both cultivated crops (*Triticum aestivum*, *Secale cereale*, *Avena sativa*, *Hordeum vulgare*, *Sorghum bicolor*, *Panicum miliaceum*, *Zea mays*) and wild grasses (e.g., *Lolium perenne*, *Bromopsis inermis*, *Agropyron cristatum*, *Festuca pratensis*). Their study showed that winter rye (a cultivated species) achieved a TPH reduction of approximately 82 % after one year of growth. However, wild grass species maintained lower residual oil levels in the soil over a three-year period and supported larger microbial populations in the rhizosphere, including hydrocarbon-degrading bacteria. These observations indicate a higher long-term remediation efficiency associated with native vegetation. Furthermore, Liste and Prutz (2006) demonstrated that cultivated grasses such as *Lolium perenne* and alfalfa, exhibited poorer germination and slower hydrocarbon degradation under diesel contamination compared to prairie wild species. The cultivated species achieved lower TPH reduction rates and supported weaker microbial colonization in the rhizosphere.

However, it is important to acknowledge the limitations inherent in the use of *Scirpus sylvaticus* and *Cirsium oleraceum* in phytoremediation processes. *Scirpus sylvaticus* is a species adapted to moist environments and may not perform effectively in dry or highly permeable soils. Nevertheless, the capacity for rapid vegetative propagation makes it a valuable candidate for the stabilization of contaminated areas, particularly in wetland or riparian zones. In contrast, *Cirsium oleraceum* exhibits greater tolerance to moderate soil drying and can be introduced into a wider range of site conditions; however, its colonization rate is comparatively slower. These habitat-specific characteristics must be carefully considered when selecting and planning field-scale phytoremediation applications involving wild species.

The chromatographic results indicated that the aliphatic hydrocarbons were removed in the following order: n-C_{10}–n-C_{21}>n-C_6–n-C_9>n-C_{22}–n-C_{30}>n-C_{31}–n-C_{36}. A similar trend was observed for PAHs, with biodegradation levels decreasing as the number of rings in the molecule increased: 2-ring PAHs>3-ring-PAHs>4-ring-PAHs>5-ring-PAHs>6-ring-PAHs (Fig. 1, Table S3 in Supplementary Materials). This pattern is related to the chemical structure of hydrocarbons. Short-chain compounds are more easily biodegradable than longer ones, which are more hydrophobic and less bioavailable due to greater sorption on organic material. Hydrocarbon biodegradability is also influenced by factors such as chain branching, the presence and position of substituents, and the structure and number of rings (Guo et al., 2010; Abbasian et al., 2015; Li et al., 2023). These results showed a similar hydrocarbon biodegradation trend to our earlier studies, albeit at a slightly higher level (Steliga and Kluk, 2021; Wojtowicz et al., 2023a, 2023b).

The presence of hydrocarbons in the soil during the phytoremediation process might inhibit plant growth and development. The use of *Scirpus sylvaticus* and *Cirsium oleraceum* helps to reduce environmental stress and facilitates adaptation to the conditions prevalent in the remediated soil. A morphological analysis of plants collected from soils SS-G6-A and CO-G6-A demonstrated vigorous plant growth, including extensive development of both the root system and shoots. The average dry biomass of plants grown in inoculated soils (SS-G6-A and CO-G6-A) was approximately 45 % higher than that of plants from uninoculated soils (SS-G6-B and CO-G6-B). It was also observed that bioaugmentation contributed to an increase in the dry biomass of *Scirpus sylvaticus* by 51.5 % in the root and 41.7 % in the shoot. In the case of *Cirsium oleraceum*, the average dry biomass of the roots in the inoculated soil was 35.0 % higher, and 59.1 % higher for the shoots, compared to the uninoculated soil (Figs. 2 and 3).

Bioaugmentation with autochthonous bacteria enhances the degradation potential of plants, thereby increasing phytoremediation efficiency, as confirmed by other studies (Vacheron et al., 2013; Bhatt et al., 2023). Orji et al. (2015) showed that inoculation of *Panicum maximum* with hydrocarbon-degrading bacteria (*Pseudomonas* sp. and *Bacillus* sp.) under petroleum contamination led to a significant increase in both shoot height and root biomass compared to uninoculated controls. Similarly, a study by Eze et al. (2022) with *Medicago sativa* inoculated with a plant growth-promoting and hydrocarbon-degrading consortium showed a 66 % increase in plant biomass and a 91 % reduction in diesel hydrocarbons within 60 days, clearly demonstrating a synergistic effect on plant growth and remediation efficacy. Plants originating from wetlands are capable of producing indole-3-acetic acid (IAA), a phytohormone that plays a key role in root elongation and contributes to increased root exudation, thereby enhancing nutrient absorption (Bennis et al., 2022). Moreover, studies have demonstrated the presence of plant growth-promoting (PGP) traits in strains isolated from *Scirpus sylvaticus*, which suggests that this plant can recruit bacteria capable of supporting its growth (Melzi et al., 2024). Our data unambiguously indicate that bioaugmentation led to better adaptation of the wild plants, naturally growing in a petroleum seep area, to the conditions present in the treated soils, as evidenced by the development of their root system and shoots, as well as their ability to accumulate hydrocarbons (Figs. 2 and 3).

Remediation of soils leads to significant changes in their microbiomes. This phenomenon is well documented and has been observed in several studies, both in soils and sediments remediated with (Dagher et al., 2019) and without the help of plants (Brzeszcz et al., 2020; Pacwa-Płociniczak et al., 2020). In this study, we recorded a clear decrease in typical obligate anaerobic microorganisms, such as methanogenic archaea and members of *Desulfobacterota* phylum, in all remediated soils. A similar trend was observed in our earlier study (Brzeszcz et al., 2020), where the abundance of anaerobic bacteria (e.g. those belonging to the Anaerolineae class) was significantly reduced in remediated microcosms. On the other hand, the decreased abundance of some aerobic and facultative anaerobic bacterial taxa may suggest that they were either outcompeted by other microorganisms or subjected to increased protozoan predation in remediated soils. This was particularly evident for members of *Agromyces*, *Cellulomonas*, *Galbitalea*, *Pseudoxanthomonas*,

Rhodoferax, and *Sulfuritalea* genera. Interestingly, some of these genera (*Pseudoxanthomonas* and *Sulfuritalea*) may actually prefer higher concentrations of hydrocarbons in their environment (Sheng et al., 2016; Brzeszcz et al., 2020, 2023; Cébron et al., 2022), which could explain their reduced numbers in the treated soils.

The remediated soils also contained members of the microbial community that were absent in the untreated, contaminated soils. Therefore, these these taxa were presumably part of the microbiomes inhabiting the rhizospheres of both plant species. While *Rhizobacter* may be considered an undesirable companion (the species is a well-known plant pathogen) (Goto and Kuwata, 1988), representatives of other genera (*Arenimonas*, *C1-B045*, *Cavicella*, and *Ellin6067*) appear to be common in hydrocarbon-impacted environments (Young et al., 2007; Nebeska et al., 2021; Fang et al., 2024), and may even participate in degradation processes (Cui et al., 2024).

We showed that bioaugmentation of hydrocarbon-contaminated soil resulted in an increased abundance of *Mycobacterium/Mycolibacterium*. This result is consistent with our earlier study (Brzeszcz et al., 2020) and with findings reported by others (Wolf et al., 2019; Zhao et al., 2021). It suggests that *Mycobacterium/Mycolicibacterium* is capable of persisting in remediated soils. The presence of *Mycobacterium* bacteria in the rhizosphere accelerates the degradation of organic pollutants and increases plant resistance to pollutants present in the soil, thereby enhancing plant tolerance to unfavourable environmental conditions (Dai et al., 2020).

The applied set of toxicological tests includes bioindicators representing all trophic groups: producers, consumers, and decomposers. This approach enables a comprehensive assessment of the soil quality and determination of the phytoremediation efficiency (Foucault et al., 2013; Oleszczuk et al., 2014; Steliga and Kluk, 2020, 2021). The G6 soil (TPH = 4320.73 mg/kg dm, PAHs = 7.19 mg/kg dm) was characterised by elevated toxicity, as evidenced by the EC_{50} -derived toxicity unit (TU): Microtox: (10.3), Ostracodtoxkit(9.2), MARA(11.3), and Phytotoxkit (6.2–8.2). After phytoremediation with *Scirpus sylvaticus* (SS-G6-B) toxicity decreased notably: Microtox to TU = 7.7; Ostracodtoxkit to TU = 6.5; MARA to TU = 8.0; Phytotoxkit to TU = 6.1–7.0. A slightly greater drop in toxicity was observed for *Cirsium oleraceum* (CO–G6–A). The application of bioaugmentation further reduced toxicity in both variants (SS-G6-A: TU = 2.1–2.4 and CO-G6-A: TU = 1.3–1.7). Earlier studies using *Echinacea purpurea* and *Zea mays*, supported by microbial consortia with comparable hydrocarbon degradation capabilities, reported similar TU values in all applied ecotoxicological tests after six months of remediation. In soils planted with *Echinacea purpurea* and supported by bioaugmentation with microbial consortia and γ-PGA, toxicity significantly decreased in all tests. In the Phytotoxkit, TU values dropped from 10,5 to 12.2 to 3,3–4.0 (Soil DW) and from 8,5 to 9.8 to 2, 9–3.7 (Soil OS); in the Ostracodtoxkit, from 11.2 to 4.2 and 10.9 to 3.6, respectively; and in the Microtox assay, from 14.1 to 4.4 and 11.4 to 3.9 (Wojtowicz et al., 2023a). Similarly, in soils planted with *Zea mays*, a significant decrease in toxicity was observed in the Microtox test, with the toxicity unit (TU) value dropping from 15.7 to 2.7 after six months of phytoremediation supported by bioaugmentation using the B2 biopreparation. (Wojtowicz et al., 2023a).

The decreased toxicity correlated with reductions in hydrocarbon content and increased plant biomass (roots and shoots) (Table 1 and Figs. 2 and 3). A particularly important source of information was the Phytotoxkit test, which directly reflects the phytotoxicity of soils towards higher plants. The assay, based on seed germination and root growth inhibition of *Lepidium sativum*, *Sorghum saccharatum*, and *Sinapis alba*, demonstrated that these plants responded sensitively to petroleum-derived contaminants. The strongest inhibition of root growth was observed in *Sorghum saccharatum*, while *Lepidium sativum* showed the lowest sensitivity among the applied bioindicators. After phytoremediation supported by bioaugmentation, root growth inhibition was clearly reduced, indicating that the soils became less phytotoxic and more favourable for plant growth. Thus, the decrease in TU values translates into a real improvement in soil quality and better adaptation of plants to growth and development in remediated areas. The ecotoxicological analyses also revealed the relative sensitivity of the applied tests. In the Phytotoxkit assay, plant responses followed the order *Lepidium sativum* < *Sorghum saccharatum* < *Sinapis alba*. The application of plants with different sensitivity levels in the Phytotoxkit test proved useful for detecting subtle changes in soil toxicity during the remediation process. Across the applied bioassays, sensitivity decreased in the following order: Ostracodtoxkit < Microtox < MARA. This observation is consistent with the findings of other researchers (Oleszczuk et al., 2014; Stefaniuk et al., 2015; Steliga and Kluk, 2021; Paskuliakova et al., 2018).

Considering their functional diversity, protozoa (ciliates) regulate various soil microorganisms and influence the cycling of soil components. The biological properties of the rhizosphere, such as bacterial diversity and abundance, can significantly affect the structure of the ciliate community. These results are consistent with the mutualism model between plants and protozoa, mediated by soil bacteria (Acosta-Mercado et al., 2004; Lara et al., 2007; Li et al., 2024a,b). To the best of our knowledge, this study represents the first attempt to comprehensively assess the response of the ciliate community in phytoremediation processes using wild plants in soils contaminated with petroleum hydrocarbons. This innovative approach is of considerable scientific and practical importance, as it allows not only for the assessment of remediation effectiveness using microbial indicators, but also for the identification of potential bioindicators of soil ecosystem stabilization. Although knowledge of the vast functional diversity of soil ciliates is still incomplete, the existing database provides a substantial starting point for bioindication tests (Lara and Acosta-Mercado, 2012; Xu et al., 2022).

Across the various phytoremediation treatments and the control, distinct patterns were observed in ciliate abundance, species richness, and functional group composition. In the reference soil (G6), nine protozoan species were detected, including *Colpoda inflata*, a cosmopolitan bacterivore playing a critical role in soil ecosystem functioning and plant growth (Li et al., 2024a,b). However, its rapid decline in abundance likely reflects encystment due to limited food availability or predatory/competitive pressure from Hypotrichida, highlighting the strong influence of soil conditions on ciliate community composition (Bharti et al., 2024). In soils inoculated with a microbial consortium and planted with *Scirpus sylvaticus* (SS-G6-A), total ciliate abundance was reduced, with dominance by *Cyrtolophosis muscicola* and *Tachysoma pellionellum*, and absence of Colpodida. This suggests that bioaugmentation shifts the bacterial food base, thereby favouring selected functional groups. In contrast, *Scirpus sylvaticus* alone (SS-G6-B) supported higher species richness, decreasing from 12 species on the third day to 10 species on the twelfth day of the study, and included both omnivores and predators, indicating a more balanced community assembly and greater functional stability. Similarly, *Cirsium oleraceum* with bioaugmentation (CO-G6-A) sustained both species richness and abundance, except for a transient dip on day 6, followed by recovery of bacterivorous taxa. In contrast, *C. oleraceum* alone (CO-G6-B) exhibited a marked increase in abundance, from 130 to 740 ind./cm^3, while species richness fluctuated between 8, 11, and 5 species on consecutive days, with clear dominance of *Cinetochilum margaritaceum*. This indicates that native vegetation alone may support high abundance but less stable richness over time.

Biodiversity indices corroborated these trends: the Shannon index showed the most stable values in soils CO-G6-A and CO-G6-B throughout the experiment (Fig. 8). The observed differences between treatments underscore how phytoremediation with wild plants, with or without bioaugmentation, shapes soil protozoan communities. Phytoremediation alone fostered higher initial richness and functional diversity (including omnivores and predators), whereas bioaugmentation suppressed early richness but allowed eventual recovery in the abundance of key bacterivores. These findings align with previous studies (Lara et al., 2007; Mondragón-Camarillo et al., 2020; Cao et al., 2022),

which demonstrate that plant identity and soil management practices significantly influence ciliate community structure via rhizosphere-driven variation in microbial substrates. Overall, the integration of phytoremediation and bioaugmentation in petroleum-contaminated soils highlights the dual role of these treatments in both promoting contaminant degradation and reshaping protozoan communities. Functional group responses, rather than species richness alone, may serve as sensitive indicators of soil recovery and ecological stability.

Comparing these results with our previous studies on phytoremediation using other plant species: *Melilotus officinalis* (Steliga and Kluk, 2021), *Zea mays* (Wojtowicz et al., 2023b), and *Echinacea purpurea* (Wojtowicz et al., 2023a), it can be concluded that the application of wild-growing plants from a natural petroleum seep contributed to achieving a higher level of TPH and PAHs removal. This, in turn, correlates with a reduction in soil toxicity following remediation. The wild-growing species *Scirpus sylvaticus* and *Cirsium oleraceum*, used as phytoremediators, are suitable candidates for such applications, as they are capable of withstanding contamination and enhancing the soil's biodegradation potential by enriching it with microbial taxa that form part of rhizospheres of both plant species.

5. Conclusions

Our results unambiguously confirm that *Scirpus sylvaticus* and *Cirsium oleraceum*, transplanted from a petroleum seep area, are capable of remediating soil contaminated with petroleum hydrocarbons. To enhance the remediation efficiency, phytoremediation-assisted bioaugmentation proves to be a more effective strategy than phytoremediation alone, as demonstrated by our chromatographic, toxicological, and plant morphology analyses. The presence of the applied plant species altered the composition and structure of both bacterial and ciliate communities, however, the available data are insufficient to draw definitive conclusions regarding bacterial–protozoan interactions. To sum up, wild-growing plants from highly contaminated sites appear to be promising candidates for phytoremediation and their use as a key component of a phytoremediation-assisted bioaugmentation strategy is well justified. Moreover, these findings provide a necessary foundation for further research aimed at better understanding the interactions within the plant-soil microbiota system and their effect on soil remediation. Furthermore, this research suggests the potential for extending future research to include the co-cultivation of *Scirpus sylvaticus* and *Cirsium oleraceum*, which could offer additional insights into the synergistic effects of mixed planting on remediation efficiency, microbial diversity, and ecosystem resilience.

CRediT authorship contribution statement

Katarzyna Wojtowicz: Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis, Conceptualization. **Teresa Steliga:** Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis, Conceptualization. **Joanna Brzeszcz:** Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis, Conceptualization. **Janusz Fyda:** Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis, Conceptualization. **Tomasz Skalski:** Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis, Conceptualization. **Piotr Kapusta:** Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis, Conceptualization.

Funding

"This research was financially supported by Polish Ministry of Science and Higher Education within statutory funding for Oil and Gas Institute-National Research Institute" DK:4100-16-24.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary data to this article can be found online at https://doi.org/10.1016/j.ibiod.2025.106217.

Data availability

Data will be made available on request.

References

Abbasian, F., Lockington, R., Mallavarapu, M., Naidu, R., 2015. A comprehensive review of aliphatic hydrocarbon biodegradation by bacteria. Appl. Biochem. Biotechnol. 176, 670–699. https://doi.org/10.1007/s12010-015-1603-5.

Al-Baldawi, I.A., Sheikh Abdullah, S.R., Anuar, N., Suja, F., Idris, M., 2013. Performance assessment of pilot horizontal sub-surface flow constructed wetlands for removal of diesel from wastewater by Scirpus grossus. Water Sci. Technol. 68, 2271–2278. https://doi.org/10.2166/wst.2013.484.

Al-Baldawi, I.A.W., Sheikh Abdullah, S.R., Abu Hasan, H., Suja, F., Anuar, N., Mushrifah, I., 2014. Optimized conditions for phytoremediation of diesel by Scirpus grossus in horizontal subsurface flow constructed wetlands (HSFCWs) using response surface methodology. J. Environ. Manag. 140, 152–159. https://doi.org/10.1016/j.jenvman.2014.03.007.

Acosta-Mercado, D., Lynn, D.H., 2004. Soil ciliate species richness and abundance associated with the rhizosphere of different subtropical plant species. J. Eukaryot. Microbiol. 51, 582–588. https://doi.org/10.1111/j.1550-7408.2004.tb00295.x.

Afzal, M., Yousaf, S., Reichenauer, T.G., Kuffner, M., Sessitsch, A., 2011. Soil type affects plant colonization, activity and catabolic gene expression of inoculated bacterial strains during phytoremediation of diesel. J. Hazard. Mater. 186, 1568–1575. https://doi.org/10.1016/j.jhazmat.2010.12.040.

Afzal, M., Khan, S., Iqbal, S., Mirza, M.S., Khan, Q.M., 2013. Inoculation method affects colonization and activity of Burkholderia phytofirmans PsJN during phytoremediation of diesel-contaminated soil. Int. Biodeter.Biodegr. 85, 331–336. https://doi.org/10.1016/j.ibiod.2013.08.022.

Afzal, M., Khan, Q.M., Sessitsch, A., 2014. Endophytic bacteria: prospects and applications for the phytoremediation of organic pollutants. Chemosphere 117, 232–242. https://doi.org/10.1016/j.chemosphere.2014.06.078.

Al-Baldawi, I.A., Abdullah, S.R.S., Anuar, N., Suja, F., Mushrifah, I., 2015. Phytodegradation of total petroleum hydrocarbon (TPH) in diesel-contaminated water using Scirpus grossus. Ecol. Eng. 74, 463–473. https://doi.org/10.1016/j.ecoleng.2014.11.007.

Aydin, S., Karaçay, H.A., Shahi, A., Gökçe, S., Ince, B., Ince, O., 2017. Aerobic and anaerobic fungal metabolism and omics insights for increasing polycyclic aromatic hydrocarbons biodegradation. Fungal Biol. Rev. 31, 61–72. https://doi.org/10.1016/j.fbr.2016.12.001.

Bennis, M., Perez-Tapia, V., Alami, S., Bouhnik, O., Lamin, H., Abdelmoumen, H., Bedmar, E.J., Missbah El Idrissi, M., 2022. Characterization of plant growth-promoting bacteria isolated from the rhizosphere of Robinia pseudoacacia growing in metal-contaminated mine tailings in eastern Morocco. J. Environ. Manag. 304, 114321. https://doi.org/10.1016/j.jenvman.2021.114321.

Bharti, D., Kumar, S., Basuri, C.K., La Terza, A., 2024. Ciliated protist communities in soil: contrasting patterns in natural sites and arable lands across Italy. Soil Syst. 8 (2), 64. https://doi.org/10.3390/soilsystems8020064.

Bhatt, P., Ganesan, S., Santhose, I., Durairaj, T., 2023. Phytoremediation as an effective tool to handle emerging contaminants. Phys. Sci. Rev. 8, 2321–2340. https://doi.org/10.1515/psr-2021-0062.

Brzeszcz, J., Kapusta, P., Steliga, T., Turkiewicz, A., 2020. Hydrocarbon removal by two differently developed microbial inoculants and comparing their actions with biostimulationtreatment. Molecules 25, 661. https://doi.org/10.3390/molecules25030661.

Brzeszcz, J., Kaszycki, P., 2018. Aerobic bacteria degrading both n-alkanes and aromatic hydrocarbons: an undervalued strategy for metabolic diversity and flexibility. Biodegradation 29, 359–407. https://doi.org/10.1007/s10532-018-9837-x.

Brzeszcz, J., Skalski, T., Jankowski, L., Kapusta, P., 2023. How do microbial communities deal with chronic hydrocarbon presence in oil seep soils? Data from historical hand-dug oil Wells. Land Degrad. Dev. 34, 1283–1296. https://doi.org/10.1002/ldr.4531.

Brzeszcz, J., Steliga, T., Ryszka, P., Kaszycki, P., Kapusta, P., 2024. Bacteria degrading both n-alkanes and aromatic hydrocarbons are prevalent in soils. Environ. Sci. Pollut. Res. 31, 5668–5683. https://doi.org/10.1007/s11356-023-31405-8.

Cao, X., Cui, X., Xie, M., Zhao, R., Xu, L., Ni, S., Cui, Z., 2022. Amendments and bioaugmentation enhanced phytoremediation and micro-ecology for PAHs and heavy metals co-contaminated soils. J. Hazard. Mater. 426, 128096. https://doi.org/10.1016/j.jhazmat.2021.128096. Epub 2021 Dec 16. PMID: 34952500.

Cébron, A., Borreca, A., Beguiristain, T., Biache, C., Faure, P., 2022. Taxonomic and functional trait-based approaches suggest that aerobic and anaerobic soil

microorganisms allow the natural attenuation of oil from natural seeps. Sci. Rep. 12, 7245. https://doi.org/10.1038/s41598-022-10850-4.
Chaudhary, D.K., Kim, J., 2017. Sphingomonas olei sp. Nov., with the ability to degrade aliphatic hydrocarbons, isolated from oil-contaminated soil. Int. J. Syst. Evol. Microbiol. 67, 2731–2738. https://doi.org/10.1099/ijsem.0.002010.
Cherian, S., Oliveira, M.M., 2005. Transgenic plants in phytoremediation: recent advances and new possibilities. Environ. Sci. Technol. 39, 9377–9390. https://doi.org/10.1021/es051134l.
Colwell, R.K., Chao, A., Gotelli, N.J., Lin, S.-Y., Mao, C.X., Chazdon, R.L., Longino, J.T., 2012. Models and estimators linking individual-based and sample-based rarefaction, extrapolation and comparison of assemblages. J. Plant Ecol. 5, 3–21. https://doi.org/10.1093/jpe/rtr044.
Cui, Z., Li, Y., Jing, X., Luan, X., Liu, N., Liu, J., Meng, Y., Xu, J., Valentine, D.L., 2024. Cycloalkane degradation by an uncultivated novel genus of Gammaproteobacteria derived from China's marginal seas. J. Hazard. Mater. 469, 133904. https://doi.org/10.1016/j.jhazmat.2024.133904.
Cunliffe, M., Kertesz, M.A., 2006. Autecological properties of soil sphingomonads involved in the degradation of polycyclic aromatic hydrocarbons. Appl. Microbiol. Biotechnol. 72, 1083–1089. https://doi.org/10.1007/s00253-006-0374-x.
Dagher, D.J., De La Providencia, I.E., Pitre, F.E., St-Arnaud, M., Hijri, M., 2019. Plant identity shaped rhizospheric microbial communities more strongly than bacterial bioaugmentation in petroleum hydrocarbon-polluted sediments. Front. Microbiol. 10, 2144. https://doi.org/10.3389/fmicb.2019.02144.
Dai, Y., Liu, R., Zhou, Y., Li, N., Hou, L., Ma, Q., Gao, B., 2020. Fire phoenix facilitates phytoremediation of PAH-Cd co-contaminated soil through promotion of beneficial rhizosphere bacterial communities. Environ. Int. 136, 105421. https://doi.org/10.1016/j.envint.2019.105421.
Diefenbach, T., Sumetzberger-Hasinger, M., Braunschmid, V., Konegger, H., Heipieper, H.J., Guebitz, G.M., Lackner, M., Ribitsch, D., Loibner, A.P., 2024. Laccase-mediated degradation of petroleum hydrocarbons in historically contaminated soil. Chemosphere 348, 140733. https://doi.org/10.1016/j.chemosphere.2023.140733.
Ehiosun, K.I., Godin, S., Urios, L., Lobinski, R., Grimaud, R., 2022. Degradation of long-chain alkanes through biofilm formation by bacteria isolated from oil-polluted soil. Int. Biodeterior. Biodegrad. 175, 105508. https://doi.org/10.1016/j.ibiod.2022.105508.
Escalante-Espinosa, E., Gallegos-Martínez, M.E., Favela-Torres, E., Gutiérrez-Rojas, M., 2005. Improvement of the hydrocarbon phytoremediation rate by Cyperus laxus lam. inoculated with a microbial consortium in a model system. Chemosphere 59, 405–413. https://doi.org/10.1016/j.chemosphere.2004.10.034.
Eze, M.O., Thiel, V., Hose, G.C., George, S.C., Daniel, R., 2022. Bacteria-plant interactions synergistically enhance biodegradation of diesel fuel hydrocarbons. Commun. Earth Environ. 3, 192. https://doi.org/10.1038/s43247-022-00526-2.
Fang, X., Zhang, M., Zheng, P., Wang, H., Wang, K., Lv, J., Shi, F., 2024. Biochar-bacteria-plant combined potential for remediation of oil-contaminated soil. Front. Microbiol. 15, 1343366. https://doi.org/10.3389/fmicb.2024.1343366.
Fatima, K., Afzal, M., Imran, A., Khan, Q.M., 2015. Bacterial rhizosphere and endospherepopulations associated with grasses and trees to be used for phytoremediation of crude oil contaminated soil. Bull. Environ. Contam. Toxicol. 94, 314–320. https://doi.org/10.1007/s00128-015-1489-5.
Finlay, B.J., Black, H.I.J., Brown, S., Clarke, K.J., Esteban, G.F., Hindle, R.M., Olmo, J.L., Rollett, A., Vickerman, K., 2000. Estimating the growth potential of the soil protozoan community. Protist 151, 69–80. https://doi.org/10.1078/1434-4610-00008.
Foissner, W., Berger, H., 1996. A user-friendly guide to the ciliates (protozoa, ciliophora) commonly used by hydrobiologists as bioindicators in rivers, lakes, and waste waters, with notes on their ecology. Freshw. Biol. 35, 375–482. https://doi.org/10.1111/j.1365-2427.1996.tb01775.x.
Foissner, W., Berger, H., Kohmann, F., 1992. Taxonomische Und Ökologische Revision Der Ciliaten Des Saprobien Systems - Band II: Peritrichia, Heterotrichida, Odontostomatida. BayerischesLandesamt für Wasserwirtschaft, München, Germany.
Foissner, W., Berger, H., Schaumburg, J., 1999. Identification and Ecology of Limnetic Plankton Ciliates. BayerischesLandesamt für Wasserwirtschaft, München, Germany.
Foissner, W., Blatterer, H., Berger, H., Kohmann, F., 1991. Taxonomische und ökologischerevision der Ciliaten des saprobiensystems. Band I: Cyrtophorida, Oligotrichida, Hypotrichia, Colpodea. BayerischesLandesamt für Wasserwirtschaft, München, Germany.
Foucault, Y., Durand, M.-J., Tack, K., Schreck, E., Geret, F., Leveque, T., Pradere, P., Goix, S., Dumat, C., 2013. Use of ecotoxicity test and ecoscores to improve the management of polluted soils: case of a secondary lead smelter plant. J. Hazard. Mater. 246–247, 291–299. https://doi.org/10.1016/j.jhazmat.2012.12.042.
Gan, J., Bilal, M., Li, X., Hussain Shah, S.Z., Mohamed, B.A., Hadibarata, T., Cheng, H., 2022. Peroxidases-based enticing biotechnological platforms for biodegradation and biotransformation of emerging contaminants. Chemosphere 307, 136035. https://doi.org/10.1016/j.chemosphere.2022.136035.
Gerhardt, K.E., Huang, X.-D., Glick, B.R., Greenberg, B.M., 2009. Phytoremediation and rhizoremediation of organic soil contaminants: potential and challenges. Plant Sci. 176, 20–30. https://doi.org/10.1016/j.plantsci.2008.09.014.
Germida, J.J., Frick, C.M., Farrell, R.E., 2002. Phytoremediation of oil-contaminated soils. In: Developments in Soil Science. Elsevier, pp. 169–186. https://doi.org/10.1016/S0166-2481(02)80015-0.
Glick, B.R., Stearns, J.C., 2011. Making phytoremediation work better: maximizing a plant's growth potential in the midst of adversity. Int. J. Phytoremediation 13, 4–16. https://doi.org/10.1080/15226514.2011.568533.
Goto, M., Kuwata, H., 1988. Rhizobacter daucus gen. nov., sp. Nov., the causal agent of carrot bacterial gall. Int. J. Syst. Bacteriol. 38, 233–239. https://doi.org/10.1099/00207713-38-3-233.
Graj, W., Lisiecki, P., Szulc, A., Chrzanowski, Ł., Wojtera-Kwiczor, J., 2013. Bioaugmentation with petroleum-degrading consortia has a selective growth-promoting impact on crop plants germinated in diesel oil-contaminated soil. Water air. Soil Pollut. 224, 1676. https://doi.org/10.1007/s11270-013-1676-0.
Guarino, C., Spada, V., Sciarrillo, R., 2017. Assessment of three approaches of bioremediation (natural attenuation, landfarming and bioaugumentation – assistitedlandfarming) for a petroleum hydrocarbons contaminated soil. Chemosphere 170, 10–16. https://doi.org/10.1016/j.chemosphere.2016.11.165.
Guo, X., Luo, L., Ma, Y., Zhang, S., 2010. Sorption of polycyclic aromatic hydrocarbons on particulate organic matters. J. Hazard. Mater. 173, 130–136. https://doi.org/10.1016/j.jhazmat.2009.08.065.
Hammer, Ø., Harper, D.A.T., Ryan, P.D., 2001. PAST: paleontological statistics software package for education and data analysis. Palaeontol. Electron. 4, 1–9.
Hoang, S.A., Lamb, D., Seshadri, B., Sarkar, B., Choppala, G., Kirkham, M.B., Bolan, N.S., 2021. Rhizoremediation as a green technology for the remediation of petroleum hydrocarbon-contaminated soils. J. Hazard. Mater. 401, 123282. https://doi.org/10.1016/j.jhazmat.2020.123282.
Hou, J., Liu, W., Wang, B., Wang, Q., Luo, Y., Franks, A.E., 2015. PGPR enhanced phytoremediation of petroleum contaminated soil and rhizosphere microbial community response. Chemosphere 138, 592–598. https://doi.org/10.1016/j.chemosphere.2015.07.025.
Hou, J., Wang, Q., Liu, W., Zhong, D., Ge, Y., Christie, P., Luo, Y., 2021. Soil microbial community and association network shift induced by several tall fescue cultivars during the phytoremediation of a petroleum hydrocarbon-contaminated soil. Sci. Total Environ. 792, 148411. https://doi.org/10.1016/j.scitotenv.2021.148411.
Hunt, L.J., Duca, D., Dan, T., Knopper, L.D., 2019. Petroleum hydrocarbon (PHC) uptake in plants: a literature review. Environ. Pollut. 245, 472–484. https://doi.org/10.1016/j.envpol.2018.11.012.
Jia, T., Liang, X., Guo, T., Chai, B., 2021. Impact of nutrients on protozoa community diversity and structure in litter of two natural grass species in a copper tailings dam, China. Microorganisms 9, 2250. https://doi.org/10.3390/microorganisms9112250.
Kang, F., Mao, X., Wang, X., Wang, J., Yang, B., Gao, Y., 2017. Sources and health risks of polycyclic aromatic hydrocarbons during haze days in eastern China: a 1-year case study in nanjing city. Ecotoxicol. Environ. Saf. 140, 76–83. https://doi.org/10.1016/j.ecoenv.2017.02.022.
Kang, Y.-H., Liu, S.-H., Wan, S.-Q., Wang, R.-S., 2013. Assessment of soil enzyme activities of saline–sodic soil under drip irrigation in the songnen plain. Paddy Water Environ. 11, 87–95. https://doi.org/10.1007/s10333-011-0295-x.
Kuzyakov, Y., Xu, X., 2013. Competition between roots and microorganisms for nitrogen: mechanisms and ecological relevance. New Phytol. 198, 656–669. https://doi.org/10.1111/nph.12235.
Khan, S., Afzal, M., Iqbal, S., Khan, Q.M., 2013. Plant–bacteria partnerships for the remediation of hydrocarbon contaminated soils. Chemosphere 90, 1317–1332. https://doi.org/10.1016/j.chemosphere.2012.09.045.
Kirk, J.L., Klironomos, J.N., Lee, H., Trevors, J.T., 2005. The effects of perennial ryegrass and alfalfa on microbial abundance and diversity in petroleum contaminated soil. Environ. Pollut. 133, 455–465. https://doi.org/10.1016/j.envpol.2004.06.002.
Kluk, D., Steliga, T., 2019. Potential of Helianthus annuus for phytoremediation of lead, zinc, total petroleum hydrocarbons (TPH) and polycyclic aromatic hydrocarbons (PAHs) contaminated soil. Nafta Gaz. 75, 379–387. https://doi.org/10.18668/NG.2019.07.01.
Kuiper, I., Lagendijk, E.L., Bloemberg, G.V., Lugtenberg, B.J.J., 2004. Rhizoremediation: a beneficial plant-microbe interaction. MPMI (Mol. Plant-Microbe Interact.) 17, 6–15. https://doi.org/10.1094/MPMI.2004.17.1.6.
Kumar, N., Jeena, N., Kumar, N., Gangola, S., Singh, H., 2019. Phytoremediation facilitating enzymes: an enzymatic approach for enhancing remediation process. In: Smart Bioremediation Technologies. Elsevier, pp. 289–306. https://doi.org/10.1016/B978-0-12-818307-6.00015-9.
Kvenvolden, K.A., Cooper, C.K., 2003. Natural seepage of crude oil into the marine environment. Geo Mar. Lett. 23, 140–146. https://doi.org/10.1007/s00367-003-0135-0.
Lara, E., Acosta-Mercado, D., 2012. A molecular perspective on ciliates as soil bioindicators. Eur. J. Soil Biol. 49, 107–111. https://doi.org/10.1016/j.ejsobi.2011.11.001.
Lara, E., Berney, C., Ekelund, F., Harms, H., Chatzinotas, A., 2007. Molecular comparison of cultivable protozoa from a pristine and a polycyclic aromatic hydrocarbon polluted site. Soil Biol. Biochem. 39, 139–148. https://doi.org/10.1016/j.soilbio.2006.06.017.
Lepš, J., Šmilauer, P., 2003. Multivariate Analysis of Ecological Data Using CANOCO, first ed. Cambridge University Press. https://doi.org/10.1017/CBO9780511615146.
Li, C., Cui, C., Zhang, J., Shen, J., He, B., Long, Y., Ye, J., 2023. Biodegradation of petroleum hydrocarbons based pollutants in contaminated soil by exogenous effective microorganisms and indigenous microbiome. Ecotoxicol. Environ. Saf. 253, 114673. https://doi.org/10.1016/j.ecoenv.2023.114673.
Li, H., Hu, X., Geng, X., Xiao, B., Miao, W., Xu, Z., Deng, Y., Jiang, B., Hou, Y., 2024a. Competition mode and soil nutrient status shape the role of soil microbes in the diversity–invasibility relationship. Ecol. Evol. 14, e11425. https://doi.org/10.1002/ece3.11425.
Li, H., Wu, K., Feng, Y., Gao, C., Wang, Y., Zhang, Yuanyuan, Pan, J., Shen, X., Zufall, R.A., Zhang, Yu, Zhang, W., Sun, J., Ye, Z., Li, W., Lynch, M., Long, H., 2024b. Integrative analyses on the ciliates colpoda illuminate the life history evolution of soil microorganisms. mSystems 9, e0137923. https://doi.org/10.1128/msystems.01379-23.

Liste, H.-H., Prutz, I., 2006. Plant performance, dioxygenase-expressing rhizosphere bacteria, and biodegradation of weathered hydrocarbons in contaminated soil. Chemosphere 62, 1411–1420. https://doi.org/10.1016/j.chemosphere.2005.05.018.

Liu, W., Hou, J., Wang, Q., Ding, L., Luo, Y., 2014. Isolation and characterization of plant growth-promoting rhizobacteria and their effects on phytoremediation of petroleum-contaminated saline-alkali soil. Chemosphere 117, 303–308. https://doi.org/10.1016/j.chemosphere.2014.07.026.

Macci, C., Doni, S., Peruzzi, E., Mennone, C., Masciandaro, G., 2016. Biostimulation of soil microbial activity through organic fertilizer and almond tree association. Land Degrad. Dev. 27, 335–345. https://doi.org/10.1002/ldr.2234.

Maliszewska-Kordybach, B., 2005. Dissipation of polycyclic aromatic hydrocarbons in freshly contaminated soils – the effect of soil physicochemical properties and aging. Water Air Soil Pollut. 168, 113–128. https://doi.org/10.1007/s11270-005-0940-3.

Melzi, A., Zecchin, S., Gomarasca, S., Abruzzese, A., Cavalca, L., 2024. Ecological indicators and biological resources for hydrocarbon rhizoremediation in a protected area. Front. Bioeng. Biotechnol. 12. https://doi.org/10.3389/fbioe.2024.1379947.

Mondragón-Camarillo, L., Zaragoza, S.R., Mendoza-López, MaR., Cabirol, N., Macek, M., 2020. Recovery of soil protozoan community structure promoted by M. sativaafter a strong pulse of hydrocarbon contamination. Water Air Soil Pollut. 231, 283. https://doi.org/10.1007/s11270-020-04618-7.

Mujakić, I., Piwosz, K., Koblížek, M., 2022. Phylum gemmatimonadota and its role in the environment. Microorganisms 10, 151. https://doi.org/10.3390/microorganisms10010151.

Muratova, A.Yu, Dmitrieva, T.V., Panchenko, L.V., Turkovskaya, O.V., 2008. Phytoremediation of oil-Sludge-Contaminated soil. Int. J. Phytoremediation 10, 486–502. https://doi.org/10.1080/15226510802114920.

Nebeská, D., Trögl, J., Ševců, A., Špánek, R., Marková, K., Davis, L., Burdová, H., Pidlisnyuk, V., 2021. Miscanthus x giganteus role in phytodegradation and changes in bacterial community of soil contaminated by petroleum industry. Ecotoxicol. Environ. Saf. 224, 112630. https://doi.org/10.1016/j.ecoenv.2021.112630.

Nemati, B., Baneshi, M.M., Akbari, H., Dehghani, R., Mostafaii, G., 2024. Phytoremediation of pollutants in oil-contaminated soils by alhagi camelorum: evaluation and modeling. Sci. Rep. 14, 5502. https://doi.org/10.1038/s41598-024-56214-y.

Oleszczuk, P., Jośko, I., Kuśmierz, M., Futa, B., Wielgosz, E., Ligęza, S., Pranagal, J., 2014. Microbiological, biochemical and ecotoxicological evaluation of soils in the area of biochar production in relation to polycyclic aromatic hydrocarbon content. Geoderma 213, 502–511. https://doi.org/10.1016/j.geoderma.2013.08.027.

Pacwa-Plociniczak, M., Biniecka, P., Bondarczuk, K., Piotrowska-Seget, Z., 2020. Metagenomic functional profiling reveals differences in bacterial composition and function during bioaugmentation of aged petroleum-contaminated soil. Front. Microbiol. 11, 2106. https://doi.org/10.3389/fmicb.2020.02106.

Paskuliakova, A., McGowan, T., Tonry, S., Touzet, N., 2018. Phycoremediation of landfill leachate with the chlorophyte Chlamydomonas sp. SW15aRL and evaluation of toxicity pre and post treatment. Ecotoxicol. Environ. Saf. 147, 622–630. https://doi.org/10.1016/j.ecoenv.2017.09.010.

Patel, V., Cheturvedula, S., Madamwar, D., 2012. Phenanthrene degradation by pseudoxanthomonas sp. DMVP2 isolated from hydrocarbon contaminated sediment of amlakhadi canal, Gujarat, India. J. Hazard. Mater. 201–202, 43–51. https://doi.org/10.1016/j.jhazmat.2011.11.002.

Peng, S., Zhou, Q., Cai, Z., Zhang, Z., 2009. Phytoremediation of petroleum contaminated soils by mirabilis jalapa L. in a greenhouse plot experiment. J. Hazard. Mater. 168, 1490–1496. https://doi.org/10.1016/j.jhazmat.2009.03.036.

Pourbabaee, A.A., Khazaei, M., Alikhani, H.A., Emami, S., 2021. Root nodulation of alfalfa by ensifermeliloti in petroleum contaminated soil. Rhizosphere 17, 100305. https://doi.org/10.1016/j.rhisph.2021.100305.

Purwanti, I.F., Abdullah, S.R.S., Hamzah, A., Idris, M., Basri, H., Latif, M.T., Mukhlisin, M., Kurniawan, S.B., Imron, M.F., 2023. Maximizing diesel removal from contaminated sand using Scirpus mucronatus and assessment of rhizobacteria addition effect. Heliyon 9, e21737. https://doi.org/10.1016/j.heliyon.2023.e21737.

Shang, X., Wu, S., Liu, Y., Zhang, K., Guo, M., Zhou, Y., Zhu, J., Li, X., Miao, R., 2024. Rice husk and its derived biochar assist phytoremediation of heavy metals and PAHs co-contaminated soils but differently affect bacterial community. J. Hazard Mater. 466, 133684. https://doi.org/10.1016/j.jhazmat.2024.133684.

Sheng, X., Chen, X., He, L., 2008. Characteristics of an endophytic pyrene-degrading bacterium of enterobacter sp. 12J1 from Allium macrostemon bunge. Int. Biodeterior. Biodegrad. 62, 88–95. https://doi.org/10.1016/j.ibiod.2007.12.003.

Sheng, Y., Wang, G., Hao, C., Xie, Q., Zhang, Q., 2016. Microbial community structures in petroleum contaminated soils at an oil field, Hebei, China. CLEAN Soil Air Water 44, 829–839. https://doi.org/10.1002/clen.201500142.

Stefaniuk, M., Bartmiński, P., Różyło, K., Dębicki, R., Oleszczuk, P., 2015. Ecotoxicological assessment of residues from different biogas production plants used as fertilizer for soil. J. Hazard. Mater. 298, 195–202. https://doi.org/10.1016/j.jhazmat.2015.05.026.

Steliga, T., Jakubowicz, P., Kapusta, P., 2012. Changes in toxicity during in situ bioremediation of weathered drill wastes contaminated with petroleum hydrocarbons. Bioresour. Technol. 125, 1–10. https://doi.org/10.1016/j.biortech.2012.08.092.

Steliga, T., Kluk, D., 2021. Assessment of the suitability of Melilotus officinalis for phytoremediation of soil contaminated with petroleum hydrocarbons (TPH and PAH), Zn, Pb and Cd based on toxicological tests. Toxics 9, 148. https://doi.org/10.3390/toxics9070148.

Steliga, T., Kluk, D., 2020. Application of Festuca arundinacea in phytoremediation of soils contaminated with Pb, Ni, Cd and petroleum hydrocarbons. Ecotoxicol. Environ. Saf. 194, 110409. https://doi.org/10.1016/j.ecoenv.2020.110409.

Sun, T.-R., Cang, L., Wang, Q.-Y., Zhou, D.-M., Cheng, J.-M., Xu, H., 2010. Roles of abiotic losses, microbes, plant roots, and root exudates on phytoremediation of PAHs in a barren soil. J. Hazard. Mater. 176, 919–925. https://doi.org/10.1016/j.jhazmat.2009.11.124.

Ter Braak, C.J.F., Smilauer, P.N., 2002. CANOCO reference manual and CanoDraw for windows user's guide: software for canonical community ordination (version 4.5) (Microcomputer Power). www.canoco.com. Ithaca NY, USA.

Thion, C., Cébron, A., Beguiristain, T., Leyval, C., 2013. Inoculation of PAH-degrading strains of Fusarium solani and Arthrobacter oxydans in rhizospheric sand and soil microcosms: microbial interactions and PAH dissipation. Biodegradation 24, 569–581. https://doi.org/10.1007/s10532-013-9628-3.

Vacheron, J., Desbrosses, G., Bouffaud, M.-L., Touraine, B., Moënne-Loccoz, Y., Muller, D., Legendre, L., Wisniewski-Dyé, F., Prigent-Combaret, C., 2013. Plant growth-promoting rhizobacteria and root system functioning. Front. Plant Sci. 4. https://doi.org/10.3389/fpls.2013.00356.

Wei, J., Liu, X., Zhang, X., Chen, X., Liu, S., Chen, L., 2014. Rhizosphere effect of Scirpus triqueter on soil microbial structure during phytoremediation of diesel-contaminated wetland. Environ. Technol. 35, 514–520. https://doi.org/10.1080/09593330.2013.836549.

Weyens, N., Croes, S., Dupae, J., Newman, L., Van Der Lelie, D., Carleer, R., Vangronsveld, J., 2010. Endophytic bacteria improve phytoremediation of Ni and TCE co-contamination. Environ. Pollut. 158, 2422–2427. https://doi.org/10.1016/j.envpol.2010.04.004.

Weyens, N., Van Der Lelie, D., Taghavi, S., Newman, L., Vangronsveld, J., 2009. Exploiting plant–microbe partnerships to improve biomass production and remediation. Trends Biotechnol. 27, 591–598. https://doi.org/10.1016/j.tibtech.2009.07.006.

Wojtowicz, K., Steliga, T., Kapusta, P., 2023a. Evaluation of the effectiveness of bioaugmentation-assisted phytoremediation of soils contaminated with petroleum hydrocarbons using Echinacea purpurea. Appl. Sci. 13, 13077. https://doi.org/10.3390/app132413077.

Wojtowicz, K., Steliga, T., Kapusta, P., Brzeszcz, J., 2023b. Oil-contaminated soil remediation with biodegradation by autochthonous microorganisms and phytoremediation by maize (zea mays). Molecules 28, 6104. https://doi.org/10.3390/molecules28166104.

Wojtowicz, K., Steliga, T., Kapusta, P., Brzeszcz, J., Skalski, T., 2022. Evaluation of the effectiveness of the biopreparation in combination with the polymer γ-PGA for the biodegradation of petroleum contaminants in soil. Materials 15, 400. https://doi.org/10.3390/ma15020400.

Wolf, D.C., Cryder, Z., Gan, J., 2019. Soil bacterial community dynamics following surfactant addition and bioaugmentation in pyrene-contaminated soils. Chemosphere 231, 93–102. https://doi.org/10.1016/j.chemosphere.2019.05.145.

Wu, M., Li, W., Dick, W.A., Ye, X., Chen, K., Kost, D., Chen, L., 2017. Bioremediation of hydrocarbon degradation in a petroleum-contaminated soil and microbial population and activity determination. Chemosphere 169, 124–130. https://doi.org/10.1016/j.chemosphere.2016.11.059.

Xie, W., Zhang, Y., Li, R., Yang, H., Wu, T., Zhao, L., Lu, Z., 2017. The responses of two native plant species to soil petroleum contamination in the yellow river Delta, China. Environ. Sci. Pollut. Res. 24, 24438–24446. https://doi.org/10.1007/s11356-017-0085-0.

Xu, R., Zhang, M., Lin, H., Gao, P., Yang, Z., Wang, D., Sun, X., Li, B., Wang, Q., Sun, W., 2022. Response of soil protozoa to acid mine drainage in a contaminated terrace. J. Hazard. Mater. 421, 126790. https://doi.org/10.1016/j.jhazmat.2021.126790.

Young, C.-C., Kämpfer, P., Ho, M.-J., Busse, H.-J., Huber, B.E., Arun, A.B., Shen, F.-T., Lai, W.-A., Rekha, P.D., 2007. Arenimonasmalthae sp. Nov., a gammaproteobacterium isolated from an oil-contaminated site. Int. J. Syst. Evol. Microbiol. 57, 2790–2793. https://doi.org/10.1099/ijs.0.64975-0.

Yuan, L., Wu, Y., Fan, Q., Li, P., Liang, J., Liu, Y., Ma, R., Li, R., Shi, L., 2023. Remediating petroleum hydrocarbons in highly saline–alkali soils using three native plant species. J. Environ. Manag. 339, 117928. https://doi.org/10.1016/j.jenvman.2023.117928.

Žaltauskaitė, J., Meištininkas, R., Dikšaitytė, A., Degutytė-Fomins, L., Mildažienė, V., Naučienė, Z., Žūkienė, R., Koga, K., 2024. Heavy fuel oil-contaminated soil remediation by individual and bioaugmentation-assisted phytoremediation with Medicago sativa and with cold plasma-treated M. sativa. Environ. Sci. Pollut. Res. 31, 30026–30038. https://doi.org/10.1007/s11356-024-33182-4.

Zhang, H., Sun, H., Yang, R., Li, S., Zhou, M., Gao, T., An, L., Chen, X., Dyson, P., 2016. Complete genome sequence of a psychotrophicPseudarthrobactersulfonivorans strain Ar51 (CGMCC 4.7316), a novel crude oil and multi benzene compounds degradation strain. J. Biotechnol. 231, 81–82. https://doi.org/10.1016/j.jbiotec.2016.04.010.

Zhao, X., Miao, R., Guo, M., Zhou, Y., 2021. Effects of fire phoenix (a genotype mixture of fesctucaarundinecea L.) and mycobacterium sp. on the degradation of PAHs and bacterial community in soil. Environ. Sci. Pollut. Res. 28, 25692–25700. https://doi.org/10.1007/s11356-021-12432-9.

Zheng, M., Wang, W., Hayes, M., Nydell, A., Tarr, M.A., Van Bael, S.A., Papadopoulos, K., 2018. Degradation of macondo 252 oil by endophytic Pseudomonas putida. J. Environ. Chem. Eng. 6, 643–648. https://doi.org/10.1016/j.jece.2017.12.071.

Suplementary materials

Phytoremediation of soil contaminated with petroleum hydrocarbons using the wild plants *Scirpus sylvaticus* and *Cirsium oleraceum*, supported by bioaugmentation

Katarzyna Wojtowicz [1*], Teresa Steliga [1], Joanna Brzeszcz [2], Janusz Fyda[3] Tomasz Skalski[4], Piotr Kapusta[2]

[1] Department of Production Technology of Reservoir Fluids, Oil and Gas Institute – National Research Institute, Lubicz 25 A, 31-503 Kraków, Poland

[2] Department of Microbiology, Oil and Gas Institute – National Research Institute, Lubicz 25 A, 31-503 Kraków, Poland

[3] Institute of Environmental Sciences, Jagiellonian University in Kraków, Gronostajowa 7, 30-387, Kraków, Poland

[4] Biotechnology Centre, Silesian University of Technology, Krzywoustego 8, 44-100 Gliwice, Poland

* Correspondence: katarzyna.wojtowicz@inig.pl

Table S1 A summary of the results of physical and chemical analyses of raw soil G6

Parameter		Unit	Value
pH of water extract of soil (1:10)		-	**6.57**
Moisture		**%**	**24.1**
TPH		mg/kg dry mass	**4320.73**
∑PAH		mg/kg dry mass	**7.19**
Cresols		mg/kg dry mass	1.64
Phenols		mg/kg dry mass	1.35
2,4-dimethylphenol		mg/kg dry mass	0.13
3,4-dimethylphenol		mg/kg dry mass	2.01
Cyanides		mg/kg dry mass	0.07
Biphenyls		mg/kg dry mass	741.36
Cl^-		mg/kg dry mass	188.9
SO_4^{2-}		mg/kg dry mass	742.6
NH_4^+		mg/kg dry mass	**39.1**
NO_3^-		mg/kg dry mass	**158.7**
PO_4^{3-}		mg/kg dry mass	**12.7**
S^{2-}		mg/kg dry mass	10.1
SiO_2		mg/kg dry mass	446 756
Al_2O_3		mg/kg dry mass	891 007
Fe_2O_3		mg/kg dry mass	5 014
CaO		mg/kg dry mass	6 952
MgO		mg/kg dry mass	3 757
Mn		mg/kg dry mass	91.11
Heavy metal content	*As*	mg/kg dry mass	0.70
	Ba	mg/kg dry mass	27.2
	Cd	mg/kg dry mass	0.7
	Cr	mg/kg dry mass	9.5
	Co	mg/kg dry mass	3.2
	Cu	mg/kg dry mass	26.6
	Hg	mg/kg dry mass	0.3
	Pb	mg/kg dry mass	24.3
	Mo	mg/kg dry mass	1.4
	Sn	mg/kg dry mass	4.8
	Zn	mg/kg dry mass	22.3
	Ni	mg/kg dry mass	11.0




Fig. S1. Content (%) of individual components of petroleum pollutants in the soil: A) aliphatic hydrocarbons; B) PAHs ($n = 7–10$, $p < 0.05$).

Materials and Methods

1. Analysis of TPH and PAHs in soil and plant tissues

1.1 TPH Extraction and Chromatographic Analysis

The extraction of total petroleum hydrocarbons (TPH) from soil samples was performed using ultrasonic-assisted solvent extraction. A 5 g portion of air-dried soil was weighed into a 200 mL Erlenmeyer flask, followed by the addition of 20 mL of dichloromethane (Chempur, Poland). The extraction process was repeated on three occasions, with each extraction lasting 15 minutes, utilising an ultrasonic bath (Sonoswiss SW 6H, Sonoswiss, Switzerland). Following extraction, the combined extracts were purified by solid-phase extraction (SPE) using Chromabond Florisil columns (No. 730081, Macherey-Nagel, Germany). The eluates were subsequently concentrated using a rotary evaporator (Chemland, Poland). The final extracts were analysed using gas chromatography (GC) with a Clarus 500 gas chromatograph equipped with a flame ionisation detector (FID) (Perkin Elmer, USA). The chromatographic separation was performed on a fused silica capillary column (RTX-1, 30 m × 0.53 mm; Restek, Bellefonte, PA, USA). The GC operating conditions were as follows: injector temperature, 290°C; detector (FID) temperature, 320°C. The sequence of events that occurred during the experimental process is outlined below. Firstly, the initial isothermal hold was initiated at 30°C for a period of two minutes. This was followed by a temperature ramp that increased from 30°C to 105°C at a rate of 10°C/min. The subsequent ramp increased the temperature from 105°C to 285°C at a rate of 5°C/min. Finally, the oven was left in a final isothermal hold at 285°C for a duration of five minutes (Wojtowicz et al. 2022, Wojtowicz et al. 2023) .

1.2 PAHs Extraction and Chromatographic Analysis

Polycyclic aromatic hydrocarbons (PAHs) were extracted from soil samples using a modified QuEChERS method, as described by Wojtowicz (2022). Dried soil samples were placed into 50 mL vials, followed by the addition of

5 mL of distilled water, 10 mL of acetonitrile (Chempur, Poland), and a mixture of extraction salts containing magnesium sulfate (MgSO4) and sodium chloride (NaCl) (SST640, Interchim, France).

The vials were hermetically sealed and subjected to vigorous shaking for a period of five minutes. Subsequently, the samples were subjected to centrifugation at 3500 rpm for 10 minutes using an MPW-223e centrifuge (MPW, Poland). The resulting mixture was then purified using a Clean-up Kit (No. JO3937, Interchim, France), and subsequently filtered through a 0.2 µm nylon syringe filter.

The purified extracts were analysed using a Vanquish Core liquid chromatograph (Thermo Scientific, USA). The chromatographic separation was carried out utilising a NUCLEODUR C18 PAH column (125 × 4 mm, 3 µm; Macherey-Nagel, Düren, Germany). The detection process was conducted utilising a UV-VIS detector in conjunction with a fluorescence detector (FLD) that was equipped with an automatic sampler. The injection volume was 10 µL.

The mobile phases employed in this study were eluent A, composed of methanol (70% purity, obtained from POCH S.A., Poland) and eluent B, which was acetonitrile (purity 99%, obtained from POCH S.A., Gliwice, Poland). The flow rate was set to 1.5 mL/min. The gradient program is outlined as follows: The initial phase of the process involved a linear increase from 20% to 50% B over a period of 1.5 minutes, followed by a transition to 100% B over 1 minute. The process was then maintained at 100% B for 1 minute, after which it was returned to 0% B over 3 minutes. The final stage involved a final hold at 100% A for 3 minutes (Wojtowicz 2022, Wojtowicz et al. 2023).

2.Ecotoxicological analyses

2.1 Phytotoxkit™

The phytotoxicity of the tested soil samples was evaluated using the Phytotoxkit™ chronic toxicity test (MicroBioTests Inc.), based on the assessment of seed germination and early root growth inhibition. The test was conducted in accordance with the standardised procedure delineated in ISO 18763, employing three plant species that were selected on the basis of their rapid and sensitive growth responses: the monocotyledonous Sorghum saccharatum (sorghum) and the two dicotyledonous species Lepidium sativum (garden cress) and Sinapis alba (white mustard).

The experiment was conducted in triplicate for each species, employing transparent polystyrene test plates to facilitate direct observation and measurement of root length through image analysis. The incubator was maintained under controlled conditions, with a temperature of 25°C and a lack of light, for a period of 72 hours. Following the conclusion of the exposure period, the inhibition of root elongation and germination was determined as indicators of phytotoxicity (Mamindy-Pajany et al., 2011; Blinova et al., 2012; Baran and Tarnawskiet al., 2013).

2.2 Ostracodtoxkit FTM

The Ostracodtoxkit F™ is a six-day chronic toxicity microbiotest based on mortality and growth inhibition of the benthic crustacean Heterocypris incongruens. This test is a preliminary bioassay for contaminated sediments and soils, and it complies with the ISO 14371:2012 standard. The assay utilises young *Heterocypris incongruens* that have been hatched from dormant cysts within approximately 52 hours. The organisms are exposed to the sediment under investigation in a six-well plate configuration (three wells by two wells) at a temperature of 25°C for a

period of six days. At the conclusion of the exposure period, mortality and growth inhibition are measured in relation to a non-toxic control sediment to assess toxicity (Rouvalis et al., 2013; Niyommaneerat et al., 2013)

2.3 Microtox®SPT– Acute Toxicity Test

The Microtox® Solid Phase Test (SPT) was utilised in order to assess the toxicity of the samples. This test is based on the analysis of the luminescent activity of *Vibrio fischeri* bacteria. Lyophilized *Vibrio fischeri* strains used in the study were supplied by SDI (USA). The bacteria were stored at a temperature of –20°C and subsequently reactivated in deionized water prior to use, in accordance with the manufacturer's instructions. Under standard metabolic conditions, approximately 10% of the bacterial energy budget is allocated to the process of photosynthesis. The intensity of this bioluminescence is proportional to the metabolic activity of the cells and can be accurately measured using a photometer. The phenomenon of bioluminescence is the result of the activity of the enzyme luciferase (also known as alkane oxygenase), which functions as a catalyst in the oxidation of reduced compounds (such as FMNH2, riboflavin phosphate, or FADH2) within the electron transport chain. This process occurs in the presence of molecular oxygen and a long-chain aldehyde. This reaction results in the emission of visible light. The presence of toxic substances, which interfere with bacterial metabolism, has been shown to cause a rapid decrease in luminescence. These alterations can be discerned following a 15-minute exposure to the sample, and are subsequently documented through the utilisation of the DeltaTox® analyser. The Microtox® SPT facilitates direct contact between the bacteria and solid-phase samples, enabling the detection of both water-soluble substances and poorly soluble or lipophilic compounds. This enhancement is of particular relevance to environmental assessments, particularly in the context of evaluating soil contamination. Toxicity measurements were conducted in accordance with the standard test protocol, and the results were expressed as EC_{50} values, defined as the concentration of the tested sample causing a 50% reduction in luminescence compared to the control. (Lima et al.. 2010; Foucaulty et al.. 2013; Oleszczuk et al.. 2014).

2.4MARA (Microbial Assay for Risk Assessment)

The MARA test (Microbial Assay for Risk Assessment) is a standardized microbiological assay used to evaluate the chronic toxicity and environmental risk of chemical substances and environmental samples. This method employs a multispecies bioassay format incorporating eleven microbial strains, of which ten are prokaryotic organisms (bacteria from diverse taxonomic groups) and one is an eukaryotic organism (yeast). The bacterial strains utilised in this study encompass *Microbacterium* sp., *Brevundimonas diminuta, Citrobacter freundii, Comamonas testosteroni, Enterococcus casseliflavus, Delftia acidovorans, Kurthia gibsoni, Staphylococcus warneri, Pseudomonas aurantiaca*, and *Serratia rubidaea*. The eukaryotic strain employed in this study is the yeast Pichia anomala. The test is conducted using a pre-prepared 96-well polystyrene microplate supplied by the manufacturer, in which lyophilized microorganisms are individually placed in each well under aseptic conditions. The test organisms are exposed to the test sample, which has been diluted in a nutrient medium, and then incubated for 18 hours at a controlled temperature. The toxicity of a substance is determined by the extent to which it inhibits microbial growth. This is assessed by measuring the reduction of a redox-sensitive dye (tetrazolium red). The presence of healthy, metabolically active microorganisms has been shown to reduce the dye, leading to the formation of an insoluble red precipitate that accumulates at the bottom of the wells. The intensity

of this red colouration is directly proportional to microbial activity and inversely proportional to the toxicity of the tested sample.Subsequent to the incubation period, the microplate is scanned using a flatbed scanner, and the resulting image is analysed using dedicated MARA software. The software quantitatively evaluates the colour development in each well, calculates the percentage inhibition of growth for each strain, and determines the EC50 values – that is, the effective concentration of the sample that causes 50% inhibition of microbial growth. The calculation of individual EC_{50} values is performed independently for each test strain. The mean toxicity concentration (MTC) is derived from these results, representing the average EC50 across all eleven organisms. In addition to providing an overall toxicity value, the MARA test generates a characteristic toxicity profile (fingerprint) based on the differential sensitivity of each strain. This allows for more detailed interpretation of toxic effects and potential identification of toxicant classes.The assay was performed in accordance with the standardised protocol that has been described in a number of previous studies (Gabrielson, 2004; Wadhia et al., 2007; Fau et al., 2010; Stefaniuk et al., 2015; Paskuliakowa et al., 2018).

3 DNA extraction, bacterial 16S rRNA gene Illumina MiSeq-based sequencing, and sequence processing

The DNA isolation, 16S sequencing and analysis were performed as described previously (Brzeszcz et al. 2020). Total DNA was extracted using Genomic Mini AX Bacteria kit according to the manufacturer's protocol. Additional mechanical lysis with zirconium beads was performed using FastPrep-24 instrument (MP Biomedicals). Next, the DNA from each replicate was pooled and then purified using and Anty-Inhibitor Kit (A&A Biotechnology, Poland). The DNA quality and concentration were verified by electrophoresis and fluorometry (Qubit 4 Fluorometer, Thermofisher Scientific, USA). Library preparation and sequencing was performed by Genomed S.A. (Warsaw, Poland). The library was prepared for the V3-V4 16S rDNA region amplified using 341F and 785R primers (Klindworth et al. 2013). A polymerase chain reaction (PCR) was carried out using Q5 Hot Start High-Fidelity 2x Master Mix (New England Biolabs, USA). An indexing PCR was performed using the Nextera XT index kit (Illumina). Next, the samples were purified using AMPure XP beads, verified in a bioanalyser (Agilent Technologies, USA) and with qPCR. The libraries were sequenced on a MiSeq platform (Illumina) with a MiSeq Reagent kit v3 in 2 x 300 bp paired-end technology. Automatic analysis and de-multiplexing of the raw reads were performed on MiSeq with the use of a MiSeqReporter (MSR) ver. 2.6. The DADA2 pipeline (ver. 1.22.0; Callahan et al. 2016) was then used in R (version 4.1.1; R Core Team, 2016) to recover the amplicon sequence variants (ASVs) from the data. The reads were quality-filtered and trimmed using the function filterAndTrim with the following parameters; truncLen = c (295, 205), trimLeft c = (17,21), maxN = 0, maxEE = c (4,7), truncQ = 2. The error rates were estimated using learnErrors, sequences were dereplicated using derepFastq with default parameters and exact sequence variants were resolved using dada. Chimeric sequences were removed using remove Bimera Denovo function, employing the "consensus" method. The identified chimeras were removed. The taxonomy was assigned to ASVs using the Naïve Bayesian classifier on the DADA2-formatted reference SILVA 16S rRNA database ver. 138 (Quast et al., 2012) with the default minimum bootstrap confidence. Multiple sequence alignment of the ASVs were performed using DECIPHER ver. 2.22.0 (Wright et al. 2015), and a phylogenetic tree was constructed using phangorn package ver. 2.11.1 (Schliep 2011). The frequency table, taxonomy assignment and phylogenetic tree information were used to create a phyloseq object, and bacterial community analyses were performed using phyloseq R package ver. 1.38.0 (McMurdie et al. 2013).

Table S2 Identification of bacterial strains used in microbial consortium

Stain (NCBI Number)	The closest relative based on 16S rRNA accession number. (% of identity) *	The closest relative for which the genome sequence is available in NCBI GenBank. accession number. (% of Identity) *	Putative gene encoding for the enzymes catalysing the breakdown of hydrocarbons
Dietzia sp. IN118 (JAUMSR000000000)	*Dietzia* sp. LJ3 MG049763 99.93%	*Dietzia kunjamensis* 313 CP099712 99.78%	Alkane 1-monooxygenase, aromatic ring-hydroxylating dioxygenase subunit alpha
Gordonia sp. IN101 (KT923337)	*Gordonia* sp. Tm-B24 MT533993 99.63%	*Gordonia terrae* RL-JC02 CP049836 99.63%	Alkane 1-monooxygenase, pentachlorophenol monooxygenase, naphthalene 1,2-dioxygenase subunit alpha (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Mycolicibacterium frederiksbergense IN53 (JAUKNU000000000)	*Mycolicibacterium frederiksbergense* DSM 44346 NR_025393.1 99.58%	*Mycolicibacterium frederiksbergense* LB 501T 99.58%	Alkane 1-monooxygenase (2 copies), pentachlorophenol monooxygenase, naphthalene 1,2-dioxygenase subunit alpha (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Rhodococcus erythropolis IN119 (KT923331)	*Rhodococcus erythropolis* KD-1 CP050124 99.42%	*Rhodococcus erythropolis* KD-1 CP050124 99.42%	Alkane 1-monooxygenase (5 copies), pentachlorophenol monooxygenase, cyclohexanone monooxygenase (2 copies), biphenyl 2,3-dioxygenase (2 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies)
Rhodococcus globerulus IN113 (KT923347)	*Rhodococcus globerulus* D757 CP079698 99.86%	*Rhodococcus globerulus* D757 CP079698 99.86%	Alkane 1-monooxygenase (3 copies), 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies), aromatic ring-hydroxylating dioxygenase subunit alpha (7 copies)
Raoultella sp. IN109 (KT923339)	*Raoultella planticola* SCLZS62 CP082168 99.08%	*Raoultella planticola* SCLZS62 CP082168 99.08%	aromatic ring-hydroxylating dioxygenase subunit alpha (2 copies)
Pseudomonas sp. IN132 (KT923299)	*Pseudomonas citronellolis* PY-1 MH685460 99.49%	*Pseudomonas citronellolis* SJTE-3 CP015878 99.49%	Alkane 1-monooxygenase, phenol 2-monooxygenase, 2,4-dichlorophenol 6-monooxygenase, naphthalene 1,2-dioxygenase, biphenyl 2,3-dioxygenase, 2,3-dihydroxybiphenyl 1,2-dioxygenase

Table S3 Contents of selected groups of aliphatic hydrocarbons and polycyclic aromatic hydrocarbons in the initial soil (G6) and in the remediated soils.

Hydrocarbons	**Content ± SD (mg/kg dry mass soil)**					
		After 180 Days				
	Initial Soil G6	control soil G6X	soil SS-G6X-A	soil SS-G6X-B	soil CO-G6X-A	soil CO-G6X-B
TPH	4,320.73 ± 258	4,158.17 ± 228	455.05 ± 25	2,755.10 ± 108	261.13 ± 13	2,436.57 ± 95
Unidentified aliphatic hydrocarbons	2,049.43 ± 125	1,977.91 ± 118	120.37 ± 7	1,382.77 ± 81	48.87 ± 3	1,235.76 ± 73
$\sum nC_6–nC_8$	27.87 ± 1.6	26.06 ± 1.5	1.01 ± 0.06	13.62 ± 0.75	0.35 ± 0.02	11.68 ± 0.66
$\sum nC_9–nC_{21}$	1,129.59 ± 65	1,071.73 ± 62	47.00 ± 2.8	586.30 ± 34	16.86 ± 1.0	500.56 ± 29
$\sum nC_{22}–nC_{30}$	793.30 ± 47	768.84 ± 46	130.94 ± 8.0	514.08 ± 30	76.65 ± 4.6	454.21 ± 27
$\sum nC_{31}–nC_{40}$	225.02 ± 13	219.37 ± 13	80.75 ± 5.0	175.27 ± 10	46.61 ± 2.8	154.65 ± 9.2
PAHs	7.19 ± 0.35	7.00 ± 0.31	1.53 ± 0.08	4.89 ± 0.25	1.14 ± 0.06	4.31 ± 0.21
∑2 -ring PAHs	2.36 ± 0.14	2.28 ± 0.13	0.18 ± 0.01	1.37 ± 0.08	0.09 ± 0.01	1.14 ± 0.07
∑ 3 - ring PAHs	2.09 ± 0.12	2.03 ± 0.12	0.33 ± 0.02	1.35 ± 0.08	0.20 ± 0.01	1.17 ± 0.07
∑ 4- ring PAHs	1.93 ± 0.11	1.88 ± 0.11	0.62 ± 0.04	1.47 ± 0.09	0.50 ± 0.03	1.35 ± 0.08
∑ 5- ring PAHs	0.61 ± 0.04	0.60 ± 0.04	0.29 ± 0.02	0.52 ± 0.03	0.24 ± 0.02	0.47 ± 0.03
∑6-ring PAHs	0.20 ± 0.01	0.20 ± 0.01	0.11 ± 0.01	0.18 ± 0.01	0.11 ± 0.01	0.17 ± 0.01

Table S4 Explanation of abbreviations in Fig. 6B) Redundancy analysis of environmental factors (arrows) and microbial phyla (circles) in two dimensional space

No.	Abberation	Full name
1	*Abdiii*	*Abditibacteriota*
2	*Acidob*	*Acidobacteriota*
3	*Actinob*	*Actinobacteriota*
4	*Armati*	*Armatimonadota*
5	*Bactero*	*Bacteroidota*
6	*Bdell*	*Bdellovibrionota*
7	*Bacter*	*unidentified Bacteria*
8	*Caldatri*	*Caldatribacteriota*
9	*Caldi*	*Caldisericota*
10	*Campi*	*Campilobacterota*
11	*Chlor*	*Chloroflexi*
12	*Cren*	*Crenarchaeota*
13	*Cyano*	*Cyanobacteria*
14	*Deino*	*Deinococcota*
15	*Dependе*	*Dependentiae*
16	*Desulfo*	*Desulfobacterota*
17	*Elusim*	*Elusimicrobiota*
18	*Entoiheo*	*Entotheonellaeota*
19	*Eury*	*Euryarchaeota*
20	*Fibrob*	*Fibrobacterota*
21	*Firmic*	*Firmicutes*
22	*Gemma*	*Gemmatimonadota*
23	*Halo*	*Halobacterota*
24	*Hydrogen*	*Hydrogenedentota*
25	*Latescib*	*Latescibacterota*
26	*Methylom*	*Methylomirabilota*
27	*Myxococc*	*Myxococcota*
28	*Nitrosp*	*Nitrospirota*
29	*Pates*	*Patescibacteria*
30	*Planctom*	*Planctomycetota*
31	*Proteo*	*Proteobacteria*
32	*Spirocha*	*Spirochaetota*
33	*Sumerl*	*Sumerlaeota*
34	*Synergi*	*Synergistota*
35	*Therm*	*Thermoplasmatota*
36	*Verruco*	*Verrucomicrobiota*

Table S5 Species of ciliated protozoa found in the initial and treatments during the studies.

No.	Species	soil G6	soil SS-G6X-A	soil SS-G6X-B	soil CO-G6X-A	soil CO-G6X-B
1	*Blepharisma lateritium*			+	+	+
2	*Cinetochilum margaritaceum*		+	+	+	+
3	*Colpoda cucullus*	+		+		+
4	*Colpoda inflata*	+		+	+	+
5	*Ctedoctema acanthocryptum*				+	+
6	*Cyrtolophosis muscicola*	+	+	+	+	
7	*Enchelyodon* sp.			+		
8	*Enchelys* sp.			+		+
9	*Epispathidium amphoriforme*		+			+
10	*Gonostomum affine*	+	+	+	+	+
11	*Halteria grandinella*			+	+	
12	*Holosticha multistilata*	+	+	+	+	
13	*Holosticha pulaster*		+		+	
14	*Kahlilembus attenuatus*			+		
15	*Leptopharynx costatus*			+		+
16	*Metacystis* sp.			+		+
17	*Perisincirra* sp.		+	+		+
18	*Platyophrya vorax*			+	+	+
19	*Pseudochilodontopsis fluviatilis*	+			+	
20	*Sathrophilus muscorum*			+	+	+
21	*Spathidium spathula*		+	+		+
22	*Stichotricha aculeata*			+		
23	*Tachysoma pellionellum*	+	+	+	+	+
24	*Tetrahymena* sp.					+
25	*Uroleptus* sp.	+		+	+	
26	*Uroleptus musculus*	+	+	+	+	+
	Total number of species	9	10	21	15	17

Table S6 Time course of the most important ecological indices in the G6 and in the other treatments after 3. 6. 9 and 12 days.

	G6 day of analysis			
Index	**3**	**6**	**9**	**12**
Dominance_D	0.90	0.40	0.67	0.29
Simpson_1-D	0.10	0.60	0.33	0.71
Shannon_H	0.28	1.16	0.69	1.30
Evenness_e^H/S	0.26	0.53	0.50	0.92
Margalef	0.50	0.78	0.51	0.58
Equitability_J	0.17	0.64	0.49	0.94
	SS-G6X-A day of analysis			
Index	3	6	9	12
Dominance_D	0.78	0.17	0.43	0.54
Simpson_1-D	0.22	0.83	0.57	0.46
Shannon_H	0.45	1.87	0.97	0.82
Evenness_e^H/S	0.39	0.92	0.88	0.76
Margalef	0.44	1.28	0.51	0.56
Equitability_J	0.32	0.96	0.88	0.75
	SS-G6X-B day of analysis			
Index	3	6	9	12
Dominance_D	0.18	0.57	0.96	0.85
Simpson_1-D	0.82	0.43	0.04	0.15
Shannon_H	2.00	1.09	0.12	0.41
Evenness_e^H/S	0.67	0.27	0.16	0.17
Margalef	1.69	1.56	0.73	1.10
Equitability_J	0.83	0.45	0.06	0.19
	CO-G6X-A day of analysis			
Index	3	6	9	12
Dominance_D	0.24	0.30	0.30	0.43
Simpson_1-D	0.76	0.70	0.70	0.57
Shannon_H	1.70	1.29	1.41	1.27
Evenness_e^H/S	0.55	0.90	0.51	0.44
Margalef	1.43	0.56	1.11	1.07
Equitability_J	0.74	0.93	0.68	0.61
	CO-G6X-B day of analysis			
Index	3	6	9	12
Dominance_D	0.37	0.33	0.33	0.49
Simpson_1-D	0.63	0.67	0.67	0.51
Shannon_H	1.39	1.32	1.47	0.76
Evenness_e^H/S	0.50	0.75	0.44	0.54
Margalef	1.24	0.82	1.48	0.45
Equitability_J	0.67	0.82	0.64	0.55

References:

1. Baran, A., & Tarnawski, M. (2013). Phytotoxkit/Phytotestkit and Microtox® as tools for toxicity assessment of sediments. *Ecotoxicology and Environmental Safety, 98*, 19–27. https://doi.org/10.1016/j.ecoenv.2013.10.010
2. Blinova, I., Niskanen, J., Kajankari, P., Kanarbik, L., Käkinen, A., Tenhu, H., Penttinen, O. P., & Kahru, A. (2013). Toxicity of two types of silver nanoparticles to aquatic crustaceans Daphnia magna and Thamnocephalus platyurus. *Environmental Science and Pollution Research, 20*(5), 3456–3463.
3. Brzeszcz, J., Kapusta, P., Steliga, T., & Turkiewicz, A. (2020). Hydrocarbon removal by two differently developed microbial inoculants and comparing their actions with biostimulation treatment. *Molecules, 25*, 661. https://doi.org/10.3390/molecules25030661

4. Fai, P. B., & Grant, A. (2010). An assessment of the potential of the Microbial Assay for Risk Assessment (MARA) for ecotoxicological testing. *Ecotoxicology, 19*, 1626–1633. https://doi.org/10.1007/s10646-010-0548-2

5. Foucault, Y., Durand, M.-J., Tack, K., Schreck, E., Geret, F., Leveque, T., Pradere, P., Goix, S., & Dumat, C. (2013). Use of ecotoxicity test and ecoscores to improve the management of polluted soils: Case of a secondary lead smelter plant. *Journal of Hazardous Materials, 246–247*, 291–299. https://doi.org/10.1016/j.jhazmat.2012.12.042

6. Gabrielson, J. (2004). *Assessing the toxic impact of chemicals using bacteria*. Karolinska University Press: Stockholm. ISBN: 9789171401434.

7. Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O. (2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-generation sequencing-based diversity studies. *Nucleic Acids Research, 41*, e1–e1. https://doi.org/10.1093/nar/gks808

8. Mamindy-Pajany, Y., Hamer, B., Roméo, M., Géret, F., Galgani, F., Durmiši, E., Hurel, C., & Marmier, N. (2011). The toxicity of composted sediments from Mediterranean ports evaluated by several bioassays. *Chemosphere, 82*, 362–369. https://doi.org/10.1016/j.chemosphere.2010.10.005

9. McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLOS ONE, 8*(4), e61217. https://doi.org/10.1371/journal.pone.0061217

10. Niyommaneerat, W., Nakajima, F., Tobino, T., & Yamamoto, K. (2017). Development of a chronic sediment toxicity test using the benthic ostracod Heterocypris incongruens and their application to toxicity assessments of urban road dust. *Ecotoxicology and Environmental Safety, 143*, 266–274. https://doi.org/10.1016/j.ecoenv.2017.05.011

11. Oleszczuk, P., Jośko, I., Kuśmierz, M., Futa, B., Wielgosz, E., Ligęza, S., & Pranagal, J. (2014). Microbiological, biochemical and ecotoxicological evaluation of soils in the area of biochar production in relation to polycyclic aromatic hydrocarbon content. *Geoderma, 213*, 502–511. https://doi.org/10.1016/j.geoderma.2013.08.027

12. Paskuliakova, A., McGowan, T., Tonry, S., & Touzet, N. (2018). Phycoremediation of landfill leachate with the chlorophyte Chlamydomonas sp. SW15aRL and evaluation of toxicity pre and post treatment. *Ecotoxicology and Environmental Safety, 147*, 622–630. https://doi.org/10.1016/j.ecoenv.2017.09.010

13. Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research, 41*, D590–D596. https://doi.org/10.1093/nar/gks1219

14. Rouvalis, A., Theodoropoulos, C., & Georgudaki, J. I. (2013). Assessment of toxicity of the untreated and Pleurotus ostreatus treated olive mill wastewater by using microbiotests. *International Journal of Environmental Engineering, 5*, 373. https://doi.org/10.1504/IJEE.2013.056623

15. Schliep, K. P. (2011). phangorn: Phylogenetic analysis in R. *Bioinformatics, 27*, 592–593. https://doi.org/10.1093/bioinformatics/btq706

16. Stefaniuk, M., Bartmiński, P., Różyło, K., Dębicki, R., & Oleszczuk, P. (2015). Ecotoxicological assessment of residues from different biogas production plants used as fertilizer for soil. *Journal of Hazardous Materials, 298*, 195–202. https://doi.org/10.1016/j.jhazmat.2015.05.026

17. Wadhia, K., Dando, T., & Thompson, K. C. (2007). Intra-laboratory evaluation of microbial assay for risk assessment (MARA) for potential application in the implementation of the Water Framework Directive (WFD). *Journal of Environmental Monitoring, 9*, 953. https://doi.org/10.1039/b704059h

18. Wojtowicz, K. (2022). Opracowanie metodyki oznaczania WWA w próbkach gleb z wykorzystaniem chromatografii cieczowej HPLC. *Nafta-Gaz, 78*, 141–153. https://doi.org/10.18668/NG.2022.02.06

19. Wojtowicz, K., Steliga, T., Kapusta, P., & Brzeszcz, J. (2022). Evaluation of the effectiveness of the biopreparation in combination with the polymer γ-PGA for the biodegradation of petroleum contaminants in soil. *Materials, 15*, 400. https://doi.org/10.3390/ma15020400

20. Wojtowicz, K., Steliga, T., Kapusta, P., & Brzeszcz, J. (2023). Oil-contaminated soil remediation with biodegradation by autochthonous microorganisms and phytoremediation by maize (*Zea mays*). *Molecules, 28*, 6104. https://doi.org/10.3390/molecules28166104

21. Wright, E. S. (2015). DECIPHER: Harnessing local sequence context to improve multiple sequence alignment. *BMC Bioinformatics, 16*, 322. https://doi.org/10.1186/s12859-015-0749-z

molecules

MDPI

Article

The Role of Graphene Oxide and Zinc Oxide Nanoparticles in Enhancing the Effectiveness of Phytoremediation of Petroleum Hydrocarbon-Contaminated Soils Using *Lolium perenne*

Katarzyna Wojtowicz [1], Teresa Steliga [1,*], Piotr Kapusta [2] and Joanna Brzeszcz [2]

[1] Department of Reservoir Fluid Production Technology, Oil and Gas Institute–National Research Institute, Lubicz 25 A, 31-503 Krakow, Poland; katarzyna.wojtowicz@inig.pl
[2] Department of Microbiology, Oil and Gas Institute–National Research Institute, Lubicz 25 A, 31-503 Krakow, Poland; piotr.kapusta@inig.pl (P.K.); joanna.brzeszcz@inig.pl (J.B.)
* Correspondence: teresa.steliga@inig.pl

Abstract

Nanomaterials are gaining increasing importance in various scientific and technological fields, including ecological strategies for environmental remediation, such as the treatment of soils contaminated with petroleum hydrocarbons. This study aimed to evaluate the effectiveness of hydrocarbon-contaminated soil remediation using graphene oxide nanoparticles (GO-NPs) and zinc oxide nanoparticles (ZnO-NPs) in combination with bacterial consortium inoculation and phytoremediation with *Lolium perenne*. The study was conducted in two stages: laboratory-scale biodegradation experiments and semi-technical scale phytoremediation. The laboratory stage determined optimal nanomaterial doses based on respirometric and chromatographic analyses. During phytoremediation, the contents of total petroleum hydrocarbons (TPHs) and polycyclic aromatic hydrocarbons (PAHs) in soil, roots, and shoots were monitored. Biomass growth was recorded, and environmental toxicity was assessed using Phytotoxkit, Microtox, and Ostracodtoxkit tests. The addition of nanomaterials significantly enhanced soil remediation, with improvements in TPH and PAHs removal resulting from microbial biodegradation in both stages and, additionally, in the phytoremediation stage, from phytoextraction by plants. TPH biodegradation reached up to 81.85% in GO-NP variants and 80.9% in ZnO-NP treatments, while PAHs reached 73.19% and 70.66%, respectively. The biomass of *Lolium perenne* increased by 28.63% in GO-NP variants and by 22.21% in ZnO-NP treatments compared to the control. Total accumulation of TPH increased by 80.86% and 74.15%, and PAHs by 71.26% and 65.35%. Nanomaterial-amended variants also showed a reduction in toxicity units to 2.50–3.30. These results indicate that combining nanomaterials with bioaugmentation significantly enhances phytoremediation efficiency while reducing soil toxicity.

Keywords: bacterial consortium; biodegradation; ecotoxicological assessment; graphene oxide nanoparticles (GO-NPs); *Lolium perenne*; petroleum hydrocarbons; phytoremediation; zinc oxide nanoparticles (ZnO-NPs)

Check for updates

Academic Editors: Kunlei Wang and Zhishun Wei

Received: 30 January 2026
Revised: 4 March 2026
Accepted: 5 March 2026
Published: 7 March 2026

Copyright: © 2026 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license.

1. Introduction

Bioremediation is currently one of the most important and frequently used methods of remediation for soils contaminated with petroleum hydrocarbons as a result of industrial, mining, or transport activities [1,2]. The utilisation of natural biological processes involving microorganisms and plants constitutes the basis of these methods. The transformation or

elimination of pollutants is thereby enabled, whilst simultaneously reducing interference with the soil environment and the costs of reclamation [3]. One of the fundamental mechanisms of bioremediation is biodegradation, which refers to the process of the enzymatic decomposition of petroleum hydrocarbons by soil microorganisms. This process results in the formation of compounds with reduced toxicity, and under optimal conditions, these compounds can undergo complete mineralisation [4,5]. The effectiveness of this process is contingent on numerous factors, including the composition and activity of soil microbiota, the availability of oxygen and nutrients, and the bioavailability of pollutants [6]. In the case of petroleum hydrocarbons, which are hydrophobic in nature, this bioavailability is often limited due to their strong sorption to the soil matrix, which significantly slows down biodegradation processes [7,8]. A complementary approach to bioremediation is phytoremediation, which harnesses the capacity of plants and rhizosphere microorganisms to catalyse the degradation, transformation, or immobilisation of organic pollutants [9,10]. The rhizosphere plays a pivotal role in this process, where intense biological activity induced by root exudates promotes the development of populations of microorganisms capable of decomposing hydrocarbons [11,12]. Notwithstanding the numerous advantages associated with phytoremediation of soils contaminated with petroleum compounds, the effectiveness of this process is sometimes constrained by several factors. These include the sluggish pace of the processes involved, the limited spatial range of influence, which is primarily confined to the surface layer of the soil, and the limited bioavailability of hydrophobic fractions of pollutants [13]. In the field of phytoremediation, particular emphasis is placed on grass species, which are characterised by rapid growth and well-developed, dense root systems [14–16]. One species frequently identified as having potential in the remediation of soils contaminated with petroleum hydrocarbons is *Lolium perenne* [17]. This species demonstrates high tolerance to environmental stress, alongside the capacity for substantial biomass production. Furthermore, it exhibits effective densification of the rhizosphere zone, which enhances soil microbial activity and intensifies biodegradation processes [18,19]. However, even with the incorporation of *Lolium perenne*, the effectiveness of phytoremediation may be inadequate without supplementary technological support. In order to enhance the efficacy of biodegradation and phytoremediation, there has been an increasing tendency to modify conventional bioremediation methods by employing integrated approaches [20–22]. The most common methods employed include bioaugmentation with biopreparations or microbial consortia containing selected strains of microorganisms capable of decomposing petroleum-derived hydrocarbons [22,23], as well as the addition of surfactants [21,24], which increase the solubility, mobility, and bioavailability of pollutants in the soil environment. In recent years, however, the use of nanomaterials as an innovative tool to support biological processes has attracted particular interest [25,26]. Of particular interest are nanoparticles of metals and their oxides, such as zinc oxide (ZnO) [27,28], magnetite (Fe_3O_4) [29,30], hematite (α-Fe_2O_3), maghemite (γ-Fe_2O_3) [31], silicon (IV) oxide (SiO_2), aluminium oxide (Al_2O_3) [32], and titanium oxide (TiO_2) [32–34], as well as modern carbon materials [35], including graphene [36,37], graphene oxide [38,39], carbon nanotubes (CNTs) [40,41], and various types of nanopolymers [42]. These materials exhibit unique physicochemical properties, such as a very large specific surface area, high reactivity, and the ability to adsorb and catalyse the transformation of organic compounds [43–45], which enables their effective use in removing a wide range of environmental pollutants [46]. Nanomaterials can enhance the efficacy of biodegradation and phytoremediation processes through the multiple functions they perform. A significant mechanism of their action is the enhancement of the bioavailability of hydrophobic pollutants by desorbing them from the soil matrix or transforming them into forms that are more readily available to rhizosphere microorganisms and plant root systems [47,48]. It has been demonstrated that metal and

https://doi.org/10.3390/molecules31050890

metal oxide nanoparticles can catalyse the transformation of petroleum-derived hydrocarbons, thereby facilitating their subsequent biodegradation [49,50]. Moreover, nanomaterials can bind toxic substances, thereby reducing their phytotoxicity and protecting plants from the deleterious effects of pollution [50,51]. Furthermore, nanomaterials have been shown to enhance plant growth and development, increase resilience to environmental stress, and stimulate natural defence mechanisms [48,50]. Numerous studies indicate that they support root system development, improve soil structure and aeration, and intensify the enzymatic activity of both plants and soil microorganisms [52,53]. When appropriately engineered, nanomaterials can also function as carriers of nutrients, growth promoters, or other bioactive substances, thereby supporting organism performance under contaminated conditions [54,55]. A particularly promising area of research is nanomaterial-assisted bioaugmentation, which involves the simultaneous introduction into the soil of selected microorganisms capable of decomposing specific hydrocarbon fractions and nanomaterials that increase their survival and metabolic activity under adverse environmental conditions [56,57]. Such a synergistic combination can significantly accelerate the remediation of soils contaminated with petroleum-derived substances. Nevertheless, despite the promising outcomes of previous studies, the application of nanomaterials in environmental remediation requires further in-depth investigation. Reports indicate potential risks related to their persistence in the environment and their long-term effects on living organisms [49,58]. Therefore, a comprehensive evaluation of biodegradation and phytoremediation assisted by bioaugmentation and nanomaterials should include not only the efficiency of hydrocarbon removal but also a thorough ecotoxicological assessment of the soil following the remediation process. Building on this, the primary aims of the present study were to develop effective methods for the remediation of soils contaminated with petroleum hydrocarbons, to investigate the influence of nanomaterials on remediation efficiency, and to enhance the overall effectiveness of biodegradation and phytoremediation strategies.

2. Results

2.1. Biodegradation Studies–Laboratory Scale

2.1.1. Respirometric Tests

Biodegradation tests were conducted under laboratory conditions using the OxiTop respirometric system. This system enables the assessment of the intensity of biodegradation processes indirectly, by means of measuring oxygen consumption by microorganisms. An increase in microbial activity in the reaction environment is indicative of microorganisms effectively utilising the carbon contained in aliphatic molecules and polycyclic aromatic hydrocarbons as a source of energy and building material. Increased oxygen consumption is a direct consequence of intensified metabolic processes and is directly related to the intensification of the biodegradation of petroleum-derived substances (Figure 1).

Respirometric analyses demonstrated that the inoculation of soil with a biopreparation, in conjunction with the incorporation of nanomaterials, exerted a substantial influence on the progression of biodegradation processes. In the control sample, which comprised soil inoculated with a biopreparation devoid of nanomaterials, oxygen consumption of 1944.8 mg O_2/dm^3 was documented following 90 days of respirometric testing. This finding signifies moderate activity of microorganisms responsible for the degradation of petroleum-derived hydrocarbons. The incorporation of graphene oxide nanoparticles (GO-NPs) into the reaction systems resulted in a significant increase in oxygen consumption, with this effect strongly dependent on the nanomaterial dosage (Figure 1a). For a dose of 10 mg GO-NP/kg dry mass (DM), oxygen consumption increased to 4304.8 mg O_2/dm^3, indicating a significant stimulation of microbial activity. The highest oxygen consumption value, 5147.8 mg O_2/dm^3, was observed for a dose of 20 mg GO-NP/kg DM, reflecting

https://doi.org/10.3390/molecules31050890

the most pronounced biodegradation processes within the examined system. Further increases in the dose of GO-NPs led to a decrease in microbial activity. At an initial concentration of 50 mg GO-NP/kg DM, oxygen consumption decreased to 4840.2 mg O_2/dm^3, while at the highest dose of 100 mg GO-NP/kg DM, this value dropped to 2244.1 mg O_2/dm^3. The observed decrease in oxygen consumption may indicate a toxic effect of nanomaterials at high concentrations, leading to inhibition of microbial activity. These findings suggest that GO-NPs, when administered in low and moderate doses, exert a stimulatory effect on biodegradation processes. Conversely, excessive GO-NPs may have an inhibitory effect. A similar relationship between nanomaterial dose and biodegradation intensity was observed for zinc oxide nanoparticles (ZnO-NPs) (Figure 1b). The incorporation of 10 mg ZnO-NP/kg DM led to an increase in oxygen consumption, reaching 3885.1 mg O_2/dm^3. A dose of 20 mg ZnO-NP/kg DM yielded 4836.6 mg O_2/dm^3, confirming the favourable impact of moderate ZnO-NP doses on microorganisms involved in petroleum hydrocarbon degradation. As ZnO-NP concentrations increased, a gradual decrease in microbial respiration was observed. At a dose of 50 mg ZnO-NP/kg DM, oxygen consumption was 4102.9 mg O_2/dm^3, whereas at 100 mg ZnO-NP/kg DM, it fell to 2025.3 mg O_2/dm^3. In systems containing 50 mg ZnO-NP/kg DM, a slight downward trend in oxygen consumption was observed during the final stage of the respirometric test, suggesting a gradual weakening of microbial activity over time, likely due to the cumulative toxic effect of the nanomaterial.




Figure 1. Results of respirometric analysis of oxygen consumption over time for soil sample G6 inoculated with a bacterial consortium and enriched with nanomaterials: (**a**) graphene oxide nanoparticles (GONPs) and (**b**) zinc oxide nanoparticles (ZnO-NPs) at different concentrations of these nanomaterials, (repetition number $n = 5$, $p < 0.05$).

2.1.2. Chromatographic Analysis

Chromatographic analyses were utilised to identify and quantitatively determine aliphatic hydrocarbons in soil contaminated with petroleum-derived substances, both before and after a 90-day biodegradation test. The initial total petroleum hydrocarbon (TPH) concentration in the tested soil was 6062.59 mg/kg (DM). In the control variant, where only bacterial consortium inoculation was applied (without nanomaterials), this concentration

https://doi.org/10.3390/molecules31050890

decreased to 3381.05 mg/kg DM, corresponding to a biodegradation of 44.23%. This outcome confirms the ability of the bacterial consortium to initiate biodegradation processes, while also highlighting the limited efficiency of aliphatic hydrocarbon decomposition in the absence of supplementary supporting factors. The application of GO-NPs in combination with bacterial inoculation significantly enhanced the biodegradation of aliphatic hydrocarbons. In the sample containing 10 mg GO-NP/kg DM, TPH concentration decreased to 2390.32 mg/kg DM, corresponding to 60.57% biodegradation. The highest decomposition efficiency was observed at a dose of 20 mg GO-NP/kg DM, where TPH content decreased to 1790.35 mg/kg DM, and the degree of biodegradation reached 70.47%. Further increases in GO-NP dosage led to reduced process efficiency. At 50 mg GO-NP/kg DM, biodegradation was 65.13%, whereas at 100 mg GO-NP/kg DM, it decreased to 50.41%. This trend indicates that excessive GO-NPs may inhibit microbial activity, potentially due to disruption of soil gas exchange, adsorption of nutrients, or adverse changes in the microstructure of the soil environment. These observations are consistent with respirometric analysis results.

A detailed chromatographic analysis of individual aliphatic hydrocarbon groups revealed that biodegradation efficiency strongly depended on both carbon chain length and the applied GO-NP dose (Figure 2a). In the control sample, the highest degradation was observed for light n-alkane fractions (n-C6–n-C8 and n-C9–n-C21), whereas heavier fractions (n-C22–n-C30 and n-C31–n-C36) exhibited markedly lower biodegradation. The incorporation of GO-NPs improved the biodegradation of all analysed hydrocarbon groups. Even at 10 mg GO-NP/kg DM, the degradation of each fraction increased, confirming that GO-NPs enhance pollutant bioavailability and support microbial activity. The highest biodegradation efficiency of individual fractions was observed at 20 mg GO-NP/kg of DM. Short- and medium-chain n-alkanes (n-C6–n-C21) underwent 81.57–99.85% biodegradation, while heavier hydrocarbons (n-C22–n-C30) exhibited over 66.58%. Even the most recalcitrant fraction (n-C31–n-C36) decreased by 40.68–48.69%, demonstrating the capacity of microorganisms to decompose more stable and branched compounds in the presence of an optimal dose of GO-NPs. Increasing the GO-NP dose above 20 mg/kg (50 and 100 mg/kg variants) reduced biodegradation efficiency across all the analysed fractions. At 50 mg/kg, the process remained relatively active, but a downward trend was already apparent, whereas the highest GO-NP concentration led to a marked reduction in microflora activity. This may result from the agglomeration of GO-NPs, restricting hydrocarbon availability for microorganisms or exerting toxic effects on bacterial cells.

Furthermore, soil variants inoculated with a bacterial consortium and enriched with ZnO-NP also demonstrated a significant effect of the nanomaterial on the biodegradation efficiency of aliphatic hydrocarbons (Figure 2b). The incorporation of 10 mg ZnO-NP/kg DM resulted in a significant enhancement in the degree of TPH biodegradation, reaching 58.30%, with a final concentration of 2527.89 mg/kg DM. The optimal efficiency of the process was attained at a dose of 20 mg of ZnO-NP/kg DM, where a substantial reduction in TPH content to 1959.53 mg/kg DM was observed, corresponding to a biodegradation degree of 67.68%. As in the case of GO-NPs, further increases in ZnO-NP concentration (50–100 mg/kg DM) resulted in a decrease in biodegradation efficiency, namely to 61.51% at 50 mg/kg DM and 49.69% at 100 mg/kg DM. These results indicate that excessive amounts of ZnO-NP in soil may limit microbial activity, potentially due to the partial toxicity of nanoparticles or disturbances in nutrient availability. The analysis of individual hydrocarbon fractions confirmed a marked increase in biodegradation efficiency across all groups in the presence of ZnO-NP compared to the control sample. In the variant with 20 mg ZnO-NP/kg DM, light n-alkanes (n-C6–n-C8) were biodegraded in the range of 88.79–89.33%, medium-chain n-alkanes(n-C9–n-C21) in 78.34–94.12%, heavier hydrocarbons (n-C22–n-C30) in the range of 63.95–71.53%, and the heaviest n-alkanes (n-C31–n-C36)

https://doi.org/10.3390/molecules31050890

in the range of 39.07–43.45%. An increase in ZnO-NP concentration above 20 mg/kg DM resulted in a gradual decrease in the biodegradation efficiency of all analysed fractions. At a dose of 50 mg/kg DM, the decomposition of light and medium fractions remained relatively intense; however, for heavier hydrocarbons, the decline in efficiency was more pronounced. The highest ZnO-NP concentration (100 mg/kg DM) led to a significant reduction in microbial activity, and the biodegradation degree of most fractions approached values observed in the control sample. As shown in Figure 2, the mean degree of biodegradation of selected groups of aliphatic hydrocarbons was determined for soil sample G6 inoculated with a bacterial consortium and enriched with nanomaterials.




Figure 2. Average degree of biodegradation of selected aliphatic hydrocarbon groups in soil sample G6 inoculated with a bacterial consortium and enriched with nanomaterials: (**a**) graphene oxide nanoparticles (GO-NPs) and (**b**) zinc oxide nanoparticles (ZnO-NPs) at different concentrations of these nanomaterials. Values are means ± SD ($n = 7$–10). Different lowercase letters indicate significant differences among treatments according to one-way ANOVA followed by Tukey's HSD test ($p < 0.05$).

As demonstrated by HPLC chromatographic analyses, the total initial concentration of polycyclic aromatic hydrocarbons (PAHs) in the tested soil sample was 12.05 mg/kg DM. In the control sample, in which inoculation was performed with the biopreparation alone, the PAHs content decreased to 7.60 mg/kg DM, corresponding to a biodegradation rate of 36.98%. This result indicates that the microorganisms contained in the bacterial consortium were capable of partially degrading aromatic compounds; however, the efficiency of this process was clearly limited, which confirms high persistence and low bioavailability of PAHs in the soil environment. The incorporation of GO-NPs at various doses resulted in a substantial enhancement of the effectiveness of PAHs biodegradation compared with the control sample. At a dose of 10 mg GO-NP/kg DM, the PAHs concentration decreased to 6.32 mg/kg DM, which corresponded to 47.56% biodegradation. The highest decomposition efficiency was observed in the variant with 20 mg GO-NP/kg DM, where the PAHs content decreased to 5.28 mg/kg DM, and the degree of biodegradation reached 56.19%. As was the case for aliphatic hydrocarbons, a further increase in the GO-NP dose resulted in a decrease in process effectiveness. At a dosage of 50 mg GO-NP/kg DM, the biodegradation rate was 52.55%, whereas at the highest dose of 100 mg GO-NP/kg DM,

https://doi.org/10.3390/molecules31050890

it decreased to 40.48%. The results demonstrate that moderate concentrations of GO-NPs promote the activity of microorganisms responsible for PAHs degradation. Conversely, excessive concentrations of GO-NPs may limit the biodegradation process, likely due to disturbances in environmental conditions or adverse effects of the nanomaterial on soil microbiota. An analysis of PAHs, with respect to the number of aromatic rings, enabled the assessment of the differential susceptibility of individual compound groups to biodegradation. The highest degree of degradation was observed for the two-ring naphthalene, which can be attributed to its lower chemical stability and greater bioavailability. In the control sample, naphthalene biodegradation reached 45.04%, whereas in the presence of 10 mg GO-NP/kg DM, this value increased to 57.65%. The maximum biodegradation efficiency of naphthalene was achieved at a dose of 20 mg GO-NP/kg DM, reaching 68.01%. Application of higher GO-NP doses (50 and 100 mg/kg DM) resulted in a decline in biodegradation efficiency to 63.95% and 48.44%, respectively. For three-ring PAHs, the degree of biodegradation increased from 33.22 to 35.23% in the control sample to 51.16–53.20% at a dose of 20 mg GO-NP/kg DM. For four-ring compounds, biodegradation increased from 26.11 to 28.36% to 40.42–42.85%. Five-ring PAHs were degraded to a lesser extent, with biodegradation ranging from 19.47 to 19.51% in the control sample to 30.42–31.47% in the sample with 20 mg GO-NP/kg DM. The most stable six-ring aromatic compounds exhibited an increase in biodegradation from 16.80 to 17.98% in the control sample to 25.37–26.14% at a dose of 20 mg GO-NP/kg DM. These findings indicate that GO-NP promoted the degradation of both lighter and more resistant PAHs fractions, although the effectiveness of the process decreased with increasing aromatic ring number. The addition of ZnO-NP also significantly affected the efficiency of PAHs biodegradation in soil. At a dose of 10 mg ZnO-NP/kg DM, the PAHs concentration decreased to 6.53 mg/kg DM, corresponding to 45.80% biodegradation. The most favourable results were obtained at a dose of 20 mg ZnO-NP/kg DM, where the PAHs content decreased to 5.43 mg/kg DM, and the biodegradation degree reached 54.94%. Increasing the ZnO-NP dose to 50 mg/kg DM led to a decrease in biodegradation efficiency to 48.38%, while at 100 mg/kg DM, it declined to 40.45%. This trend suggests that, similarly to GO-NP, excessive concentrations of ZnO-NP may limit microbial activity, potentially due to toxic effects or reduced bioavailability of aromatic compounds. Among all analysed PAHs, two-ring naphthalene exhibited the highest degree of degradation. In the variant with 20 mg ZnO-NP/kg DM, its biodegradation reached 65.96%, whereas at a dose of 10 mg/kg DM, it was 55.34%. Increasing the ZnO-NP dose to 50 mg/kg DM resulted in a decrease in degradation efficiency to 58.38% and to 49.52% at 100 mg/kg DM. Three-ring PAHs also showed a clear improvement in biodegradation in the presence of ZnO-NP, ranging from 41.85% to 43.29% at a dose of 10 mg/kg and from 50.62% to 53.60% at 20 mg/kg DM. Higher doses of the nanomaterial reduced degradation efficiency to 44.07–47.22% at 50 mg/kg DM and to 35.73–37.92% at 100 mg/kg DM. Four-ring PAHs, characterised by greater chemical stability, underwent biodegradation in the range of 34.60–34.85% at 10 mg ZnO-NP/kg DM, and 39.91–42.53% in the optimal dose. At higher concentrations, the degree of degradation decreased to 34.68–36.52% and 28.78–30.57%. Five-ring PAHs exhibited limited susceptibility to biodegradation. The highest degradation degree, 29.51–30.52%, was observed at 20 mg ZnO-NP/kg DM, whereas at doses of 10, 50, and 100 mg/kg DM the values ranged from 23.92 to 24.98%, 24.80–25.71%, and 19.98–20.23%, respectively. The most resistant six-ring PAHs were biodegraded in the range of 24.71–25.36% in the variant with 20 mg ZnO-NP/kg DM, while at higher nanomaterial concentrations, the degradation degree decreased to 21.26–22.88% and 17.15–18.41%. The ring-profile analysis clearly indicates that moderate ZnO-NP doses enhance PAHs bioavailability and microbial degradation of both lighter and heavier, more stable aromatic compounds. In contrast, excessive nanomaterial concentrations reduce biodegradation

https://doi.org/10.3390/molecules31050890

efficiency, likely due to adverse effects on soil microbiota or limited substrate availability. Figure 3 presents the mean degree of biodegradation of PAHs with different numbers of aromatic rings in soil sample G6 inoculated with a bacterial consortium and enriched with nanomaterials. Figure 3 presents the mean degree of biodegradation of PAHs with different numbers of aromatic rings in soil sample G6 inoculated with a bacterial consortium and enriched with nanomaterials.




Figure 3. Average degree of biodegradation of selected polycyclic aromatic hydrocarbon groups in soil sample G6 inoculated with a bacterial consortium and enriched with nanomaterials: (**a**) graphene oxide nanoparticles (GO-NPs) and (**b**) zinc oxide nanoparticles (ZnO-NPs) at different concentrations of these nanomaterials. Values are means ± SD (n = 7–10). Different lowercase letters indicate significant differences among treatments according to one-way ANOVA followed by Tukey's HSD test ($p < 0.05$).

2.2. Phytoremediation Studies–Semi-Technical Scale

2.2.1. Chromatographic Analysis

A study of the degree of TPH biodegradation under phytoremediation conditions using *Lolium perenne* demonstrated clear variations in the effectiveness of the process depending on the soil enrichment variant employed. In the system involving *Lolium perenne* and inoculation with a bacterial consortium, without the addition of nanomaterials, the degree of TPH biodegradation was 63.22%. The addition of nanomaterials significantly enhanced the efficiency of the biodegradation process. The variant enriched with Zn-O-NP exhibited a TPH degradation rate of 80.39%, whereas in the presence of GO-NPs, this value increased to 81.85%. This indicates that the incorporation of nanomaterials increased the efficiency of TPH biodegradation by approximately 17–19 percentage points compared with the system lacking additives. Figure 4 illustrates the mean degree of biodegradation of selected groups of hydrocarbons in soil sample G6 subjected to phytoremediation using *Lolium perenne*, including (a) aliphatic hydrocarbons and (b) PAHs, depending on the soil enrichment with nanomaterials.

A detailed analysis of the n-alkane profile indicated that the degree of biodegradation was strongly dependent on the length of the carbon chain and the phytoremediation variant employed (Figure 4a). In the *Lolium perenne* + bacterial consortium system (without the

https://doi.org/10.3390/molecules31050890

addition of nanomaterials), light n-alkane fractions (n-C6–n-C8) underwent biodegradation in the range of 82.26–91.42%, while medium-chain hydrocarbons (n-C9–n-C21) were biodegraded in the range of 68.65–97.35%. For heavier fractions, a marked decrease in biodegradation efficiency was observed. Aliphatic hydrocarbons with carbon chain length of n-C22–n-C30 underwent biodegradation only in the range of 43.16–62.49%, whereas the heaviest n-alkanes (n-C31–n-C36) were degraded in only 21.64–29.21%. The introduction of nanomaterials into soil substantially broadened the biodegradation ranges across all analysed variants. In systems containing GO-NPs and ZnO-NPs, light n-alkanes (n-C6–n-C8) underwent biodegradation in the range of 98.61–99.73%, while medium-chain fractions (n-C9–n-C21) were biodegraded at 84.52–100%. A particularly pronounced effect of nanomaterials was observed for heavier hydrocarbons, for which the biodegradation of n-C22–n-C30 increased to 61.59–78.76%, whereas that of n-C31–n-C36 increased to 34.91–44.79%. Comparison of the obtained ranges clearly indicates that the presence of nanomaterials not only enhanced the biodegradation of light and medium fractions but also significantly improved the decomposition of heavy hydrocarbons, which under natural conditions exhibit limited bioavailability and low susceptibility to biological degradation.




Figure 4. Average degree of biodegradation of selected groups: **(a)** aliphatic hydrocarbons and **(b)** polycyclic aromatic hydrocarbons in soil sample G6 subjected to phytoremediation using *Lolium perenne* under different soil enrichment variants. Values are means ± SD (n = 7–10). Different lowercase letters indicate significant differences among treatments according to one-way ANOVA followed by Tukey's HSD test ($p < 0.05$).

Analogous relationships were also observed for PAHs. In the system comprising *Lolium perenne* and a bacterial consortium, without the addition of nanomaterials, the total degree of PAHs biodegradation reached 61.86%. The incorporation of nanomaterials increased the efficiency of the process to 70.66–73.19%. Compared with the system without additives, the effectiveness of PAHs degradation was higher by approximately 9–11 percentage points, with slightly favourable results obtained for GO-NPs.

Analysis of the aromatic ring profile revealed pronounced differences in the susceptibility of individual PAH groups to biodegradation (Figure 4b). In the variant without nanomaterials, the highest degree of decomposition was observed for two-ring naphthalene,

https://doi.org/10.3390/molecules31050890

which underwent biodegradation at 74.14%. Three-ring PAHs were degraded in the range of 49.81–68.47%, while a clear decrease in biodegradation efficiency was observed with an increasing number of aromatic rings. Four-ring compounds were biodegraded in the range of 27.95–48.41%, five-ring compounds in 31.01–32.29%, and six-ring compounds only in 25.07–29.62%. Variants enriched with nanomaterials were characterised by a marked expansion of biodegradation ranges across all analysed PAH groups. In the presence of GO-NPs, the biodegradation of naphthalene increased to 82.44%, while that of three-ring PAHs reached 58.57–70.56%. For compounds exhibiting greater resistance to degradation, namely four-, five-, and six-ring PAHs, biodegradation efficiencies ranged from 49.97 to 52.91%, 33.73–36.38%, and 29.25–32.34%, respectively. A similar, though slightly less pronounced, effect was observed in the variant amended with ZnO-NP. The biodegradation of 2-ring naphthalene reached 80.56%, while three-ring PAHs were degraded in the range of 57.75–70.19%. In the case of four-, five- and six-ring PAHs, biodegradation efficiencies ranged from 39.49 to 52.29%, 34.89–54%, and 28.56–31.61%, respectively.

2.2.2. Analysis of Plant Material

Lolium perenne demonstrated a high degree of adaptation to soil conditions and a notable level of tolerance to petroleum-derived pollutants, thus confirming its suitability for phytoremediation processes. The germination and initial plant growth were observed in all experimental variants within the first week of the experiment (Figure 5a). In the system comprising *Lolium perenne* and a bacterial consortium, without the addition of nanomaterials, the length of seedlings after seven days was on average 5–7 cm. In contrast, in variants enriched with nanomaterials, including both GO-NPs and ZnO-NPs, seedlings reached a length of approximately 10 cm. Plants growing in the presence of GO-NPs and ZnO-NPs were characterised by greater shoot density and more intense green leaf coloration, indicating a beneficial effect of nanomaterials on the initial stages of growth and physiological condition of plants. After one month of the experiment, *Lolium perenne* in the variant inoculated only with the bacterial consortium reached a height of 25–30 cm, while in the variants enriched with nanomaterials, plant height ranged from 30 to 37 cm. These plants were well developed, rigid, and evenly distributed across the soil surface (Figure 5b). A particularly favourable development of the above-ground biomass was observed in the system amended with GO-NPs, which may indicate its stronger stimulatory effect on plant physiological processes. After a three-month phytoremediation period, the root system of perennial rye grass was well developed, exhibiting a typical fibrous structure with numerous thin lateral roots. Variants enriched with nanomaterials showed enhanced root development, with roots penetrating the entire soil volume, thereby improving soil structure, increasing permeability, and intensifying gas exchange. Such root system development could indirectly stimulate rhizosphere microbial activity and enhance the biodegradation of petroleum-derived hydrocarbons.

The analysis of dry biomass confirmed the visual observations regarding plant condition. In the variant comprising *Lolium perenne* and a bacterial consortium, the dry root biomass amounted to 1171.90 mg/kg DM, while the shoot biomass reached 2463.10 mg/kg DM. In the presence of GO-NPs, these values increased to 1533.30 mg/kg DM for roots and 3142.50 mg/kg DM for shoots. In turn, in the variant amended with ZnO-NP, dry root biomass reached 1519.90 mg/kg DM, while shoot biomass amounted to 2922.40 mg/kg DM. The results clearly indicate that soil enrichment with nanomaterials in combination with bacterial consortium inoculation promoted the growth of *Lolium perenne* and increased biomass accumulation in both above- and below-ground plant parts. The higher biomass values recorded in the variant amended with GO-NPs confirm its more favourable effect on plant development and its potential role in supporting phytoremediation processes. The

https://doi.org/10.3390/molecules31050890

distribution of dry biomass of *Lolium perenne* roots and shoots across individual experiment variants is presented in Figure 6.




Figure 5. Photographs of *Lolium perenne* after (**a**) 1 week and (**b**) 1 month of phytoremediation of soil sample G6 under different soil enrichment variants, where (1) soil G6 + *Lolium perenne* + bacterial consortium; (2) soil G6 + *Lolium perenne* + bacterial consortium + GO-NP (20 mg/kg DM); (3) soil G6 + *Lolium perenne* + bacterial consortium + ZnO-NP (20 mg/kg DM).

A comprehensive analysis of petroleum hydrocarbon content in *Lolium perenne* tissues revealed clear differences among individual phytoremediation variants, depending on the applied soil enrichment strategy. In the system comprising perennial ryegrass and a bacterial consortium, without the addition of nanomaterials, the concentration of TPH in the roots amounted to 110.83 mg/kg DM, while in shoots it reached 66.03 mg/kg DM. The application of GO-NPs in combination with bioaugmentation resulted in a significant increase in TPH accumulation in both plant organs, with concentration reaching 193.50 mg/kg DM in roots and 125.90 mg/kg DM in shoots. In the variant enriched with ZnO-NP, an

https://doi.org/10.3390/molecules31050890

elevated accumulation of petroleum hydrocarbons was also observed, with TPH contents of 187.70 mg/kg DM in roots and 120.30 mg/kg DM in shoots. The obtained results indicate that *Lolium perenne* exhibits a greater capacity for accumulating under conditions supported by nanomaterial addition. In all experimental variants, higher concentrations of petroleum-derived compounds were consistently detected in root tissues compared to above-ground parts (Figure 7). This pattern suggests a dominant role of the root system in the uptake and binding of hydrocarbons from the soil, as well as limited translocation of these compounds to shoots.




Figure 6. Biomass production of roots and shoots of *Lolium perenne* collected from soil subjected to phytoremediation under different soil enrichment variants. Values are means ± SD (n = 7–10). Different lowercase letters indicate significant differences among treatments according to one-way ANOVA followed by Tukey's HSD test ($p < 0.05$).

Analogous relationships were observed for PAHs in *Lolium perenne* tissues. In the variant containing only the bacterial consortium, PAHs concentration amounted to 77.20 µg/kg DM in roots, and 44.60 µg/kg DM in shoots. Soil enrichment with GO-NPs led to a pronounced increase in aromatic compound accumulation; with PAHs, concentration rose to 123.40 µg/kg DM in roots and 85.20 µg/kg DM in shoots. In the variant combining ZnO-NP with bioaugmentation, PAHs content reached 119.90 µg/kg DM in roots and 81.50 µg/kg DM in shoots. These results confirm that the presence of nanomaterials promoted enhanced PAHs accumulation in *Lolium perenne* tissues, with higher concentrations observed in the root system, similar to the case of TPH (Figure 7b). The highest accumulation levels of both TPH and PAHs were recorded in the variant enriched with GO-NP, suggesting its more effective role in modifying the bioavailability and stimulating pollutants uptake and binding by plants. The findings indicate that the combined application of *Lolium perenne*, a bacterial consortium, and nanomaterials, particularly GO-NPs, can substantially enhance phytoremediation efficiency through the simultaneous processes of biodegradation and phytoaccumulation of petroleum-derived and aromatic hydrocarbons.

https://doi.org/10.3390/molecules31050890




Figure 7. Concentration of (**a**) total petroleum hydrocarbons (TPH) and (**b**) polycyclic aromatic hydrocarbons (PAHs) in the roots and shoots of *Lolium perenne* collected from soil subjected to phytoremediation under different soil enrichment variants. Values are means ± SD (n = 7–10). Different lowercase letters indicate significant differences among treatments according to one-way ANOVA followed by Tukey's HSD test ($p < 0.05$).

2.3. Toxicological Analyses

Soil toxicity was assessed using the Microtox, Ostracodtoxkit, and Phytotoxkit tests, which included two biological parameters: seed germination and the inhibition of root growth in test plants (*Sinapis alba*, *Sorghum saccharatum*, and *Lepidium sativum*). The test organisms comprised *Vibrio fischeri* bacteria, *Heterocypris incongruens* crustaceans, and higher plants, respectively. To facilitate comparison of results and ensure unambiguous interpretation of toxicity levels, all outcomes were expressed as toxicity units (TUs).

The reference soil (Soil G6) was characterised by elevated biological toxicity, as evidenced by increased TU values across all applied tests. In the Microtox and Ostracodtoxkit tests, toxicity indices ranged from 10.50 to 13.05 TU, indicating a pronounced adverse effect of soil contamination on bacteria and crustaceans. The application of phytoremediation involving *Lolium perenne* in combination with a bacterial consortium resulted in a clear improvement in soil quality and a substantial reduction in biological toxicity. TU values decreased to 6.81 in the Microtox test and 5.99 in the Ostracodtoxkit test, corresponding to a toxicity reduction of approximately 45–50% relative to the initial soil. Further enhancement in soil quality was observed in variants enriched with nanomaterials. In the presence of GO-NPs, TU values declined to 2.82 in the Microtox test and 2.66 in the Ostracodtoxkit test, representingan additional decrease in toxicity of approximately 59% and 56%, respectively, compared to the variant planted with *Lolium perenne* and inoculated with the bacterial consortium. The variant enriched with ZnO-NP also exhibited a significant reduction in toxicity, with TU values of 3.06 (Microtox) and 2.81 (Ostracodtoxkit), corresponding to a reduction of approximately 53–55% compared to the planted and inoculated soil. These results confirm that the addition of nanomaterials effectively enhanced soil detoxification, with GO-NPs demonstrating slightly higher efficiency. Analogous relationships were observed in the Phytotoxkit test for both seed germination and root growth inhibition. The initial soil exhibited elevated phytotoxicity, with TU values ranging from 9.02 to 11.25 for

https://doi.org/10.3390/molecules31050890

seed germination and from 8.70 to 12.56 for root growth inhibition, depending on the plant species. The use of *Lolium perenne* together with the bacterial consortium reduced toxicity levels to 5.14 to 6.40 TU, confirming the beneficial effect of phytoremediation on plant growth conditions. Variants enriched with nanomaterials showed a further decrease in toxicity indices. In the presence of GO-NPs, TU values for both phytotoxicity parameters ranged from 2.50 to 2.92, corresponding to a reduction in toxicity of approximately 43–61% compared to the planted and inoculated variant. In the case of ZnO-NP, TU values ranged from 2.70 to 3.30, reflecting a toxicity reduction of approximately 36–58%. The obtained results clearly indicate that the application of phytoremediation supported by bacterial consortium inoculation significantly improved soil quality, while additional enrichment with nanomaterials, particularly GO-NPs, led to a further and pronounced reduction in biological toxicity. The consistent decrease in TU values across all applied tests confirms the effectiveness of the investigated remediation systems in mitigating the adverse effects of soil contamination on test organisms, thereby indicating an improvement in the environmental properties of the soil (Figure 8).




Figure 8. Toxicological analysis results for soil samples subjected to phytoremediation using *Lolium perenne* under different soil enrichment variants, with toxicity units (TUs) values determined by Microtox, Ostracodtoxkit, and Phytotoxkit tests. Values are means ± SD (n = 7–10). Different lowercase letters indicate significant differences among treatments according to one-way ANOVA followed by Tukey's HSD test ($p < 0.05$).

3. Discussion

Despite the prevalence and numerous environmental advantages of biological methods for the remediation of soil contaminated with petroleum hydrocarbons, further optimisation is required to increase their effectiveness and shorten remediation times. Numerous

https://doi.org/10.3390/molecules31050890

studies have shown that optimal outcomes are achieved through the integration of complementary strategies, including inoculation with microbial consortia, phytoremediation, or the application of additives that enhance pollutant bioavailability [20–23,59]. In recent years, increasing attention has been directed toward the use of nanomaterials to intensify biodegradation processes [60,61]. The present paper proposes an integrated approach combining bioaugmentation with the addition of GO-NPs or ZnO-NPs, followed by verification under both laboratory conditions and on a semi-technical scale with the participation of *Lolium perenne*. This approach enabled the assessment of the synergistic effect of these factors on soil remediation efficiency.

The results demonstrate that the application of an integrated strategy combining microbial biodegradation, phytoremediation, and the use of nanomaterials significantly improved the quality of soil contaminated with petroleum hydrocarbons. During the laboratory stage, a marked increase in microbial metabolic activity was observed in samples inoculated with a bacterial consortium, as confirmed by respirometric analyses. In variants containing GO-NPs and ZnO-NPs at a dose of 20 mg/kg DM, the highest oxygen consumption values were recorded, reaching 5147.8 mg O_2/dm^3 for GO-NPs and 4836.6 mg O_2/dm^3 for ZnO-NPs, clearly indicating the stimulation of microbiological processes [62]. For comparison, under laboratory conditions analogous to those applied in this study, the addition of surfactants to soil resulted in oxygen consumption levels of 5164.85 mg O_2/dm^3 for the PSR mixture, 5087.2 mg O_2/dm^3 for JBR425, and 4956.2 mg O_2/dm^3 for JBR320 [20]. These results indicate that the stimulatory effects of nanomaterials are comparable to those of conventional biodegradation-enhancing additives. The elevated oxygen consumption observed in this study corresponds to the active decomposition of organic compounds, including TPH and PAH fractions, and is consistent with reports indicating that respirometric parameters are reliable indicators of organic pollutant biodegradation [63,64]. These effects can be attributed to increased pollutant bioavailability, favourable conditions for microorganism colonisation, and the formation of surfaces conducive to biofilm development. The interaction between nanomaterials and soil microflora has been extensively documented in the literature [65,66]. At the same time, reductions in biological activity documented at higher nanomaterial doses confirm the dose-dependent nature of their effects and highlight the importance of moderate concentrations in eliciting stimulatory rather than toxic responses [67,68].

Chromatographic analyses conducted during laboratory experiments showed that the enhanced microbial activity resulted in a measurable decrease in petroleum hydrocarbon concentration in soil. The most effective degradation of the TPH fraction was observed in the variant enriched with GO-NPs at a dose of 20 mg/kg DM, where the reduction reached 70.47%. This effect is attributed to the adsorption of hydrophobic organic compounds onto the surface of GO-NPs, which increased their bioavailability to microorganisms [69,70]. A similar, though slightly lower, effect was obtained in variants amended with ZnO-NP. The highest TPH degradation in these systems (67.68%) was also recorded at a dose of 20 mg/kg DM, while in the other variants, TPH reduction ranged from 49.69% to 67.68%, confirming the dose-dependent influence of nanomaterials on biodegradation processes. The observed differences in susceptibility of individual aliphatic hydrocarbon fractions to biodegradation are consistent with reports indicating that light and medium-chain n-alkanes degrade more rapidly than heavy fractions [20–23,71,72]. Importantly, the presence of nanomaterials also promoted the degradation of more resistant compounds, including PAHs. Under laboratory conditions, PAH reduction ranged from 40.48 to 56.19% in variants containing GO-NPs and from 40.45 to 54.94% in those amended with ZnO-NP. These results are in good agreement with findings reported by other researchers. Xue et al. [73] demonstrated that GO-NPs increased hydrocarbon availability and accelerated degradation,

https://doi.org/10.3390/molecules31050890

achieving a 62.86% reduction in PAHs within 35 days. Similar trends were observed in the present study, confirming the effectiveness of GO-NPs in intensifying aromatic hydrocarbon biodegradation. PAHs removal efficiency was highest for compounds with fewer aromatic rings (2–4), whereas degradation of compounds with more than four rings was substantially lower, although still enhanced by the presence of GO-NPs and ZnO-NPs. Comparable relationships have been described in the literature, where graphene nanostructures increased PAHs desorption from soil matrices and facilitated their biological degradation [38,66]. ZnO-NP has likewise been reported to stimulate soil microbiota involved in pollutant removal [74].

Phytoremediation experiments confirmed that *Lolium perenne*, in combination with a bacterial consortium, effectively supported soil remediation, achieving 63.22% TPH biodegradation and 61.86% PAHs biodegradation. The extensive root system of this species promoted the formation of an active rhizosphere, enhancing microbial processes and pollutant bioavailability [75–77]. These results are consistent with literature data obtained under similar conditions with other plant species, such as *Echinacea purpurea* (35.56% TPHs; 30.82% PAHs after 180 days) [23], *Scirpus sylvaticus* (89.47% TPHs; 75.52% PAHs), and *Cirsium oleraceum* (93.96% TPHs; 79.55% PAHs) [21]. The efficacy of *Lolium perenne* in the removal of hydrocarbons was also substantiated in studies by Muratova et al., conducted on a former oil sludge deposit. In these studies, both cultivated plants (*Triticum aestivum*, *Secale cereale*, *Avena sativa*, *Hordeum vulgare*, *Sorghum bicolor*, *Panicum miliaceum*, and *Zea mays*) and wild grasses (including *Lolium perenne*, *Bromopsis inermis*, *Agropyron cristatum*, and *Festuca pratensis*) demonstrated the ability to reduce TPH, with *Secale cereale* exhibiting the highest reduction (82%) after 365 days, while *Lolium perenne* reduced TPH by 61% [78]. In a separate study, Tang et al. demonstrated that combining *Lolium perenne* with a mixed consortium of microorganisms (*Bacillus subtilis*, *Sphingobacterium multivolume*, *Acinetobacter radioresistens*, *Rhodococcus erythropolis*, and *Pseudomonas fluorescens*) enabled 58% TPH degradation after 162 days of phytoremediation, confirming the key role of synergy between plants and microorganisms in hydrocarbon removal processes [17]. Analogous trends were described by Hussain et al., where soil contaminated with TPH was treated using *Lolium perenne* in combination with additives that stimulated microbial activity. The highest hydrocarbon removal (40%) was achieved in variants containing biochar (5%), compost (5%), and immobilised microorganisms [79]. These results indicate that the efficacy of phytoremediation involving *Lolium perenne* can be enhanced by integrating supplementary strategies that facilitate microbiological processes, as realised in this study through the implementation of nanomaterials. The incorporation of nanomaterials further increased the efficacy of phytoremediation, as evidenced by TPH biodegradation rising to 81.85% in the presence of GO-NPs and to 80.9% with ZnO-NP, along with PAHs biodegradation increasing to 73.19% and 70.66%, respectively. These findings demonstrate a clear synergistic interaction between plants, microorganisms, and nanoparticles. In the case of GO-NPs, a particularly pronounced effect supporting phytoremediation processes was observed, likely due to their capacity to modify soil physicochemical properties and stimulate microbial activity in the rhizosphere [80]. Wang et al. and Li et al. reported that GO-NPs exhibit high porosity and adsorption, enhancing soil structure and creating favourable habitats for soil microorganisms [81,82]. Mu et al. observed that the addition of GO-NPs increased the removal rate of phenanthrene and anthracene from 1.7% to 52.7%, highlighting the role of GO-NPs in activating native bacterial communities [39]. Furthermore, Wang et al. demonstrated that GO-NPs applied in the phytoremediation of soil heavily contaminated with PAHs with *Fire Phoenix* plants led to an increase in the removal of six PAHs to 67.44% after 120 days, with the maximum efficiency at a GO-NP dose of 500 mg/kg. Under these conditions, benzo(k)fluoranthene degradation reached 86.40%, an increase of 11.77%

https://doi.org/10.3390/molecules31050890

compared to the variant without GO-NP [81]. The effectiveness of GO-NPs in accelerating PAHs degradation is also confirmed by Zhou et al., who analysed the impact of GO-NPs on the decomposition of 4- and 5-ring PAHs.

ZnO nanoparticles are more commonly reported in the context of heavy metals removal, such as arsenic [27], cadmium [83], and lead [84], but studies also indicate their effect on the degradation of organic compounds. Oluyinka-Edunro et al. showed that the application of ZnO-NP in soil contaminated with PAHs, lanted with *Panicum maximum* and *Aspilia africana*, resulted in the degradation of anthracene (56.5–60.9%), benzo[a]anthracene (71.3–74.6%), benzo[k]fluoranthene (39.7–45.8%), chrysene (54.2–58.8%), and naphthalene (83.5–85.2%) [85]. The authors also demonstrated that increasing ZnO-NP concentration enhanced PAH degradation. The efficacy of nanomaterials in remediating hydrocarbon-contaminated soils is supported by employing disparate types of nanostructures. Moses et al. reported that multi-walled carbon nanotubes (MWCNT) used in the phytoremediation of oil-contaminated soil with *Zea mays* enabled 90.72% TPH removal. Combining MWCNT with iron oxide nanoparticles (MWCNT + IONP) enhanced the efficiency to 91.37%, and applying a polymer nanocomposite (MWCNT + IONP + PEG) archieved 85.56% TPH reduction. These results underscore the potential of nanomaterials as effective additives supporting phytoremediation and nanoremediation of hydrocarbon-contaminated soils [86].

Analysis of *Lolium perenne* biomass showed that nanomaterials did not induce phytotoxicity; rather, they promoted plant growth. In this study, biomass increased by 28.63% in variants with GO-NPs and by 22.21% with ZnO-NP compared to the variant without nanomaterials. The higher dry weight of roots and shoots indicates that low concentrations of nanomaterials exert a stimulatory effect, consistent with the literature reports [49]. Analogous effects on plant growth have been observed in other species, such as rice and wheat, where the application of Fe_3O_4 and ZnO nanoparticles enhanced tolerance and biomass [87,88]. The relationship between nanomaterials and plant biomass was also observed in ryegrass studies, where combining nanoparticles with microorganisms increased plant biomass [89]. At the same time, the literature emphasizes that the effect of nanomaterials on plant growth strongly depends on their type and dosage. Negative phytotoxic effects of improperly selected nanomaterials or at high concentrations have been reported, for example, in studies using nZVI [90–92] and at high doses of GO [93], highlighting the need for precise selection of nanomaterial type and concentration in assisted phytoremediation processes.Increased biomass was accompanied by greater accumulation of TPH and PAHs in *Lolium perenne* tissues, confirming that phytoremediation mainly involved phytostabilization and rhizodegradation. The application of nanomaterials resulted in a significant increase in pollutant accumulation throughout the entire plant, including both roots and shoots. Total TPH accumulation in treatments with GO-NPs and ZnO-NP was 80.86% and 74.15% higher, respectively, than in the system without nanomaterials. A comparable trend was observed for PAHs, with total accumulation increasing by 71.26% in the presence of GO-NPs and by 65.35% with ZnO-NPs. Regardless of the variant, higher pollutant concentrations were observed in roots than in shoots, typical for grasses, indicating limited translocation. The absence of a negative growth effect confirms that pollutant levels did not exceed the phytotoxicity threshold.

Final toxicological assessment confirmed the efficacy of applied remediation strategies. Elevated toxicity units (TUs) were recorded in the initial soil: 8.70–12.56 (Phytotoxkit), 10.50 (Ostracodtoxkit) (13.05) (Microtox). indicating negative impacts on test organisms. Phytoremediation with *Lolium perenne* and a bacterial consortium significantly reduced soil toxicity, while additional nanomaterial amendments further decreased TU values in all tests. In GO-NP and ZnO-NP variants, TU values declined to 2.50–3.30 (Phytotoxkit), 2.66–2.81 (Ostracodtoxkit), and 2.82–3.06 (Microtox), with slightly lower toxicity in GO-

https://doi.org/10.3390/molecules31050890

NPs. These results indicate that the applied nanomaterial doses did not induce secondary soil toxicity or adverse effects on test organisms. Higher toxicity unit values (3.62–4.91) were reported in studies using biosurfactants and bacterial consortium [20], while slightly lower toxicity was observed in phytoremediation with *Scirpus sylvaticus* (TU = 2.1–2.4) and *Cirsium oleraceum* (TU = 1.3–1.7) [21]. Substantial toxicity reduction was also observed in studies with *Echinacea purpurea* and *Zea mays* (TU decreased to 2.7–4.4) [22,23]. The consistency of chromatographic, respirometric, and toxicological results, combined with favourable remediation outcomes, provides strong evidence that an integrated approach combining *Lolium perenne*, bacterial consortium inoculation, and nanomaterial amendment (GO-NP and ZnO-NP) is highly effective in reducing both contaminant concentrations and associated toxicity in hydrocarbon-contaminated soils.

4. Materials and Methods

4.1. Materials

The bioremediation studies examined soil that had been heavily contaminated with petroleum compounds. These samples had been characterised and analysed in other studies. The total petroleum hydrocarbon (TPH) content in the tested material was 6062.59 mg/kg DM, while the concentration of polycyclic aromatic hydrocarbons (PAHs) was 12.05 mg/kg DM [20]. A more detailed physicochemical characterisation of the raw soil G6, including moisture, pH, organic pollutants, and selected metal and ion contents, is presented in Figure 9.




Figure 9. Comprehensive physicochemical characterisation of raw soil G6.

The raw soil G6 was subsequently prepared for bioremediation to provide optimal conditions for soil microbial growth through supplementation with lime fertiliser and Azofoska, aimed at correcting the pH and balancing the N:C:P ratio. Following this amendment, the soil exhibited a pH of 7.2, moisture content of 20%, NH_4^+ concentration of 65.1 mg/kg DM, NO_3^- of 238.9 mg/kg DM, and PO_4^{3-} of 28 mg/kg DM. This amended material constituted the final soil G6 used in the bioremediation experiments.

A multi-strain bacterial consortium was used for soil inoculation, a potential for whose degradation of petroleum-derived hydrocarbons has been confirmed in our previous studies. The following strains comprised the inoculum: *Dietzia* sp. IN118, *Gordonia* sp. IN101, *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN119, *Rhodococcus globerulus* IN113, and *Raoultella* sp. IN109. A detailed description of the procedure for

https://doi.org/10.3390/molecules31050890

isolating microorganisms, selecting strains, and constructing the bacterial consortium has been presented in our previous publications [20–23].

The study employed nanomaterials in the form of graphene nanoparticles (GO-NPs) MKCK8154 (Sigma-Aldrich, Saint Louis, MO, USA) and zinc nanoparticles (ZnO-NPs) 30-1405 (STREM Chemicals Inc., Newburyport, MA, USA).

Lolium perenne seeds from the seed company "GRANUM" Sp.j. (Łódź, Poland) were used in phytoremediation studies and were employed to sow the tested soil material.

4.2. Methods

4.2.1. Biodegradation Studies–Laboratory Scale

The biodegradation of petroleum hydrocarbons was investigated under laboratory conditions. Experimental samples consisted of 30 g of soil, to which GO-NP or ZnO-NP were added at concentrations of 10, 20, 50, and 100 mg/kg DM, along with 2 cm^3 of a bacterial consortium. Control samples included only the bacterial consortium (without nanomaterials) or a blank sample. Experiments were conducted for 90 days under controlled conditions: 20% moisture, 20 °C, with continuous air access.

Respirometric Analysis

The potential for aerobic biodegradation was evaluated using the OxiTop® Control system (WTW, Weilheim, Germany). Samples were placed in sealed respirometric bottles, and the system recorded pressure every two hours. Data were transferred to the OxiTop OC 110 controller, and graphical and statistical analyses were performed using the Achat OC software. Oxygen consumption (mg O_2/dm^3) was calculated based on the pressure decrease using the following formula:

$$m(O_2) = \frac{\Delta p \cdot V_g \cdot M(O_2) \cdot \alpha \cdot T_0}{R \cdot T_m}$$

where $m(O_2)$—mass of consumed oxygen [mg/dm^3], Δp—pressure drop [Pa], Vg—gas volume in the bottle [m^3], $M(O_2)$—molar mass of oxygen [kg/mol], R—gas constant [$J \cdot mol^{-1} \cdot K^{-1}$], T_m—measurement temperature [K], T_0 = 273.15 K—reference temperature, α = 0.03103—absorption coefficient. Based on these data, the Synergistic Index (SI) was calculated to evaluate the interaction between nanomaterials and the bacterial consortium.

TPH and PAHs Analysis

TPH was extracted using ultrasound-assisted dichloromethane extraction (POCH S.A., Gliwice, Poland) in three cycles of 20 mL each, 15 min per cycle, in a Sonoswiss SW 6H ultrasonic bath (30 kHz) (Ramsen, Switzerland). Extracts were purified with Florisil SPE columns (Bakerbond No. JTB-7213-03) (J.T. Baker®, Phillipsburg, NJ, USA), concentrated under reduced pressure, and re-dissolved in 1 mL dichloromethane. Quantification of TPH, n-alkanes (n-C6–n-C44), and isoprenoid hydrocarbons (pristane, phytane) was performed on a Clarus 500 gas chromatograph (PerkinElmer, Waltham, MA, USA) with FID detection on an RTX-1 column (30 m × 0.53 mm, Restek, Bellefonte, PA, USA). Oven programme: 30 °C for 2 min, ramp to 105 °C at 10 °C/min, then to 285 °C at 5 °C/min, 285 °C isothermal for 5 min. Injector and detector temperatures: 290 °C and 320 °C. Standards: BAM K010 (Tusnovic Instruments, Krakow, Poland) for TPH, Supelco D2807 (n-C6–n-C44) and Fuel Oil Degradation Mix A029668 (Restek, USA) for n-alkanes, Hopane No. 08189 (Sigma-Aldrich, Saint Louis, MO, USA).

PAHs were extracted using a QuEChERS method. Samples were spiked with water and acetonitrile (Chempur, Piekary Śląskie, Poland), and $MgSO_4$/NaCl salts (SST640, Interchim, Montluçon, France) were added. After centrifugation (3500 rpm, 10 min, MPW-223e, MPW,

https://doi.org/10.3390/molecules31050890

Warszawa, Poland), extracts were purified using dispersive SPE (JO3937, Interchim, France) and filtered through 0.2 µm nylon filters. PAHs were analysed on a Vanquish Core HPLC system (Thermo Scientific, Waltham, MA, USA) with UV–VIS and fluorescence detection on a NUCLEODUR C18 PAH column (125 × 4 mm, 3 µm, Macherey-Nagel, Düren, Germany). Mobile phases: 70% methanol (A) and 99% acetonitrile (B), flow rate 1.5 mL/min, gradient: 20–50% B (1.5 min), 50–100% B (1 min), 100% B (1 min), decrease to 0% B (3 min), final isothermal 100% A (3 min). Standards: PAH Standard Ref. No. 722393 (Macherey-Nagel, Düren, Germany) and certified reference material PAHs by HPLC 40g SQC017-40G (Sigma Aldrich, Saint Louis, MO, USA) [20–23]. Biodegradation of individual and total PAHs was calculated.

The schematic workflow for TPH and PAHs extraction and analysis is shown in Figure 10.




Figure 10. Schematic representation of the extraction and chromatographic analysis of TPH and PAHs.

4.2.2. Phytoremediation Studies–Semi-Technical Scale

A phytoremediation study was conducted under semi-technical conditions using 1 kg samples of soil exposed to natural atmospheric conditions. The pot method was applied: soil was placed in pots and sown with *Lolium perenne* seeds. Three experimental systems were tested: soil inoculated with the bacterial consortium, soil enriched with GO-NP (20 mg/kg DM) and inoculated with the bacterial consortium, and soil enriched with ZnO-NP (20 mg/kg DM) and inoculated with the bacterial consortium. In all cases, the pots were sown with *Lolium perenne*.

The phytoremediation period lasted 90 days, during which standard plant care was maintained. At the end of the experiment, samples were analysed for TPH, n-alkanes,

https://doi.org/10.3390/molecules31050890

and PAHs using the procedures described in the previous section. Additionally, plant biomass and concentrations of TPH and PAHs in roots and shoots of *Lolium perenne* were determined. The soils were also evaluated for potential toxicity using a combination of bioassays. Chronic toxicity was assessed with the Phytotoxkit™ assay (MicroBioTests Inc., Gent, Belgium), which measures inhibition of seed germination and early root elongation in three plant species with different germination and growth rates: *Sorghum saccharatum*, *Lepidium sativum* (cress), and *Sinapis alba* (white mustard) over a 72 h period. Toxicity to juvenile crustaceans was determined with the Ostracodtoxkit(F)™ assay (MicroBioTests Inc., Belgium), where *Heterocypris incongruens* hatched from cysts were exposed to soil samples in six-well plates, each containing 0.5 g of soil and 2 mL of standardised algal suspension as food, with ten individuals per well; reference soils served as controls during six days of incubation. The Microtox® Solid Phase Test (SDI, Newark, DE, USA) used the bioluminescent bacterium *Vibrio fischeri* to measure inhibition of luminescence catalysed by bacterial luciferase. Lyophilized bacteria were stored at −20 °C and rehydrated prior to testing, and serial dilutions of soil extracts were applied. The EC_{50} value, representing the concentration causing 50% inhibition of luminescence, was determined, with lower EC_{50} values indicating higher toxicity. Detailed procedures for the extraction and chromatographic determination of petroleum hydrocarbons in both soil and plant material, as well as the execution of the toxicological assays, have been described in our previous publications [20–23].

4.2.3. Statistical Analysis

All experimental data were processed using Statistica 14.0 (TIBCO Software Inc., Palo Alto, CA, USA). Prior to analysis, the normality of the datasets was assessed. One-way analysis of variance (ANOVA) was then performed, and where significant differences were detected, Tukey's post hoc test was applied for multiple comparisons. Additionally, standard deviation (SD) and Pearson correlation coefficients were calculated. Differences were considered statistically significant at $p < 0.05$ [20–23].

Author Contributions: Conceptualization, K.W., T.S., P.K. and J.B.; Methodology, K.W., T.S., P.K. and J.B.; Validation, K.W., T.S. and P.K.; Formal Analysis, K.W., T.S., P.K. and J.B.; Investigation, K.W.; Writing—Original Draft Preparation, K.W.; Writing—Review and Editing, K.W., T.S. and P.K. All authors have read and agreed to the published version of the manuscript.

Funding: This research was financially supported by the Polish Ministry of Science and Higher Education within statutory funding for the Oil and Gas Institute-National Research Institute. DK:4100-15-25.

Institutional Review Board Statement: Not applicable. "The studies used commercial toxicological tests not subject to ethical evaluation".

Informed Consent Statement: Not applicable.

Data Availability Statement: Dataset available on request from the authors.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Bhatia, R.K.; Sakhuja, D.; Mundhe, S.; Walia, A. Renewable Energy Products through Bioremediation of Wastewater. *Sustainability* **2020**, *12*, 7501. [CrossRef]
2. Kalia, A.; Sharma, S.; Semor, N.; Babele, P.K.; Sagar, S.; Bhatia, R.K.; Walia, A. Recent Advancements in Hydrocarbon Bioremediation and Future Challenges: A Review. *3 Biotech* **2022**, *12*, 135. [CrossRef] [PubMed]
3. Radhakrishnan, A.; Balaganesh, P.; Vasudevan, M.; Natarajan, N.; Chauhan, A.; Arora, J.; Ranjan, A.; Rajput, V.D.; Sushkova, S.; Minkina, T.; et al. Bioremediation of Hydrocarbon Pollutants: Recent Promising Sustainable Approaches, Scope, and Challenges. *Sustainability* **2023**, *15*, 5847. [CrossRef]

https://doi.org/10.3390/molecules31050890

4. Hamad, A.A.; Moubasher, H.A.; Moustafa, Y.M.; Mohamed, N.H.; Abd-El Rhim, E.H. Petroleum Hydrocarbon Bioremediation Using Native Fungal Isolates and Consortia. *Sci. World J.* **2021**, *2021*, 6641533. [CrossRef]
5. Abtahi, H.; Parhamfar, M.; Saeedi, R.; Villaseñor, J.; Sartaj, M.; Kumar, V.; Coulon, F.; Parhamfar, M.; Didehdar, M.; Seifi, H.; et al. Effect of Competition between Petroleum-Degrading Bacteria and Indigenous Compost Microorganisms on the Efficiency of Petroleum Sludge Bioremediation: Field Application of Mineral-Based Culture in the Composting Process. *J. Environ. Manag.* **2020**, *258*, 110013. [CrossRef]
6. Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J.; Skalski, T. Evaluation of the Effectiveness of the Biopreparation in Combination with the Polymer γ-PGA for the Biodegradation of Petroleum Contaminants in Soil. *Materials* **2022**, *15*, 400. [CrossRef] [PubMed]
7. Ortega-Calvo, J.J.; Lahlou, M.; Saiz-Jimenez, C. Effect of Organic Matter and Clays on the Biodegradation of Phenanthrene in Soils. *Int. Biodeterior. Biodegrad.* **1997**, *40*, 101–106. [CrossRef]
8. Weissenfels, W.D.; Klewer, H.-J.; Langhoff, J. Adsorption of Polycyclic Aromatic Hydrocarbons (PAHs) by Soil Particles: Influence on Biodegradability and Biotoxicity. *Appl. Microbiol. Biotechnol.* **1992**, *36*, 689–696. [CrossRef]
9. Kaimi, E.; Mukaidani, T.; Tamaki, M. Screening of Twelve Plant Species for Phytoremediation of Petroleum Hydrocarbon-Contaminated Soil. *Plant Prod. Sci.* **2007**, *10*, 211–218. [CrossRef]
10. Kamath, R.; Rentz, J.A.; Schnoor, J.L.; Alvarez, P.J.J. Chapter 16 Phytoremediation of Hydrocarbon-Contaminated Soils: Principles and Applications. *Stud. Surf. Sci. Catal.* **2004**, *151*, 447–478.
11. Hussain, I.; Puschenreiter, M.; Gerhard, S.; Schöftner, P.; Yousaf, S.; Wang, A.; Syed, J.H.; Reichenauer, T.G. Rhizoremediation of Petroleum Hydrocarbon-Contaminated Soils: Improvement Opportunities and Field Applications. *Environ. Exp. Bot.* **2018**, *147*, 202–219. [CrossRef]
12. Conlon, R.; Dowling, D.N.; Germaine, K.J. Assessing Microbial Activity and Rhizoremediation in Hydrocarbon and Heavy Metal-Impacted Soil. *Microorganisms* **2025**, *13*, 848. [CrossRef]
13. Cunningham, S.D.; Berti, W.R. Remediation of Contaminated Soils with Green Plants: An Overview. *Vitr. Cell. Dev. Biol. Plant* **1993**, *29*, 207–212. [CrossRef]
14. Gaskin, S.E.; Bentham, R.H. Rhizoremediation of Hydrocarbon Contaminated Soil Using Australian Native Grasses. *Sci. Total Environ.* **2010**, *408*, 3683–3688. [CrossRef]
15. Cook, R.L.; Hesterberg, D. Comparison of Trees and Grasses for Rhizoremediation of Petroleum Hydrocarbons. *Int. J. Phytoremediat.* **2013**, *15*, 844–860. [CrossRef] [PubMed]
16. Steliga, T.; Kluk, D. Application of Festuca Arundinacea in Phytoremediation of Soils Contaminated with Pb, Ni, Cd and Petroleum Hydrocarbons. *Ecotoxicol. Environ. Saf.* **2020**, *194*, 110409. [CrossRef]
17. Tang, J.; Wang, R.; Niu, X.; Zhou, Q. Enhancement of Soil Petroleum Remediation by Using a Combination of Ryegrass (*Lolium perenne*) and Different Microorganisms. *Soil Tillage Res.* **2010**, *110*, 87–93. [CrossRef]
18. Wyszkowska, J.; Borowik, A.; Kucharski, J. The Resistance of *Lolium perenne* L. × Hybridum, *Poa pratensis*, *Festuca rubra*, F. Arundinacea, *Phleum pratense* and *Dactylis glomerata* to Soil Pollution by Diesel Oil and Petroleum. *Plant Soil Environ.* **2019**, *65*, 307–312. [CrossRef]
19. He, M.; Li, Z.; Chen, C.; Mei, P. Impact of Soil Types and Root Exudates on Cadmium and Petroleum Hydrocarbon Phytoremediation by *Sorghum sudanense*, *Festuca arundinace*, and *Lolium perenne*. *Front. Ecol. Evol.* **2022**, *10*, 1036765. [CrossRef]
20. Wojtowicz, K.; Steliga, T.; Skalski, T.; Kapusta, P. Influence of Biosurfactants on the Efficiency of Petroleum Hydrocarbons Biodegradation in Soil. *Sustainability* **2025**, *17*, 6520. [CrossRef]
21. Wojtowicz, K.; Steliga, T.; Brzeszcz, J.; Fyda, J.; Skalski, T.; Kapusta, P. Phytoremediation of Soil Contaminated with Petroleum Hydrocarbons Using the Wild Plants Scirpus Sylvaticus and Cirsium Oleraceum, Supported by Bioaugmentation. *Int. Biodeterior. Biodegrad.* **2026**, *206*, 106217. [CrossRef]
22. Wojtowicz, K.; Steliga, T.; Kapusta, P.; Brzeszcz, J. Oil-Contaminated Soil Remediation with Biodegradation by Autochthonous Microorganisms and Phytoremediation by Maize (*Zea mays*). *Molecules* **2023**, *28*, 6104. [CrossRef]
23. Wojtowicz, K.; Steliga, T.; Kapusta, P. Evaluation of the Effectiveness of Bioaugmentation-Assisted Phytoremediation of Soils Contaminated with Petroleum Hydrocarbons Using *Echinacea purpurea*. *Appl. Sci.* **2023**, *13*, 13077. [CrossRef]
24. Wojtowicz, K.; Steliga, T. Ocena Skuteczności Biodegradacji Wspomaganej Inokulacją Biopreparatem z Dodatkiem Surfaktantu w Procesie Oczyszczania Gleby Historycznie Zanieczyszczonej Węglowodorami Ropopochodnymi. *Nafta-Gaz* **2025**, *81*, 381–388. [CrossRef]
25. Mohanapragash, A.G.; Kaleeswari, R.K.; Madhupriyaa, D.; Salma Santhosh, S.; Meena, S.; Baskar, M.; Umamaheswari, T.; Selvamurugan, M.; Ramesh, T. Nanotechnology in Sustainable Soil Remediation: Exploring a Review on Carbon-Based, Clay-Based, Dendrimer, Polymer, and Nanocomposite Based Nanomaterials for Environmental Cleanup. *Int. J. Environ. Anal. Chem.* **2025**, *105*, 8533–8558. [CrossRef]
26. Souza, L.R.R.; Pomarolli, L.C.; Da Veiga, M.A.M.S. From Classic Methodologies to Application of Nanomaterials for Soil Remediation: An Integrated View of Methods for Decontamination of Toxic Metal (Oid)s. *Environ. Sci. Pollut. Res.* **2020**, *27*, 10205–10227. [CrossRef] [PubMed]

https://doi.org/10.3390/molecules31050890

27. Pérez-Hernández, H.; Pérez-Moreno, A.Y.; Méndez-López, A.; Fernández-Luqueño, F. Effect of ZnO Nanoparticles During the Process of Phytoremediation of Soil Contaminated with As and Pb Cultivated with Sunflower (*Helianthus annuus* L.). *Int. J. Environ. Res.* **2024**, *18*, 7. [CrossRef]
28. Németh, I.; Vaszita, E.; Molnár, M. Influence of nZnO on Enzyme-Mediated PAH-Removal from Contaminated Soil. *Period. Polytech. Chem. Eng.* **2025**, *69*, 361–375. [CrossRef]
29. Baragaño, D.; Alonso, J.; Gallego, J.R.; Lobo, M.C.; Gil-Díaz, M. Magnetite Nanoparticles for the Remediation of Soils Co-Contaminated with As and PAHs. *Chem. Eng. J.* **2020**, *399*, 125809. [CrossRef]
30. Mirzaee, E.; Gitipour, S.; Mousavi, M.; Amini, S. Optimization of Total Petroleum Hydrocarbons Removal from Mahshahr Contaminated Soil Using Magnetite Nanoparticle Catalyzed Fenton-like Oxidation. *Environ. Earth Sci* **2017**, *76*, 165. [CrossRef]
31. Yao, S.; Ouyang, S.; Zhou, Q.; Tao, Z.; Chen, Y.; Zheng, T. Environmental Remediation and Sustainable Design of Iron Oxide Nanoparticles for Removal of Petroleum-Derived Pollutants from Water: A Critical Review. *Environ. Res.* **2024**, *263*, 120009. [CrossRef]
32. Naderi Peikam, E.; Jalali, M. Application of Three Nanoparticles (Al_2O_3, SiO_2 and TiO_2) for Metal-Contaminated Soil Remediation (Measuring and Modeling). *Int. J. Environ. Sci. Technol.* **2019**, *16*, 7207–7220. [CrossRef]
33. Zhang, Y.; Wang, J.; Liu, X.; Zhang, L.; Wang, J.; Xu, C.; Qian, T.; Guo, J. Nano-TiO_2 with Humic Acid to Promote Remediation of Cadmium/Pyrene Composite Contaminated Soil by *Hylotelephium spectabile*. *J. Environ. Chem. Eng.* **2025**, *13*, 116044. [CrossRef]
34. Zand, A.D.; Mikaeili Tabrizi, A.; Vaezi Heir, A. Application of Titanium Dioxide Nanoparticles to Promote Phytoremediation of Cd-Polluted Soil: Contribution of PGPR Inoculation. *Bioremediat. J.* **2020**, *24*, 171–189. [CrossRef]
35. Cheng, J.; Sun, Z.; Yu, Y.; Li, X.; Li, T. Effects of Modified Carbon Black Nanoparticles on Plant-Microbe Remediation of Petroleum and Heavy Metal Co-Contaminated Soils. *Int. J. Phytoremediat.* **2019**, *21*, 634–642. [CrossRef]
36. Zhao, L.; Yang, S.-T.; Yilihamu, A.; Wu, D. Advances in the Applications of Graphene Adsorbents: From Water Treatment to Soil Remediation. *Rev. Inorg. Chem.* **2019**, *39*, 47–76. [CrossRef]
37. Özlü, V.; Elif Fırat, M. Fly Ash and Nano-Graphene Enhanced Stabilization of Engine Oil-Contaminated Soils. *Nanotechnol. Rev.* **2025**, *14*, 20250210. [CrossRef]
38. Zhou, X.; Zhang, B.; Meng, Q.; Li, L. Effects of Graphene Oxide on Endophytic Bacteria Population Characteristics in Plants from Soils Contaminated by Polycyclic Aromatic Hydrocarbons. *Molecules* **2024**, *29*, 2342. [CrossRef]
39. Mu, L.; Zhou, Q.; Zhao, Y.; Liu, X.; Hu, X. Graphene Oxide Quantum Dots Stimulate Indigenous Bacteria to Remove Oil Contamination. *J. Hazard. Mater.* **2019**, *366*, 694–702. [CrossRef] [PubMed]
40. Moses, H.; Odigure, J.O.; Otaru, A.J.; Oduoza, C.F. A Potential Remediation of Total Petroleum Hydrocarbon from Contaminated Soils Using Surfactant Suspended Multiwall Carbon Nanotubes. *Results Eng.* **2023**, *19*, 101371. [CrossRef]
41. Matos, M.P.S.R.; Correia, A.A.S.; Rasteiro, M.G. Application of Carbon Nanotubes to Immobilize Heavy Metals in Contaminated Soils. *J. Nanopart. Res.* **2017**, *19*, 126. [CrossRef]
42. Nwadinigwe, A.O.; Ugwu, E.C. Overview of Nano-Phytoremediation Applications. In *Phytoremediation*; Ansari, A.A., Gill, S.S., Gill, R., Lanza, G.R., Newman, L., Eds.; Springer International Publishing: Cham, Switzerland, 2018; pp. 377–382.
43. Wang, M.; Zheng, X.; Oba, B.T.; Lin, Y.; Shen, C.; Huang, X.; Yang, F.; Xiao, Q.; Ding, Y. Innovations in Nanomaterials for Remediation of Heavy Metal−polluted Soil: Advances, Mechanistic Insights, and Future Prospects. *Nano Mater. Sci.* **2025**, *8*, 11–35. [CrossRef]
44. Jiang, C.; Liu, S.; Phenrat, T.; Sui, Q. Nanomaterial-Enabled Environmental Remediation and Removal of Emerging Pollutants. *Toxics* **2025**, *13*, 810. [CrossRef]
45. Ding, N.; Han, M.; He, Y.; Wang, X.; Pan, Y.; Lin, H.; Yu, G. Advances in Application of Nanomaterials in Remediation of Heavy Metal Contaminated Soil. *E3S Web Conf.* **2021**, *261*, 04027. [CrossRef]
46. Rai, P.K.; Kim, K.-H.; Lee, S.S.; Lee, J.-H. Molecular Mechanisms in Phytoremediation of Environmental Contaminants and Prospects of Engineered Transgenic Plants/Microbes. *Sci. Total Environ.* **2020**, *705*, 135858. [CrossRef]
47. Song, B.; Xu, P.; Chen, M.; Tang, W.; Zeng, G.; Gong, J.; Zhang, P.; Ye, S. Using Nanomaterials to Facilitate the Phytoremediation of Contaminated Soil. *Crit. Rev. Environ. Sci. Technol.* **2019**, *49*, 791–824. [CrossRef]
48. Zeb, A.; Liu, W.; Zhang, Y. Effects of Nanoparticles on Alleviating Phytotoxicity of Soil Heavy Metals: Potential for Enhancing Phytoremediation. In *Sustainable Agriculture Reviews* 53; Faizan, M., Hayat, S., Yu, F., Eds.; Springer International Publishing: Cham, Switzerland, 2021; Volume 53, pp. 327–344.
49. Gomes, M.P. Nanophytoremediation: Advancing Phytoremediation Efficiency through Nanotechnology Integration. *Discov. Plants* **2025**, *2*, 8. [CrossRef]
50. Chen, X.; Wang, S.; Lai, H.; Deng, L.; Zhong, Q.; Okoye, C.O.; Niu, Q.; Jing, Y.; Wang, J.; Jiang, J. Nano-Biotechnology in Soil Remediation: Use of Nanomaterials to Promote Plant Growth and Stress Tolerance. *Nanomaterials* **2025**, *15*, 1743. [CrossRef]
51. Obi, C.C.; Abonyi, M.N.; Ohale, P.E.; Onu, C.E.; Nwabanne, J.T.; Igwegbe, C.A.; Kamuche, T.T.; Ozofor, I.H. Adsorption of Antibiotics from Aqueous Media Using Nanocomposites: Insight into the Current Status and Future Perspectives. *Chem. Eng. J.* **2024**, *497*, 154767. [CrossRef]

https://doi.org/10.3390/molecules31050890

52. Ma, C.; Han, L.; Shang, H.; Hao, Y.; Xu, X.; White, J.C.; Wang, Z.; Xing, B. Nanomaterials in Agricultural Soils: Ecotoxicity and Application. *Curr. Opin. Environ. Sci. Health* **2023**, *31*, 100432. [CrossRef]
53. Alazaiza, M.Y.D.; Alzghoul, T.M.; Ramu, M.B.; Nassani, D.E. Catalytic Nanomaterials for Soil and Groundwater Remediation: Global Research Trends (2010–2024). *Catalysts* **2025**, *15*, 981. [CrossRef]
54. Ulhassan, Z.; Khan, I.; Hussain, M.; Khan, A.R.; Hamid, Y.; Hussain, S.; Allakhverdiev, S.I.; Zhou, W. Efficacy of Metallic Nanoparticles in Attenuating the Accumulation and Toxicity of Chromium in Plants: Current Knowledge and Future Perspectives. *Environ. Pollut.* **2022**, *315*, 120390. [CrossRef]
55. Prakash, P.; Smitha Chandran, S. Nano-Phytoremediation of Heavy Metals from Soil: A Critical Review. *Pollutants* **2023**, *3*, 360–380. [CrossRef]
56. Sattar, S.; Siddiqui, S.; Shahzad, A.; Bano, A.; Naeem, M.; Hussain, R.; Khan, N.; Jan, B.L.; Yasmin, H. Comparative Analysis of Microbial Consortiums and Nanoparticles for Rehabilitating Petroleum Waste Contaminated Soils. *Molecules* **2022**, *27*, 1945. [CrossRef]
57. Ren, W.; Liu, H.; Mao, T.; Teng, Y.; Zhao, R.; Luo, Y. Enhanced Remediation of PAHs-Contaminated Site Soil by Bioaugmentation with Graphene Oxide Immobilized Bacterial Pellets. *J. Hazard. Mater.* **2022**, *433*, 128793. [CrossRef]
58. Mathur, S.; Singh, D.; Ranjan, R. Remediation of Heavy Metal (Loid) Contaminated Soil through Green Nanotechnology. *Front. Sustain. Food Syst.* **2022**, *6*, 932424. [CrossRef]
59. Sui, X.; Wang, X.; Li, Y.; Ji, H. Remediation of Petroleum-Contaminated Soils with Microbial and Microbial Combined Methods: Advances, Mechanisms, and Challenges. *Sustainability* **2021**, *13*, 9267. [CrossRef]
60. Li, M.; Luo, L. Review on Application of Nanomaterials in Soil Remediation. *J. Phys. Conf. Ser.* **2020**, *1637*, 012070. [CrossRef]
61. Hussain, A.; Kabir, M. Synergistic Nanobioremediation: Exploring Nanoparticle–Plant–Microbe Interactions for Sustainable Soil Decontamination. *Nanotechnol. Environ. Eng.* **2026**, *11*, 11. [CrossRef]
62. Simonin, M.; Richaume, A. Impact of Engineered Nanoparticles on the Activity, Abundance, and Diversity of Soil Microbial Communities: A Review. *Environ. Sci. Pollut. Res.* **2015**, *22*, 13710–13723. [CrossRef]
63. Plaza, G.; Ulfig, K.; Worsztynowicz, A.; Malina, G.; Krzeminska, B.; Brigmon, R.L. Respirometry for Assessing the Biodegradation of Petroleum Hydrocarbons. *Environ. Technol.* **2005**, *26*, 161–170. [CrossRef] [PubMed]
64. Aichberger, H.; Hasinger, M.; Braun, R.; Loibner, A.P. Potential of Preliminary Test Methods to Predict Biodegradation Performance of Petroleum Hydrocarbons in Soil. *Biodegradation* **2005**, *16*, 115–125. [CrossRef] [PubMed]
65. Ikuma, K.; Decho, A.W.; Lau, B.L.T. When Nanoparticles Meet Biofilms—Interactions Guiding the Environmental Fate and Accumulation of Nanoparticles. *Front. Microbiol.* **2015**, *6*, 591. [CrossRef] [PubMed]
66. Bhatia, R.; Gulati, D.; Sethi, G. Biofilms and Nanoparticles: Applications in Agriculture. *Folia Microbiol.* **2021**, *66*, 159–170. [CrossRef]
67. Qian, Y.; Qin, C.; Chen, M.; Lin, S. Nanotechnology in Soil Remediation—Applications vs. Implications. *Ecotoxicol. Environ. Saf.* **2020**, *201*, 110815. [CrossRef]
68. Dinesh, R.; Anandaraj, M.; Srinivasan, V.; Hamza, S. Engineered Nanoparticles in the Soil and Their Potential Implications to Microbial Activity. *Geoderma* **2012**, *173–174*, 19–27. [CrossRef]
69. Avornyo, A.; Chrysikopoulos, C.V. Applications of Graphene Oxide (GO) in Oily Wastewater Treatment: Recent Developments, Challenges, and Opportunities. *J. Environ. Manag.* **2024**, *353*, 120178. [CrossRef]
70. Eigler, S.; Hirsch, A. Chemistry with Graphene and Graphene Oxide—Challenges for Synthetic Chemists. *Angew. Chem. Int. Ed.* **2014**, *53*, 7720–7738. [CrossRef]
71. Wu, M.; Li, W.; Dick, W.A.; Ye, X.; Chen, K.; Kost, D.; Chen, L. Bioremediation of Hydrocarbon Degradation in a Petroleum-Contaminated Soil and Microbial Population and Activity Determination. *Chemosphere* **2017**, *169*, 124–130. [CrossRef]
72. Nwankwegu, A.S.; Orji, M.U.; Onwosi, C.O. Studies on Organic and In-Organic Biostimulants in Bioremediation of Diesel-Contaminated Arable Soil. *Chemosphere* **2016**, *162*, 148–156. [CrossRef]
73. Xue, C.; Li, L.; Guo, C.; Gao, Y.; Yang, C.; Deng, X.; Li, X.; Tai, P.; Sun, L. Understanding the Role of Graphene Oxide in Affecting PAHs Biodegradation by Microorganisms: An Integrated Analysis Using 16SrRNA, Metatranscriptomic, and Metabolomic Approaches. *J. Hazard. Mater.* **2023**, *457*, 131811. [CrossRef]
74. Strekalovskaya, E.I.; Perfileva, A.I.; Krutovsky, K.V. Zinc Oxide Nanoparticles in the "Soil–Bacterial Community–Plant" System: Impact on the Stability of Soil Ecosystems. *Agronomy* **2024**, *14*, 1588. [CrossRef]
75. Günther, T.; Dornberger, U.; Fritsche, W. Effects of Ryegrass on Biodegradation of Hydrocarbons in Soil. *Chemosphere* **1996**, *33*, 203–215. [CrossRef]
76. Kirk, J.L.; Klirnomos, J.N.; Lee, H.; Trevors, J.T. Phytotoxicity Assay to Assess Plant Species for Phytoremediation of Petroleum-Contaminated Soil. *Bioremediat. J.* **2002**, *6*, 57–63. [CrossRef]
77. Kirk, J.L.; Klironomos, J.N.; Lee, H.; Trevors, J.T. The Effects of Perennial Ryegrass and Alfalfa on Microbial Abundance and Diversity in Petroleum Contaminated Soil. *Environ. Pollut.* **2005**, *133*, 455–465. [CrossRef]

https://doi.org/10.3390/molecules31050890

78. Muratova, A.Y.; Dmitrieva, T.V.; Panchenko, L.V.; Turkovskaya, O.V. Phytoremediation of Oil-Sludge–Contaminated Soil. *Int. J. Phytoremediat.* **2008**, *10*, 486–502. [CrossRef]
79. Hussain, F.; Khan, A.H.A.; Hussain, I.; Farooqi, A.; Muhammad, Y.S.; Iqbal, M.; Arslan, M.; Yousaf, S. Soil Conditioners Improve Rhizodegradation of Aged Petroleum Hydrocarbons and Enhance the Growth of *Lolium multiflorum*. *Environ. Sci. Pollut. Res.* **2022**, *29*, 9097–9109. [CrossRef] [PubMed]
80. Tarigholizadeh, S.; Houshani, M.; Alizadeh, Z.; Ranjan, A.; Shende, S.S.; Rajput, V.D.; Sushkova, S.; Mandzhieva, S.; Bauer, T.; Minkina, T.; et al. Nanomaterial-Augmented Rhizoremediation: A Sustainable Strategy for Soil Bioremediation of Petroleum Hydrocarbons. *Adv. Sustain. Syst.* **2025**, *9*, e00558. [CrossRef]
81. Wang, Y.; Liu, R.; Dai, Y.; Chen, J.; Xiao, Y.; Bi, X.; Zou, P.; Li, S. Fire Phoenix Mediated Bioremediation of High-PAHs Contaminated Soil Enhanced by Graphene Oxide Application. *Environ. Technol. Innov.* **2025**, *37*, 103933. [CrossRef]
82. Li, L.; Liu, R.; Chen, J.; Tai, P.; Bi, X.; Zou, P.; Wang, Y.; Xiao, Y. Biotrophic Interactions between Plant and Endophytic Bacteria in Removal of PAHs and Cd from Contaminated Soils Enhanced by Graphene Oxide. *J. Clean. Prod.* **2023**, *417*, 137996. [CrossRef]
83. Hafez, E.M.; Alharbi, K.; Gharib, H.S.; Omara, A.E.-D.; Elatafi, E.; Hamada, M.M.; Rashwan, E.; Alshaal, T. Synergistic Effect of Sugarcane Bagasse and Zinc Oxide Nanoparticles on Eco-Remediation of Cadmium-Contaminated Saline Soils in Wheat Cultivation. *Plants* **2024**, *14*, 85. [CrossRef]
84. Hussain, F.; Hadi, F.; Rongliang, Q. Effects of Zinc Oxide Nanoparticles on Antioxidants, Chlorophyll Contents, and Proline in *Persicaria hydropiper* L. and Its Potential for Pb Phytoremediation. *Environ. Sci. Pollut. Res.* **2021**, *28*, 34697–34713. [CrossRef]
85. Oluyinka-Edunro, A.F.; Ogunsanwo, B.M.; Adeyemi, O.O. Green Synthesis of Zinc Oxide Nanoparticles UsingTerminaliaCatappa Leaf Extract and Its Application in Phytoremediation of Some Polycyclic Aromatic Hydrocarbons in Contaminated Soil. *FUOYE J. Pure Appl. Sci.* **2024**, *9*, 1–15.
86. Moses, H.; Odigure, J.O.; Otaru, A.J. Ecotoxicological Assessment of Nano-Remediated Crude Oil Contaminated Soil Using Zea Mays. *Results Eng.* **2024**, *23*, 102612. [CrossRef]
87. Bidi, H.; Fallah, H.; Niknejad, Y.; Barari Tari, D. Iron Oxide Nanoparticles Alleviate Arsenic Phytotoxicity in Rice by Improving Iron Uptake, Oxidative Stress Tolerance and Diminishing Arsenic Accumulation. *Plant Physiol. Biochem.* **2021**, *163*, 348–357. [CrossRef]
88. Ali, Q.; Zia, M.A.; Kamran, M.; Shabaan, M.; Zulfiqar, U.; Ahmad, M.; Iqbal, R.; Maqsood, M.F. Nanoremediation for Heavy Metal Contamination: A Review. *Hybrid Adv.* **2023**, *4*, 100091. [CrossRef]
89. Farooqi, A.; Ul Haq, E.; Johansen, A.; Ellegaard-Jensen, L.; Iqbal, M.; Yousaf, S.; Lackner, M. Plant Growth-Promoting Bacteria and Nanomaterials Synergism to Enhance Lolium Perenne Growth and Phytoremediation in Cadmium-Contaminated Soil. *Chem. Eng. J. Adv.* **2025**, *21*, 100695. [CrossRef]
90. Gil-Díaz, M.; González, A.; Alonso, J.; Lobo, M.C. Evaluation of the Stability of a Nanoremediation Strategy Using Barley Plants. *J. Environ. Manag.* **2016**, *165*, 150–158. [CrossRef]
91. Wang, J.; Fang, Z.; Cheng, W.; Yan, X.; Tsang, P.E.; Zhao, D. Higher Concentrations of Nanoscale Zero-Valent Iron (nZVI) in Soil Induced Rice Chlorosis Due to Inhibited Active Iron Transportation. *Environ. Pollut.* **2016**, *210*, 338–345. [CrossRef] [PubMed]
92. Sun, D.; Hu, J.; Bai, J.; Qin, H.; Wang, J.; Wang, J.; Lin, X. Arbuscular Mycorrhizal Fungus Facilitates Ryegrass (*Lolium perenne* L.) Growth and Polychlorinated Biphenyls Degradation in a Soil Applied with Nanoscale Zero-Valent Iron. *Ecotoxicol. Environ. Saf.* **2021**, *215*, 112170. [CrossRef] [PubMed]
93. Zhao, S.; Wang, W.; Chen, X.; Gao, Y.; Wu, X.; Ding, M.; Duo, L. Graphene Oxide Affected Root Growth, Anatomy, and Nutrient Uptake in Alfalfa. *Ecotoxicol. Environ. Saf.* **2023**, *250*, 114483. [CrossRef] [PubMed]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

https://doi.org/10.3390/molecules31050890

molecules

MDPI

Article

Assessment of Biodegradation Efficiency of Polychlorinated Biphenyls (PCBs) and Petroleum Hydrocarbons (TPH) in Soil Using Three Individual Bacterial Strains and Their Mixed Culture

Teresa Steliga [1,*,†], Katarzyna Wojtowicz [1,†], Piotr Kapusta [2] and Joanna Brzeszcz [2]

[1] Department of Reservoir Fluid Production Technology, Oil and Gas Institute–National Research Institute, ul. Lubicz 25 A, 31-503 Krakow, Poland; katarzyna.wojtowicz@inig.pl

[2] Department of Microbiology, Oil and Gas Institute-National Research Institute, ul. Lubicz 25 A, 31-503 Krakow, Poland; piotr.kapusta@inig.pl (P.K.); joanna.brzeszcz@inig.pl (J.B.)

* Correspondence: teresa.steliga@inig.pl

† These authors contributed equally to this work.

Academic Editor: Łukasz Chrzanowski

Received: 31 December 2019; Accepted: 5 February 2020; Published: 6 February 2020

check for updates

Abstract: Biodegradation is one of the most effective and profitable methods for the elimination of toxic polychlorinated biphenyls (PCBs) and total petroleum hydrocarbons (TPH) from the environment. In this study, aerobic degradation of the mentioned pollutants by bacterial strains *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, and *Rhodococcus* sp. IN306 and mixed culture M1 developed based on those strains at 1:1:1 ratio was analyzed. The effectiveness of individual strains and of the mixed culture was assessed based on carried out respirometric tests and chromatographic analyses. The *Rhodococcus* sp. IN306 turned out most effective in terms of 18 PCB congeners biodegradation (54.4%). The biodegradation index was decreasing with an increasing number of chlorine atoms in a molecule. Instead, the *Mycolicobacterium frederiksbergense* IN53 was the best TPH degrader (37.2%). In a sterile soil, contaminated with PCBs and TPH, the highest biodegradation effectiveness was obtained using inoculation with mixed culture M1, which allowed to reduce both the PCBs (51.8%) and TPH (34.6%) content. The PCBs and TPH biodegradation capacity of the defined mixed culture M1 was verified *ex-situ* with prism method in a non-sterile soil polluted with aged petroleum hydrocarbons (TPH) and spent transformer oil (PCBs). After inoculation with mixed culture M1, the PCBs were reduced during 6 months by 84.5% and TPH by 70.8% as well as soil toxicity was decreased.

Keywords: bacteria; polychlorinated biphenyls (PCBs); inoculation; biodegradation; total petroleum hydrocarbons (TPH); toxicological tests

1. Introduction

Polychlorinated biphenyls (PCBs) are synthetic chlorinated biphenyls, which were manufactured on a large scale for half a century and found various industrial applications, such as components of lubricants, transformer oils, dielectric fluids, and plasticizers. It is usually estimated that more than 1.5 million tons of PCBs have been produced worldwide [1]. These compounds are among the most persistent xenobiotic pollutant classes. They remain in the environment for a long time due to their low reactivity and high chemical stability in various conditions. Because of this, their production and application was banned in 1980 [2–4]. However, to a large extent they were dispersed and resulted in vast pollution of all ecosystems [5]. The most alarming adverse PCBs property is their tendency to bioaccumulate in lipid tissues and organic components of the soil and in fat tissues of animals and

people [6]. PCBs molecules consist of a biphenyl core containing 1 to 10 chlorine atoms substituted in various positions, as a result of which there are 209 various congeners featuring different degrees of chlorination and substitution. Congeners that have benzene rings in one plane and non-substituted *ortho*-positions are the most persistent and feature the highest toxicity [7,8]. The number of chlorine atoms and their arrangement in PCBs molecule are important factors for aerobic biodegradation by oxidative enzymes. The neighbouring non-chlorinated carbon atoms allow for hydroxylation of the ring and formation of arene oxide intermediate compounds, facilitating aromatic ring splitting. In general, PCB congeners with four or less chlorine atoms in a molecule are easily degraded, and with five or more, are more resistant. PCBs are usually first oxidized by biphenyl dioxygenase (BphA) to produce cis-dihydrodiol semi-products, which are subjected to ring *ortho*- or *meta*-cleavage-producing chlorobenzoates [1,9,10].

Bacteria using biphenyl (BP) as a source of carbon and energy play a crucial role in the PCBs degradation. The most active bacteria capable of PCBs metabolizing and most frequently reported in the literature include: *Achromobacter* sp. [6,11], *Acinetobacter* sp. [12], *Bacillus* sp. [13,14], *Burkholderia* sp. [15] *Janibacter* sp. [16], *Mycobacterium* sp. [17], *Ochrobactrum* sp. [18,19], *Pseudomonas* sp. [20–25], and *Rhodococcus* sp. [19,26–33].

The removal of toxic and mutagenic, hydrophobic chlorinated aromatic compounds, mainly PCBs, from the environment is one of most important goals of environmental biotechnology. The cleaning of soil polluted with PCBs using conventional methods such as thermal desorption, the plasma method, combustion methods, dechlorination, or soil washing method are frequently too expensive [34]. Biological methods are an alternative strategy for PCBs removal and now continuous development of bioremediation techniques is observed [19,31,35–38]. Bioremediation is considered an effective and profitable method for PCBs removal from the soil and its main strategies comprise bioaugmentation and/or biostimulation, which means the introduction of bacterial strains, degrading PCBs individually or as a mixed culture, and nutrients, respectively. In order to avoid antagonistic interaction with foreign natural soil microbiota on foreign microbial cultures not adapted to a given environment from a commercial mixed culture, it is preferable to prepare bacterial consortia based on previously isolated, selected and multiplied indigenous microorganisms. They do not upset the biological equilibrium, creating a high probability of survival in polluted biocenosis, and hence guarantee an appropriately high biodegradation potential [39–42]. The bioaugmentation may fail due to the following reasons: cell stress, lack of nutrients, existence of more preferred source of carbon as a substrate, competition, or inappropriate biological availability. The studies prove that the application of mixed bacterial culture better reflects real conditions as compared with the application of a pure bacterial culture [43]. There is little information in the literature about the application of bacterial consortia to clean soils historically contaminated by PCBs. Havel and Reineke [44] applied a consortium containing isolates belonging to *Pseudomonas* sp., and Egorova et al. [45] used a combination of strains *Rhodococcus ruber* P52 and *Microbacterium sp* B51. The last study carried out by Horváthová et al., [19] showed clear degradation of highly chlorinated PCB congeners by means of bacterial consortia prepared on the basis of four bacterial strains (*Achromobacter xylosoxidans, Stenotrophomonas maltophilia, Ochrobactrum anthropi and Rhodococcus ruber*) at various biomass ratios, isolated from deposits historically contaminated with PCBs. Satisfactory effects of PCBs biodegradation (60% after 8 weeks) were achieved using as an inoculum of a mixture of four bacterial strains: *Bacillus* sp., *Achromobacter* sp., *Pseudomonas stutzeri*, and *Bacillus subtilis* [38].

Apart from biological factors, the effectiveness of PCBs group biodegradation is affected by environmental factors, such as temperature, pH, oxygen availability, humidity, soil matrix properties, nutrients content [46], a surfactant addition [47], as well as pollution-related factors (physical and chemical properties of polychlorinated biphenyls, xenobiotic concentration, pollutant age, number of chlorine atoms in a biphenyl ring, and a spatial PCBs structure) [48].

Frequent historical soil contamination with polychlorinated biphenyls (PCBs) originating from a transformer oil is accompanied by the existence of total petroleum hydrocarbons (TPH) [31].

Biodegradation of petroleum pollutants in a historically polluted soil originating from waste pits with the use of a defined mixed culture based on non-pathogenic indigenous bacterial strains was studied by the authors, which is proven by numerous papers [39,49–51].

The studies were aimed at the determination of aerobic biodegradation effectiveness PCBs and TPH with the use of three single bacterial strains isolated from contaminated natural environments and a mixed culture developed on the basis of these strains at 1:1:1 ratio. The studies were carried out on a sterile soil polluted with (a) PCBs, (b) TPH, and (c) both PCBs and TPH. The effectiveness of individual strains and of the mixed culture was assessed based on carried out respirometric tests and chromatographic analyses of PCBs and TPH in the examined soil. The PCBs and TPH biodegradation capacity of the defined mixed culture was verified under semi-technical conditions by means of the ex-situ prism method on a non-sterile soil weathered polluted with petroleum hydrocarbons (TPH) and spent PCBs containing transformer oil. The course of PCBs and TPH biodegradation in the soil was monitored using chromatographic determination of the content of individual polychlorinated biphenyls and petroleum hydrocarbons as well as toxicological studies using a set of new-generation toxicological tests, in which the bioindicators belong to various taxonomic groups (bacteria, crustaceans, and higher plants).

2. Results

2.1. Bacterial Strains Used in this Study

The 16S rRNA gene sequences of strains used in this study have been deposited in the NCBI GenBank database and are available under the following accession numbers: JN572675 (*Mycolicibacterium frederiksbergense* IN53), KT923311 (*Rhodococcus erythropolis* IN129), and KX058399 (*Rhodococcus* sp. IN306). Table 1 presents the similarity of our strains with their closest relatives based on the available 16S rRNA gene sequences deposited in the NCBI GenBank database (comparison was made both for individual nucleotide sequences and the whole genomes). For the genomes, annotated genes coding putative enzymes responsible for degradation of polycholorinated biphenyls and other hydrocarbons, are shown. The presented data show that genomes of organisms closely related to strains IN53, IN129, and IN306 contain genes responsible for the catabolism of both aliphatic and aromatic hydrocarbons. All genome sequences contain genes coding for the third enzyme of the biphenyl degradation pathway (Table 1), i.e., 2,3-dihydroxybiphenyl 1,2-dioxygenase (*bphC*). Moreover, each genome contains more than one copy of this gene (Table 1). In the genome sequence of *Rhodococcus jostii* RHA1, all genes are related to the enzyme responsible for the initial stages of biphenyl degradation. Namely, genes encoding for biphenyl 2,3-dioxygenase (*bphA*, found all four components), biphenyl-2,3-dihydrodiol 2,3-dehydrogenase (*bphB*), and 2,3-dihydroxybiphenyl 1,2-dioxygenase (*bphC*) (Table 1). The genome sequence of *Mycobacterium* sp. YC-RL4 contains a gene coding for the sub-unit alpha of ring-hydroxylating dioxygenase (probably the first component of biphenyl 2,3-dioxygenase, Table 1), and there are genes coding the same enzyme and the enzyme identified as the biphenyl 2,3-dioxygenase in the genome of *Rhodococcus erythropolis* X5. In addition, all the analyzed genome sequences contain *alkB*, the gene coding for alkane 1-monooxygenase. This enzyme is responsible for the hydroxylation, which is the first reaction in the alkane degradation pathway.

For the tree construction, the 16S rRNA gene sequences of the IN53, IN129 and IN306 were aligned using SINA Aligner, [52] and further analyzed using ARB program [53] with the SILVA SSU Ref NR99 release 132 database. Phylogenetic reconstructions were performed with full-length sequences using Neighbour Joining with Jukes-Cantor correction and Maximum Likelihood (RaxML, model: GTRGAMMA). In all cases, a base frequency filter was applied. The selected tree represents a consensus topology between the different reconstructions (Figure 1).

Table 1. Genes involve in biphenyl and hydrocarbon degradation, found in genomes of the closest relatives of IN53, IN129 and IN306.

Strain	The Closest Relative Based on 16S rRNA Accession Number, (% of Identity) *	The Closest Relative for Which the Genome Sequence Is Available in NCBI GenBank, Accession Number, (% of Identity)	Gene Encoding for the Enzymes Degrading Hydrocarbons
Mycolicibacterium frederiksbergense IN53 JN572675	*Mycolicibacterium frederiksbergense* DSM 44346 (typical strain) NR_025393.1 99.58%	*Mycobacterium* sp. YC-RL4, CP015596, (99.24%)	Alkane 1-monooxygenase (2 copies), pentachlorophenol monooxygenase, 2,3-dihydroxybiphenyl 1,2-dioxygenase (2 copies), ring-hydroxylating dioxygenase alpha subunit
Rhodococcus erythropolis IN129 KT923311	*Rhodococcus erythropolis* IN104 KT923338 100%	*Rhodococcus erythropolis* X5, CP044284, (99.93%)	Alkane 1-monooxygenase (5copies), pentachlorophenol monooxygenase, phenol/toluene hydroxylase, biphenyl 2,3-dioxygenase, 2,3-dihydroxybiphenyl 1,2-dioxygenase (3 copies), ring-hydroxylating dioxygenase alpha subunit
Rhodococcus sp. IN306 KX058399	*Rhodococcus* sp. S2-17 KY765341 99.30%	*Rhodococcus jostii* RHA1, CP000431, (98.75%)	Alkane 1-monooxygenase, pentachlorophenol monooxygenase, phenol/toluene hydroxylase, biphenyl 2,3-dioxygenase, 2,3-dihydrodihydroxybiphenyl 2,3-dehydrogenase, 2,3-dihydroxybiphenyl 1,2-dioxygenase (5 copies),

* the presented data concerns the data gathered in the NCBI GenBank database (November 2019).




Figure 1. Phylogenetic consensus tree of the stains IN129, IN53 and IN306 with closely related sequences from the NCBI GenBank database. GenBank accession numbers are shown in the Figure. The scale bar corresponds to 10% estimated sequence divergence.

2.2. Assessment of PCB and TPH Biodegradation Effectiveness in Sterile Soils

2.2.1. Respirometric Tests

An increase in the microbiological activity in a reactive environment proves that microorganisms used PCBs and TPH as a source of carbon and energy. An increased biological activity in the system results in a growth of oxygen consumption by the studied bacterial strains: *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306, and mixed culture M1 (prepared from individual bacterial strains at 1:1:1 ratio), over time causing biodegradation of analytes.

Figure 2 presents the dependence of the consumed oxygen amount on the experiment duration for samples of sterile Soils A, B, and C inoculated with strains of bacteria IN53, IN129, IN306, and mixed culture M1 for control samples. The respirometric tests performed on a sterile soil contaminated with PCBs (Soil A) inoculated with strains IN53, IN129, IN306 and mixed culture M1 showed that the process of biodegradation is most intensive during inoculation with *Rhodococcus* sp. IN306, which is proved by the amount of consumed oxygen after 30 days of testing, approx. 3,274 ± 196 mg O_2/dm^3. A slightly lower oxygen consumption during the experiment was recorded in the sample inoculated with mixed culture M1 (3,161 ± 174 mg O_2/dm^3) and with *Rhodococcus erythropolis* IN129 (3,020 ± 172 mg O_2/dm^3). *Mycolicibacterium frederiksbergense* IN53 turned out to be least effective in terms of biodegradation potential of PCBs group compounds, because the oxygen consumption after 30 days of testing amounted to 2258 ± 124 mg O_2/dm^3. The results of respirometric tests performed on the sample of Soil B showed that the strain IN53 features the highest biodegradation potential in terms of TPH content reduction in the soil (4235 ± 241 mg O_2/dm^3). Slightly lower amounts of consumed oxygen after the test completion were recorded for samples inoculated with mixed culture M1 (3975 ± 226 mg O_2/dm^3) and with the IN129 strain (3636 ± 204 mg O_2/dm^3). The lowest biological activity after completion of the respirometric test was observed in the sample inoculated with the IN306 strain (3193 ± 179 mg O_2/dm^3). Results of respirometric tests carried out on the sample of soil contaminated both with PCBs and TPH (Soil C) showed that the biodegradation is most intensive in the sample inoculated with mixed culture M1 (7136 ± 357 mg O_2/dm^3). Inoculation of Soil C with strains IN53, IN129, and IN306 also resulted in an increase in the microbiological activity in the system, which is proved by the amounts of consumed oxygen after 30 days of the experiment, 6817 ± 348 mg O_2/dm^3, 6656 ± 339 mg O_2/dm^3, and 6466 ± 336 mg O_2/dm^3, respectively.

2.2.2. Assessment of PCBs and TPH Biodegradation Based on Chromatographic Analyses

PCB biodegradation potential of studied bacterial strains and mixed culture in sterile soil contaminated solely with PCBs (Soil A) was assessed by chromatographic analyses. Figure 3 and Table S1 present the content of studied PCB congeners in the PCBs contaminated soil (Soil A) before and after 30-day inoculation with bacterial strains: *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306, and mixed culture M1.




Figure 2. Oxygen consumption by microorganisms Mycolicibacterium frederiksbergense IN53, Rhodococcus erythropolis IN129, Rhodococcus sp. IN306 and mixed culture M1 to degrade (**a**) PCBs (sterile Soil A contaminated PCB), (**b**) TPH (sterile Soil B contaminated TPH), and (**c**) PCBs and TPH (sterile Soil C contaminated PCB and TPH) in the biodegradation process. Control-sterile uncontaminated "pure" soil inoculated with bacterial strains and mixed culture.




Figure 3. Residual PCB congeners content in sterile Soil A contaminated with PCBs inoculated by Mycolicibacterium frederiksbergense IN53, Rhodococcus erythropolis IN129, Rhodococcus sp. IN306 and mixed culture M1 after a 30-day biodegradation process, (repetition number $n = 8–10$, $p < 0.05$). Control-non-inoculated Soil A.

The results of chromatographic analyses showed that *Rhodococcus* sp. IN306 features the highest biodegradation potential with respect to the studied PCB congeners. The inoculation of soil A with the strain IN306 led to a significant reduction of the PCB content from 13,100 to 5,970 μg/kg of dry mass (54.4%). Satisfactory results were achieved also for mixed culture M1 from 13,100 μg/kg to 6309 μg/kg of dry mass (46.2%). *Mycolicibacterium frederiksbergense* IN53 turned out to be the least effective, for which the degree of biodegradation after 30 days of test duration was 35.9%. The PCB28 congener (containing three chlorine atoms in a molecule) turned out to be the most easily biodegradable, in which content in the sample after 30 days of testing was reduced from 494 to 202 μg/kg of dry mass, (IN53), 162 μg/kg (IN129), 123 (IN306), and 134 (M1), which is 59.1%, 67.2%, 75.1%, and 72.9%, respectively. Congeners containing four chlorine atoms in a biphenyl molecule, including: PCB52, PCB77, and PCB81 were also degraded to a comparable extent: PCB52 (46.8–62.4%), PCB77 (55.9–68.5%), and PCB81 (54.1–68.1%). PentaCBs (PCB101, PCB105, PCB114, PCB118, PCB123, and PCB126) were reduced as follows: 38.5–42.8% (IN53), 40.8–43.4% (IN129), 52.3–57.2% (IN306), and 43.9–47.4% (M1). The PCB105 congener turned out to be the most easily biodegradable among PentaCBs, in which thecontent in the sample was reduced from 733 μg/kg of dry mass to 419 μg/kg (IN53), 414 μg/kg (IN129), 313 μg/kg (IN306), and 385 μg/kg of dry mass (M1). The PCB congener containing five chlorine atoms in a molecule most difficult to degrade was PCB101 (di-*ortho* structure), because the degree of its biodegradation as a result of inoculation was: 38.5% (IN53), 40.1% (IN129), 52.3% (IN306), and 49.8% (M1). The HexaCBs content reduction ranged from 23.4 to 49.3%. Among HexaPCBs, the PCB157 inoculated with *Rhodococcus* sp. (IN306), showed the highest reduction of content after 30 days of testing—by 51.2%, while the most difficult to degrade were: PCB138 and PCB153, for which the biodegradation degrees were 46.9% and 46.5%, respectively. HeptaCBs (PCB180 and PCB189) turned out to be most difficult to degrade, for which the highest biodegradation degree after inoculation with IN306 was 31.7% and 38.7%.

Chromatographic analyses were carried out on sterile soil contaminated solely with TPH (Soil B) before and after 30-day inoculation with: *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306, and mixed culture M1, which showed that the strain *Mycolicibacterium frederiksbergense* IN53 features the highest biodegradation potential with respect to TPH. Table S2 and Figure 4 present a comparison of the content reduction for TPH; unidentified hydrocarbons; alkanes; $\Sigma nC_{10}–nC_{22}$, $\Sigma nC_{23}–nC_{40}$; and individual n-alkanes in Soil B after inoculation with strains IN53, IN129, IN306, and mixed culture M1.

The TPH content in Soil B after 30 days of Soil B inoculation with the *Mycolicibacterium frederiksbergense* IN53 was reduced by 37.2% (from 12,515 to 7863 mg/kg of dry mass). A satisfactory reduction of TPH content from 12,515 to 8309 mg/kg of dry mass (33.6%) was recorded during inoculation with mixed culture M1. The TPH content in Soil B samples inoculated with IN129 and IN306 strains was reduced by 28.5% and 24.9%, respectively. The easiest to degrade were aliphatic hydrocarbons with chain lengths $nC_{10}–nC_{22}$, within a range of: 42.7–48.7% (IN53), 38.7–41.5% (IN129), 24.4–33.4% (IN306), and 39.5–46.7% (M1). In turn, heavier homologues ($nC_{23}–nC_{40}$) were degraded less efficiently and their content was reduced by 21.0% (IN129), 15.2% (IN306), and 31.8% (M1) in samples inoculated with IN129, IN306 and M1, respectively. The best effects of heavy hydrocarbons biodegradation were obtained during Soil B inoculation with the *Mycolicibacterium frederiksbergense* IN53 strain, equal to 39.5%.

A satisfactory effect of n-alkanes biodegradation as a result of Soil B inoculation is proven by indices of C_{17}/Pr and C_{18}/Ph biodegradation, which were substantially reduced (Table S2). The highest reduction was recorded during inoculation with the IN53 strain (C_{17}/Pr from 0.818 to 0.473 and C_{18}/Ph from 1.675 to 0.985) and with mixed culture M1 (C_{17}/Pr from 0.818 to 0.522 and C_{18}/Ph from 1.675 to 1.029).




Figure 4. Residual n-alkanes content in sterile Soil B contaminated with TPH inoculated by Mycolicibacterium frederiksbergense IN53, Rhodococcus erythropolis IN129, Rhodococcus sp. IN306 and mixed culture M1 after a 30-day biodegradation process, (repetition number $n = 8–10$, $p < 0.05$). Control non-inoculated Soil B.

The comparison of PCBs and TPH content in sterile Soil C after inoculation with strains IN53, IN129, IN306, and mixed culture M1 is presented in Tables S1 and S2 and in Figure 5. Results of chromatographic analyses proved that during the inoculation of sterile Soil C with mixed culture M1 a high reduction of PCBs content was achieved from 13,100.0 to 6309 µg/kg of dry mass (51.8%) as well as of TPH content from 12,365 to 8085 mg/kg of dry mass (34.6%). The TriCBs content after 30 days of inoculation was reduced by 51.2% (*Mycolicibacterium frederiksbergense* IN53), 61.1% (*Rhodococcus erythropolis* N129), 69.4% (*Rhodococcus* sp. IN306), and 70.4% (M1). TetraCBs were biodegraded to a comparable extent: 43.3% (IN53), 53.9% (IN129), 62.3% (IN306), and 62.4% (M1). PentaCBs and HexaCBs were best degraded by the mixed culture M1, by 52.9% and 46.8%. HeptaCBs were definitely the most difficult to biodegrade among PCBs. Their content as a result of inoculation with strains IN53, IN129, IN306, and mixed culture M1 was reduced from 1680 µg/kg of dry mass to: 1449 µg/kg (13.8%), 1256 µg/kg (25.2%), 1184 µg/kg (29.5%), and 1059 µg/kg (37.0%), respectively.

Chromatographic analyses of petroleum hydrocarbons (TPH) showed that aliphatic hydrocarbons nC_{10}–nC_{22} were easiest to degrade, for which the biodegradation degree after 30 days of inoculation ranged as follows: 41.1–48.2% (IN53), 35.3–39.1% (IN129), 26.2–34.2% (IN306), and 39.1–45.4% (M1). Long chain aliphatic hydrocarbons nC_{23}–nC_{40} were reduced by 32.5% (IN53), 17.5% (IN129), 13.8% (IN306), and 35.4% (M1) accordingly. Biodegradation of unidentified hydrocarbons reached: 29.4% (IN53), 21.5% (IN129), 18.0% (IN306), and 30.0% (M1).

2.3. Assessment of PCBs and TPH Biodegradation in Non-Sterile Soil-the Ex-Situ Prism Method

The effectiveness of mixed culture M1 was verified carrying out the process of biodegradation in a non-sterile Soil D collected from waste pit G-44 with a TPH content of 22,126 mg/kg of dry mass, which was contaminated with aged transformer oil containing PCBs up to 8752 µg/kg of dry mass. Figure S1 illustrates the percentage share of identified pollutants in the sample (Soil D) comprising: (a) PCBs, (b) TPH.




Figure 5. Comparison of biodegradation (%) in sterile Soil C contaminated with both PCBs and TPH after inoculation with Mycolicibacterium frederiksbergense IN53, Rhodococcus erythropolis IN129, Rhodococcus sp. IN306, and mixed culture M1: (**a**) polychlorinated biphenyls; (**b**) petroleum hydrocarbons (repetition number $n = 8–10$, $p < 0.05$).

Comparison of identified pollutants content in the studied soil (Soil D) and after 2 (Soil D2), 4 (Soil D4), and 6 (Soil D6) months of inoculation with mixed culture M1 is presented in Figure 6: (a) PCB congeners, (b) n-alkanes. As a result of Soil D inoculation with mixed culture M1 after 6 months of the process duration, the content of identified PCBs was reduced from 8752 µg/kg of dry mass to 1357 µg/kg of dry mass (84.5%). Results of Soil D2 chromatographic analyses show that the PCB28 content was reduced from 1396 µg/kg to 371 µg/kg of dry mass (73.4%). The tetraCBs content (PCB52 and PCB77) decreased by 56.4 and 65.6% respectively. The biodegradation of pentaCBs was similar and ranged from 49.8 to 53.6%. In this case, the *di-ortho* PCB101 was an exception, in which biodegradation degree was much lower and after 2 months of the process amounted to 45.2%. *Di-ortho* congeners turned out to be most durable among hexaCBs, in which content decreased by 40.1% (PCB138) and 39.3% (PCB156). The other congeners having six chlorine atoms were reduced by 44.8 to 45.3%. The PCB180 congener turned out to definitely be the most difficult to degrade, which has seven chlorine atoms and a *di-ortho* spatial structure (29.7%).

After 4 months (Soil D4), the content of low-chlorinated biphenyls: PCB28, PCB52, and PCB77 decreased by 89.7%, 76.1%, and 81.2%, respectively. PCB101 was reduced by 69.5%, while the other PentaCBs between 72.1 and 73.3%. The HexaCBs (PCB 138, PCB 153, PCB 156, and PCB 167) content reduction ranged from 65.9 to 69.8%, while the PCB180 biodegradation in soil D4 was 40.8%. The process of Soil D inoculation with mixed culture M1 was finished after 6 months, and the recorded reduction of individual identified congeners was: PCB28, PCB52, and PCB77 (87.7–94.6%); pentaCBs (83.3–86.7%); HexaCBs (81.0–84.2%); and PCB180 (58.6%). The inoculation of non-sterile Soil D with mixed culture M1 during 6 months resulted in a TPH content reduction from 22,126 to 6470 mg/kg of dry mass. The degree of TPH content reduction during the process of inoculation with mixed culture M1 was as follows: after 2 months, 41.4%; after 4 months, 60.7%; and after 6 months, 70.8%.




Figure 6. Residual content of (**a**) PCB congeners and (**b**) n-alkanes in non-sterile soil contaminated with both PCBs and petroleum hydrocarbons (Soil D) after 2 (Soil D2), 4 (Soils D4) and 6 (Soil D6) months of inoculation with mixed culture M1 (ex-situ prism method), (repetition number $n = 7–10$, $p < 0.05$). Control non-inoculated Soil D.

The chromatographic analysis showed that during inoculation with mixed culture M1, the biodegradation of aliphatic hydrocarbons $nC_{11}–nC_{22}$ was as follows: Soil D2 (47.2–76.2%), Soil D4 (77.5–88.7%), and Soil D6 (87.2–93.2%). Hydrocarbons from the $nC_{23}–nC_{40}$ range were also degraded to a satisfactory degree: Soil D2 (13.8–41.2%), Soil D4 (20.1–62.9%), and Soil D6 (25.6–72.4%). The content of unidentified hydrocarbons in Soil D6 was reduced by 68.1%. The assessment indices of biodegradation degree proved a high level of biological decomposition of n-alkanes, because they were visibly reduced in Soil D6: nC_{17}/Pr from 1.131 to 0.192, nC_{18}/Ph from 2.081 to 0.402.

The course of PCBs and TPH biodegradation process during the carried out inoculation with mixed culture M1 of non-sterile Soil D is described by equation (2). Individual coefficients of equation (2): k, $(C/C_x)_0$, and the correlation coefficient (r^2) are specified in Table S3. A comparison of the biodegradation course is presented in a graphical form for PCBs, TriCBs, TetraCBs, PentaCBs, HexaCBs, HeptaCBs (Figure 7a), and for TPH, $\Sigma nC_{10}–nC_{22}$, and $\Sigma nC_{23}–nC_{40}$ (Figure 7b).




Figure 7. Comparison of content reduction (**a**) in PCB-209 normalized PCBs, TriCB, TetraCBs, PentaCBs, HexaCBs, and HeptaCBs (**b**) in hopane-normalized TPH, ΣnC_{10}–nC_{22}, and ΣnC_{23}–nC_{40} during inoculation of non-sterile soil D with mixed culture M1 (ex-situ prism method). (repetition number n = 7–10, $p < 0.05$).

The first-order biodegradation constants (k) during inoculation of non-sterile Soil D with mixed culture M1 are as follows: PCBs (0.0109 d^{-1}), TriPCB (0.0170 d^{-1}), TetraCBs (0.0110 d^{-1}), PentaCBs (0.0095 d^{-1}), Hexa CBs (0.0082 d^{-1}), and HeptaCBs (0.0050 d^{-1}). Instead, for petroleum pollutants, they are: TPH (0.0070 d^{-1}), Σ nC_{11}–nC_{22} (0.0130 d^{-1}), and Σ nC_{23}–nC_{40} (0.0049 d^{-1}). The presented results allow comparing the kinetics of individual pollutant groups biodegradation during inoculation with mixed culture M1 and prove its biodegradation capacity both of contamination with polychlorinated biphenyls (PCBs) and total petroleum hydrocarbons (TPH).

Ecotoxicological Assessment

During petroleum hydrocarbons biodegradation, metabolites of diversified or poorly recognized biological activity can originate as a result of chemical and microbiological changes. Because of that, it is recommended to carry out toxicological tests, which enable monitoring the toxicity of PCBs and TPH biodegradation during inoculation with mixed culture M1 in a non-sterile soil (Soil D) under semi-technical conditions by the ex-situ prism method. Toxicological tests on the trophic level of reducers were carried out by means of Microtox SPT test using luminescent bacteria *Vibrio fischeri*. The determined EC_{50} concentration causing 50% inhibition of test bacteria luminescence was 6.5% vol., which corresponds to a toxicity expressed in toxicity units of around TU = 15.4. Toxicological analyses carried out during Soil D inoculation with mixed culture M1 showed a decrease of its toxicity during 6 months to a level of TU = 3.9, at which the cleaned soil may be classified as low-toxic (Figure 8).

The next test, which was applied to estimate Soil D toxicity on the consumer level during inoculation with mixed culture M1, was the Ostracodtox kit based on the use of *Heterocypris incongruens* organisms, which are more sensitive to pollutants than the *Vibrio fisheri* bacteria. Figure 8 presents the test results. The toxicity (TU) of raw soil (Soil D) was 16.9 and was gradually going down to TU = 11.3 (after 3 months) and TU = 4.9 (after 6 months).

Moreover, an innovative MARA test was performed, estimating the environmental risk and using 11 test strains, which featured various sensitivities. The toxicity calculated based on the mean toxic concentration MCT mean, expressed in toxicity units, was 19.6. During the carried out PCBs and TPH biodegradation, its reduction to 5.2 was recorded (Figure 8).




Figure 8. Comparison of toxicity test results expressed in toxicity units (TU) from non-sterile soil during inoculation with mixed culture M1 (ex-situ prism method) ($n = 3$, $p < 0.05$): Soil D—raw soil; Soil D3—soil after 3 months; Soil D6—soil after 6 months.

In the toxicological tests on the producers level, the Phytotoxkit test was used, in which the tested plants were: *Lepidium sativum*, *Sorghum saccharatum*, and *Sinapis alba*. After the conversion of toxic concentrations to toxicity units (TU), considering the plant germination as a criterion of toxicity assessment, it was found that during the carried out biodegradation process of pollutants contained in the tested soil samples, it was gradually decreasing from 10.5 to 3.0 (*Lepidium sativum*), from 9.5 to 3.1 (*Sorghum saccharatum*), and from 8.5 to 3.0 (*Sinapis alba*). For the second tested parameter (root growth inhibition), the calculated TU value for the tested plants in the sample (Soil D) ranged from 10.1 to 15, while after inoculation with mixed culture M1 (Soil D6), it went down to a level of 3.6–4.6. The carried out tests showed that *Lepidium sativum* was the plant most sensitive to pollutants contained in the tested soil samples (Figure 8).

The Ames test series was applied to find mutagenic and carcinogenic compounds in raw and cleaned soil. A standard strain TA-100 (*Salmonella typhimirium*) was used. Its mutations exist on a histidine operon, which is not capable of amino acid synthesising. It was observed that the application of an activator (microsomal S9 fraction) had an insignificant impact on the growth of reversible mutations number. Results of tests on raw samples and in the consecutive months of soil bioremediation are presented in Figure 9 as changes of revertants number versus the pollutant concentrations. In the case of (Soil D) samples, which are considered mutagenic, the mutagenecity coefficient ranged from 7.85 to 9.85 (Figure 9a). During the carried out process of pollutants (PCBs and TPH) biodegradation, the mutagenecity coefficient was gradually decreasing: Soil D3 from 4.32 to 6.19 (Figure 9b), Soil D6 from 0.68 to 1.86 (Figure 9c), i.e., to a level at which the number of revertants induced without histidine was slightly higher (less than twice) than the number of spontaneous mutants in the control sample. Therefore, the samples cannot be classified as mutagenic.




Figure 9. Influence of PCBs and TPH concentration on revertants number ($n = 3$, $p < 0.05$): (**a**) raw soil (Soil D); (**b**) soil after 3-month biodegradation (Soil D3); (**c**) soil after 6-month biodegradation (Soil D6); fold increase over the baseline (mutagenicity index).

3. Discussion

Polychlorinated biphenyls (PCBs), despite chemical stability and toxicity, can be degraded under aerobic conditions or at least reduced to low-chlorinated congeners by microorganisms possessing genes encoding for biphenyl dioxygenase responsible for the onset of a specific metabolic degradation pathway [3,54]. The literature studies prove that the best PCBs biodegradation properties are featured by bacteria isolated from the soil/deposits historically PCBs contaminated: *Microbacterium oleivorans*, *Stenotrophomonas maltophilia*, *Brevibacterium* sp., *Ochrobactrum anthropi*, *Pseudomonas mandelii*, *Rhodococcus* sp., *Achromobacter xylosoxidans*, *Stenotrophomonas* sp., *Ochrobactrum* sp. [6], *Sinorhizobium meliloti* [35,55], *Rhodococcus ruber* and *Rhodococcus pyridinivoran* [33], *Rhodococcus* sp. and *Stenotrophomonas maltofilia* [56] *Bacillus* sp., *Achromobacter* sp., *Pseudomonas stutzeri*, [38]. Results of genes sequencing for bacterial strains isolated from the deposit polluted with transformer oil showed that *Rhodococcus* sp., *Pseudomonas* sp., *Pseudoxanthomonas* sp., *Agromyces* sp., and *Brevibacillus* sp. were the prevailing bacteria decomposing PCBs [57]. *Rhodococcus jostii* RHA1 is one of the best characterized PCBs degraders, featuring a broad PCBs degradation range and its catabolic potential and adaptation to the soil environment suggest that it can be appropriate for bioremediation of polluted Soil D [31].

Research conducted to determine the biodegradability of PCB congeners in sterile contaminated (i.e., one that does not contain other living organisms) reference soil PCB SQC068 by single bacterial strains isolated from contaminated natural environments (*Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306) and the mixed culture M1 prepared on the basis of these strains at ratio 1:1:1, confirm the anticipated ability of these strains to metabolize biphenyl and its derivatives. The mixed culture M1 confirms the anticipated ability of these strains to metabolize biphenyl and its derivatives. It was possible to show that each of the tested bacterial stains, as well as their mixture, degraded PCB congeners, where the *Rhodococcus* sp. IN306 strain turned out to be most effective, removing more than 54.4% of PCBs. *Rhodococcus* sp. IN306 strain is very closely related to the *Rhodococcus jostii* RHA-1 species (Figure 1, Table 1), which is a well-known degrader of biphenyl and its polychlorinated derivatives [31,58–60]. The other two strains featured a lower degree of degradation, but the obtained results confirm the observations made during previous metabolic tests, that they are capable of carrying out catabolic processes of biphenyl and its derivatives. In the case of *Rhodococcus erythropolis* IN129 [26,29], it is not surprising, because other species belonging to the *Rhodococcus genus* [27,30,32,33,61,62] have such abilities; in the case of *Mycolicibacterium frederiksbergense* IN53, this means that this strain has even more potential for degrading xenobiotics than previously thought. The higher biodegradation efficiency of polychlorinated biphenyls in the case of *Rhodococcus* sp. IN306 compared to the mixed culture M1 most likely results from competition between individual strains for the same substrates.

Moreover, the obtained results of chromatographic analyses show that with an increasing number of chlorine atoms in a PCB molecule, the degree of their degradation decreases (Table S1), which is

consistent with other studies [2,9,54]. PCB congeners containing up to four chlorine atoms in a molecule easily decompose as a result of carried out inoculation with all tested bacterial strains and mixed culture M1 developed based on these strains. Highly chlorinated biphenyls (more than four chlorine atoms in a molecule) are definitely more difficult to degrade [63], in the case of which satisfactory results were obtained for samples inoculated with *Rhodococcus* sp. IN306 and mixed culture M1. For example, the content of PCB-28 congener with three chlorine atoms was reduced as a result of *Rhodococcus* sp. IN306 action by 75.1%, while the content of pentaCBs, hexaCBs, and heptaCBs was reduced by 55.3%, 49.2%, and 39.3%, respectively. It should be noticed that in the case of polychlorinated biphenyls biodegradation, their spatial structure is also important, apart from the number of chlorine atoms in a molecule [64]. It was observed that so-called *di-ortho* congeners (two chlorine atoms in an *-ortho* position) are more difficult to biodegrade as compared with *non-ortho* or *mono-ortho* PCBs with the same number of chlorine atoms in the biphenyl ring [9,65]. 5 *di-ortho* PCB congeners (PCB52, PCB101, PCB138, PCB153, and PCB180) were determined in the studied sample (Soil A). For example, as a result of standard PCB Soil A inoculation with the *Rhodococcus* sp. IN306 strain, the reduction of *di-ortho* PCB52 congener (tetrachlorobiphenyl) was 62.4%, while for PCB77 and PCB81 congeners it was approx. 68.5% and 68.1%.

In the case of biodegradation of petroleum hydrocarbons (TPH) in sterile Soil B, strain *Mycolicibacterium frederiksbergense* IN53 showed the best degradability (37.2%), better than mixed culture M1 (33.6%). There was probably a similar situation to this in the case of soil A, and the effect of competition between strains for the same source of carbon and energy could occur. Then the *Rhodococcus* sp. IN306 turned out to be the weakest degrader, most likely due to specialization of its enzymatic apparatus to metabolize biphenyl and its derivatives.

Moreover, the obtained results of chromatographic analyses enabled determination of biodegradability of individual hydrocarbon groups (Table S2). The rate of individual hydrocarbon groups removal was arranged in a decreasing order nC_{10}–nC_{22} > nC_{23}–nC_{28} > nC_{29}–nC_{40}. This removal scheme is probably related to the chemical structure of alkanes. Alkanes with chain length C_{10}–C_{22} are substances most willingly used by bacteria in metabolic processes [39,46,66,67]. For the most effective bacterial *Mycolicibacterium frederiksbergense* IN53, the content of aliphatic hydrocarbons with carbon chain length nC_{10}–nC_{22} was reduced by 46.9%, and in the case of hydrocarbons with longer chains (nC_{23}–nC_{40}), the degree of their reduction was around 39.5%, which proves a high degree of their biodegradation [39,51].

The tests carried out on sterile Soil C (contaminated with PCBs and TPH) confirmed the previously obtained results and showed that the largest biodegradable potential against polychlorinated biphenyls was characterized by *Rhodococcus* sp. IN306, whereas against petroleum substances the strain *Mycolicibacterium frederiksbergense*. IN53 showed the largest potential. However, when it comes to the joint removal of both types of impurities, the mixed culture M1 proved to be the most effective. Probably in the case of soil polluted with a few (in this case two) xenobiotic types, the application of a bacterial mixed culture developed based on various (in this case three) bacterial strains allows to combine the degradation effect of all strains comprised by the mixed culture parallel to the reduction of competition between strains for the source of food (carbon contained in PCBs and hydrocarbon molecules).

In the case of non-sterile soil, there is always a justified risk that not every allochthonous strain will adapt equally well to the new environment, so there is a risk that the obtained biodegradation effect will be weaker [19,39,51]. In bioremediation techniques for historically PCBs polluted soil, it is preferred to apply mixed bacterial cultures [18,19,38,68]. The studies carried out now on acceleration of petroleum hydrocarbons (TPH) biodegradation through biotechnological processes, with the use of active bacterial cultures, prove that these are the most rational methods for bioremediation of polluted soil and due to relatively low costs and high effectiveness they are practically used on a technical scale [39,40,69].

Therefore, tests on the biodegradation process of PCBs and TPH were carried out in real Soil D (non-sterile), contaminated with petroleum derivatives and aged transformer oil, in ex-situ semi-technical prism method. As a result of the inoculation process with mixed culture M1 of non-sterile Soil D after a period of 6 months, the PCBs content was reduced by 84.5% and TPH by 70.8%.

The calculated first-order biodegradation constants (k) for inoculation with mixed culture M1 for non-sterile Soil D (Table S3) are on a level proving a satisfactory degree of individual PCB congeners biodegradation, decreasing with the increase in the number of chlorine atoms in a molecule [64]. Instead, for petroleum pollutants, they show higher biodegradability of aliphatic hydrocarbons with a carbon chain length nC_{10}–nC_{22} as compared with heavier homologous (nC_{23}–C_{40}) and are close to results of studies presented in the world literature [39,46,67].

To assess the effectiveness of bioremediation, it is preferable to conduct toxicological monitoring using biotests belonging to different trophic groups: Microtox STP [70–73] Ostracodtoxkit [74,75], Phytotoxkit [76–78], MARA environmental risk assessment test [77,79–82] and the Muta-Chromoplate test based on the Ames test [39,83,84]. In this study, unlike other authors, extensive toxicological monitoring was conducted using a set of five biotests, whose bioindicators represented all trophic groups: reducers, decomposers, and producers. It allowed not only the observation of changes in toxicity in Soil D during the inoculation process with the mixed culture M1, but also the determination of the diverse toxicity of the tested organisms to contaminants contained in the tested soil. The calculated toxicity (TU) for individual tests was in decreasing order: Phytotoxkit < Microtox SPT < Ostracodtoxkit < MARA, which shows that the most sensitive bioindicators for impurities contained in the tested Soil D (TPH and PCBs) are *Heterocypris incongruens* and MARA microorganisms. As a result of the conducted bioremediation process, Soil D's toxicity (TU) was reduced from the level of 19.6–10.5 to 5.2–3.1, while the mutagenicity coefficient from 9.85 to 0.68 proves that contaminated soil can be classified as low-toxic (Figures 8 and 9).

The presented results clearly confirmed the rightness of the adopted bioremediation concept of soils contaminated with polychlorinated biphenyls (PCBs) in the presence of total petroleum hydrocarbon (TPH).

4. Materials and Methods

4.1. Soil and Microorganisms

To determine biodegradation effects of each strain, acting independently or within a mixed culture in the absence of native microbiota, three types of sterile soils were used in this studies. Namely, Soil A, commercially available PCB-polluted soil (Certified Reference Material—PCB Congeners in Soil, Product ID: SQC068-50g, Sigma-Aldrich, Laramie, WY, USA–Certificate of analysis Soil A contains in Supplementary Materials); Soil B, petroleum hydrocarbon polluted soil collected from the G-44 waste pit situated in southern-eastern Poland (N 54°14′55″, E 46°68′51″); and Soil C, soil polluted with both PCBs and TPH. Soil B and Soil C were sterilized in an autoclave (SMS ASVE) at 15 psi and 121 °C for 20 min. To determine biodegradation effects of mixed culture M1 in the presence of native microbiota, the ex-situ prism method was applied. A non-sterile Soil D polluted with both PCB and TPH (soil originating from the G-44 waste pit polluted with spent transformer oil containing PCB) was used.

The following bacterial strains—*Mycolicibacterium frederiksbergense* IN53 (formerly *Mycobacterium frederiksbergense* IN53), *Rhodococcus erythropol* is IN129, and *Rhodococcus* sp. IN306 were used in this study. They came from the hydrocarbon-degrading microbial collection of the Department of Microbiology (at the Oil and Gas Institute–National Research Institute, Krakow, Poland) and were isolated from temperate (IN129 and IN53) and desert (IN306) hydrocarbon-exposed soils. In addition, IN53 was previously tested alone or as a member of a hydrocarbon-degrading mixed culture [51]. Diagnostic features of the strain were determined based on microscopic observations, morphology, growth on selective agar media, and biochemical profile (API-Coryne test, bioMerieux).

The strains were phylogenetically identified by sequencing of gene encoding for 16S rRNA as described previously [39,85].

Hydrocarbon-degrading capabilities towards biphenyl and other compounds (n-alkanes, pristane, and aromatic hydrocarbons) were examined using the method of Wrenn and Venosa [86] and by microscopic observations of strain growth on a Kievskaya mineral medium (1 g L^{-1} K_2HPO_4, 1 g L^{-1} KH_2PO_4, 1 g L^{-1} NH_4NO_3, 0.02 g L^{-1} $CaCl_2$, 0.05 g L^{-1} $FeCl_3$, 0.2 g L^{-1} $MgSO_4$, 1 g L^{-1} NaCl, final pH 7.0 ± 0.2) supplemented with biphenyl (1% (*w*/*v*)). After 30-day incubation at room temperature, the strain growth was verified by microscopic observations.

Pure cultures of the individual strains were grown in BD Difco™ nutrient broth (Difco, USA) supplemented with sodium acetate (0.2% (*w*/*v*), POCH, Poland), incubated at room temperature (25 °C) with shaking at 150 rpm for 3–7 days. They reached a density of 3.6 ± 2.4 × 10^8 (IN53), 8.4 ± 3.15 × 10^8 (IN129) and 4.1 ± 4.8 × 10^8 (IN306) colony forming units/mL (CFU·ml^{-1}), respectively, as judged by direct counts on BD Difco™ nutrient agar with sodium acetate (0.2% *w*/*v*). Microbial mixed culture M1 was constructed by mixing equal volumes of each strain (1:1:1) and, therefore, the obtained density was presumable 1 × 10^8–1 × 10^9 CFU·mL^{-1}.

4.2. *Experimental Setup*

4.2.1. Respirometric Tests

The respirometric tests based on the measurement of consumed oxygen (O_2) consumption and/or of released carbon dioxide (CO_2) allow to determine possibilities of aerobic biodegradation of the analyzed soil pollutants. Based on the O_2 consumption and/or CO_2 production rate, it is possible to estimate the rate of aerobic biodegradation of organic pollutant, i.e., the rate of substrate loss over time. Moreover, the results of respirometric tests illustrate the degree of metabolic activity of the polluted environment [87].

Studies on biodegradation of PCBs (Soil A), TPH (Soil B), and both PCBs and TPH (Soil C) in sterile soils were carried out in an Oxi-Top® system (WTW, Bartoszyce, Poland). Twenty grams of each sterile soil were placed in glass sample cells; the system humidity approached the level of 25%. Inoculation was performed using 2 mL of previously isolated bacterial strains IN53, IN129, and IN309 with a density of 1 × 10^8–1 × 10^9 CFU·mL^{-1} and mixed culture M1. At the same time, control samples were prepared. In respirometric tests, control samples were sterile, and uncontaminated "pure" soil samples were inoculated with IN53, IN129, IN306 bacterial stains and mixed culture M1. Control samples were prepared analogously to test samples. After the inoculation, the sample cells were tightly closed with measuring heads, placed in an incubator and thermostated at 20 °C for 30 days. Measuring heads of the Oxi-Top Control system were reading the value of the pressure in the system every 2 h. The acquired data by means of an IR interface was transferred to the Oxi-Top OC 110 controller, where it could be graphically and statistically processed by means of the Achat OC software (WTW, Wroclaw, Poland) [88]. The measured pressure was converted into the amount of used oxygen (MO_2) according to Equation (1) [89]. After the test completion (30 days), analytes were extracted from the studied soils for further analysis by means of gas chromatography. All experiments were performed three times with five repetitions in each treatment.

$$m_{O2} = \frac{M(O_2)}{RT_m} \cdot \left(V_g + \alpha \frac{T_m}{T_0}\right) \cdot \Delta p \quad (1)$$

where: $M(O_2)$—molar mass of oxygen [kg/mol], V_g—free gas volume [m^3], R—gas constant [J mol^{-1} K^{-1}], T_m—measured value of temperature [K], T_0—reference temperature (273.15 K), α—absorption coefficient (0.03103), and Δp—pressure drop in the test [Pa].

4.2.2. PCBs and TPH Biodegradation in Non-sterile Soil-the Ex-Situ Prism Method

Studies on PCBs and TPH biodegradation in non-sterile soil (Soil D) were carried out by means of the ex-situ prism method. The soil at an amount of 50 kg was placed on a specially designed test stand,

ensuring constant temperature (20–25 °C) and humidity (20–25%) during the experiment (Figure S2). A proper choice of biogenic substances and determination of their concentrations in the soil prior to starting the bioremediation is extremely important and allows to enhance the environment with nutrients and to optimize parameters of the process course, where for proper activity of microorganisms, the C:N:P ratio should be approx. 100:10:1 [39,46,67]. The soil pH was corrected, supplementing fertilizer lime at an amount of 1.0–1.5 g/kg of soil till obtaining the optimum pH 7.5–7.6. Prior to the soil inoculation, nitrogen and phosphorus were supplemented in the form of Azofoska mineral fertilizer as follows: 13.6% of total N, 5.5% of nitrate nitrogen, 8.1% of ammonia nitrogen, 6.4% of soluble P_2O_5, 19.1% of K_2O in the form of K_2SO_4, 4.4% of MgO in the form of soluble $MgSO_4$, as well as microelements (0.17% Fe, 0.27% Mn, 0.18% Cu, 0.045% Zn, 0.09% Mo) to obtain optimum proportions of nutrients C:N:P = 100:10:1. Nutrients (nitrogen and phosphorous, present in the soil as ions: NH_4-N (Merck, Spectroquant® cat. no. 114752, Darmstadt, Germany), NO_3-N (Merck, Spectroquant® cat. No. 109713), and PO_4-P (Merck, Spectroquant® cat. No. 114543) were determined by Perkin Elmer Lambda 40 spectrophotometer. Before the inoculation, the total number of heterotrophic aerobic bacteria in non-sterile soil D was $5.9 \pm 5.1 \cdot 10^5$ CFU·g^{-1} dry mass. The soil prepared in this way was inoculated with mixed culture M1 with a density of 10^9 CFU mL^{-1} (2 L mixed culture M1/50 kg soil D). After 6 months of the experiment, the number of bacteria in D6 soil was $4.1 \pm 4.6 \times 10^7$ CFU·g^{-1}. The PCBs and TPH biodegradation was monitored by means of chromatographic analyses (every 30 days during 6 months) and ecotoxicological tests. Control was non-sterile Soil D non-inoculated with mixed culture M1.

4.3. Chromatographic Analysis

4.3.1. PCB Extraction and Quantification

Ten grams of dried soil were placed in an Erlenmeyer flask and extracted using a mixture (80 mL) of n-hexane and dichloromethane (POCH S.A., Gliwice, Poland) at 1:1 ratio [90]. The PCBs were extracted on a Conbest DOS-20L shaker for 3 h, at a constant temperature of 30 °C and shaking rate of 150 rpm [91]. After 3 h, 1 g of powdered copper (Merck, Darmstadt, Germany) was added to the extract and the mixture was extracted for another hour [19]. Copper and interfering substances were removed via filtration using a set of Backerbond columns (silica gel No 7086-03 and PCB-A No 7511-04). The solvent was evaporated in a vacuum rotary evaporator and the extract was dissolved in 1 mL of isooctane (Merck, Germany) and analyzed by the GC method. An internal PCB-209 standard (2,2′,3,3′,4,4′,5,5′,6,6′-decachlorobiphenyl) (Sigma-Aldrich, USA) was used to determine the extraction yield with recovery of approx. 93%.

PCB congeners were analyzed by means of gas chromatography (GC), using a Perkin Elmer Clarus 500 chromatograph with an electron capture detector (ECD). The conditions of GC chromatograph operation were as follows: a capillary column of fused silica (Rtx-XLB 30 m × 0.32 mm ID) (Restek, Bellefonte, PA, USA) with a low-polarity proprietary phase (0.50 µm thick), with the application of the following temperature parameters: injector temperature 280 °C; ECD detector temperature 320 °C; oven temperature programme: 120 °C (2 min isothermal run), 120–280 °C (temperature increase 15 °C min^{-1}), 280 °C (5 min isothermal run), 280–320 °C (temperature increase 28 °C min^{-1}), and 320 °C (10 min isothermal run). The carrier gas was helium (He) at a constant flow of 3.5 mL·min^{-1}. The identification and quantitative analyses of 18 PCB congeners (PCB28 (2,4,4′-), PCB52 (2,2′,5,5′-), PCB77 (3,3′,4,4′-), PCB81 (3,4,4′,5-), PCB101 (2,2′,4,5,5′-), PCB105 (2,3,3′,4,4′-), PCB114 (2,3,4,4′,5-), PCB118 (2,3′,4,4′,5-), PCB123 (2,3′,4,4′,5′-), PCB126 (3,3′,4,4′,5-), PCB138 (2,2′,3,4,4′,5′-), PCB 153 (2,2′,4,4′,5,5′-), PCB156 (2,3,3′,4,4′,5-), PCB157 (2,3,3′,4,4′,5-), PCB167 (2,3′,4,4′,5,5′-), PCB169 (3,3′,4,4′,5,5′-), PCB180 (2,2′,3,4,4′,5,5′-), and PCB189 (2,3,3′,4,4′,5,5′-) were carried out using a certified reference material namely: PCB congeners in the SQC068-50g soil (Sigma-Aldrich, USA). A certified PCB-209 standard (Sigma-Aldrich, Buchs, Switzerland) was used as a biomarker.

4.3.2. TPH Extraction and Quantification

Ten grams of dried soil were placed in an Erlenmeyer flask and extracted using dichloromethane (POCH S.A., Poland) in three series (20 mL of solvent, 15 min). The extraction of petroleum hydrocarbons was carried out by means of the sonification method in an ultrasonic bath Sonoswiss SW 6H, at an ultrasonic frequency of 30 kHz [92]. Polar substances were removed by filtration through Backerbond columns with Florisil No 7213-03 packing. The solvent was evaporated in a vacuum rotary evaporator and the extract was dissolved in 1 mL of dichloromethane and analysed by the GC method. A substitute standard o-tertphenyl was used to determine the extraction yield with recovery of approx. 95.9% [39].

TPH were analysed by means of gas chromatography (GC) using a Perkin Elmer Clarus 500 GC chromatograph with a flame ionisation detector (FID). The conditions of chromatograph operation were as follows: a capillary column of fused silica (RTX-1: 30 m × 0.53 mm) (Restek, USA), applying the following temperature parameters: injector temperature = 290 °C, FID detector temperature = 320 °C, oven temperature programme: 30 °C (2 min. isothermal run), 30–105 °C (temperature increase 10 °C min^{-1}), 105–285 °C (temperature increase 5 °C min^{-1}), and 285 °C (5 min isothermal run). The carrier gas was He at a constant flow of 20 mL·min^{-1}. For the quantitative determination of total petroleum pollutants content (TPH), a set of Tusnovic Instruments calibration standards was used (certified standard: BAM K010). In turn, certified sets of Supelco and Restek standards (standard mixture No D2807 of paraffin hydrocarbons: nC_6–nC_{44} and certified standard mixture No A029668: Fuel Oil Degradation Mix n-C_{17}, pristane, n-C_{18}, phytane) were used to quantitatively determine individual n-alkanes being components of petroleum pollutants. A certified standard C_{30} 17α(H), 21β(H)-hopane No 08189 (Sigma-Aldrich, Switzerland) was used as a biomarker.

4.4. Mathematical Model of PCB and TPH Biodegradation

A simplified mathematical model of polychlorinated biphenyls and petroleum hydrocarbons biodegradation during the performance of tests by the ex-situ method, was developed. PCB-209 biomarker (Sigma-Aldrich, USA) was applied to normalize concentrations of the following analytes: PCBs, TriCB, TetraCBs, PentaCBs, HexaCBs, and HeptaCBs. To normalize concentrations of: TPH, ΣnC_{10}–nC_{22}, and ΣnC_{23}–nC_{40}, a biomarker C_{30}17α(H),21β(H)-hopane (Supelco, Bellefonte, PA, USA) was used. The biomarker application of biomarkers in biodegradation studies allows to eliminate analytical errors in chromatographic determinations of individual analyte groups. Normalized values were used to develop a mathematical first-order model describing biodegradation course, according to Equation (2) [39,64].

$$C/C_x = (C/C_x)_0 \exp(-kt) \quad (2)$$

where: C—analyte concentration, C_x—biomarker (PCB-209 or Hopane) concentration, k—first-order rate constant [d^{-1}], $(C/C_x)_0$—normalized analyte concentration at the starting point, and t—process duration [d].

The analysis of non-linear regression using Equation (2) enabled determination of the first-order biodegradation constant (k) and the correlation coefficient (r^2) defining the measuring points fit to theoretical curves of the biodegradation process: PCBs, TriCB, TetraCBs, PentaCBs, HexaCBs, HeptaCBs, TPH, ΣnC_{11}–nC_{22}, and ΣnC_{23}–nC_{40}.

4.5. Ecotoxicological Analyses

Petroleum hydrocarbon biodegradation leads to the formation of various metabolites often with poorly recognized biological activity. Because of that, it is recommended to carry out toxicological tests, which enable monitoring the changes of soil toxicity. In this study, assessment of toxicity changes during PCB and TPH biodegradation in a non-sterile soil (soil D) inoculated with mixed culture M1 (the ex-situ prism method) was performed. A set of new generation tests, which bioindicators belong to various trophic levels, was used. The specification of toxicological tests used in this studies is presented in Table 2, while the description of their performance procedures is presented in Supplementary Materials.

Table 2. Tests used in monitoring of toxicity alternations in soil under biodegradation.

Trophic Level	Life Form	Test Type	Test Reaction	Reference
Consumers	*Heterocypris incongruens*	Ostracodtoxkit(F)™ (MicroBioTest Inc., Gent, Belgium)	growth inhibition, mortality	[74,75]
Producers	*Sorghum saccharatum* *Sinapis alba* *Lepidium sativum*	Phytotoxkit™ test (MicroBioTests Inc., Belgium),	growth inhibition, germinated seeds	[76–78]
Decomposers	*Vibrio fischeri*	Microtox® Solid Phase Test (SDI, New Castle, DE, USA)	luminescence inhibitionTU = 100/EC_{50}	[70–73]
Decomposers	*Set of 11 bacteria strains*	MARA environmental risk test	growth inhibition	[77,79–82]
		oil contaminants were extracted (SPE) and vaporised, deposit was dissolved in dimethylsulfooxygen (DMSO)		
	Microbacterium sp., *Brevundimonas diminuta, Citrobacter freudii, Comamonas testosterone, Enterococcus casseliflavus, Delftia acidovorans, Kurthia gibsoni, Staphylococcus warnerii, Pseudomonas aurantiaca, Serratia rubidaea, Pichia anomala)*			
Decomposers	*Salmonella typhimurium* (*TA-98, TA-100*)	AMES (Muta-ChromoPlate™ Kit) (EBPI, Mississauga, ON, Canada).	revertants number	[39,83,84]
		oil contaminants were extracted (SPE) and vaporised, deposit was dissolved in DMSO		

4.6. Data Analysis and Statistical Information

Statistical analysis was performed with Statistica 13.3 (StatSoft, Krakow, Poland). Standard deviation (SD), relative standard deviation (RSD) (%) and Pearson correlation coefficient were calculated. Statistically significant differences were evaluated with a one-way ANOVA followed by a post-hoc pairwise Tukey test (when the ANOVA produced significant results). Significance was set at $p < 0.05$.

5. Conclusions

The study was aimed at the assessment of bacterial strains application possibilities: *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306, and mixed culture M1 developed based on these strains. The results of both respirometric tests and chromatographic analyses revealed that chosen bacterial strains with potential PCB decomposition capacity are actually capable of removing this type of pollution from the soil, as well as they are capable of decomposing petroleum hydrocarbons (in particular aliphatic ones). *Rhodococcus* sp. IN306 appeared as the most effective PCB-degrader among the studied strains. Moreover, IN306 is very closely related to *Rhodococcus jostii* RHA-1 (bacterial strains with great PCB-transforming potential). It was shown that the degree of PCB congeners biodegradation was decreasing with increasing number of chlorine atoms and depended on their spatial structure. Petroleum pollutants, in particular n-alkanes, were degraded by *Mycolicibacterium frederiksbergense* IN53 in the most efficient way. Its effect on low-, medium- (nC_{10}–nC_{22}) and long-chain compounds (nC_{23}–nC_{40}) was the greatest among all tested bioaugmentation variants. In turn, mixed culture M1 developed on the basis of tested bacterial strains, featured satisfactory biodegradation capacity both for individual PCB congeners and TPH. The action of mixed culture M1 was verified in a semi-technical scale experiment under ex-situ conditions on a soil polluted with aged petroleum hydrocarbons (TPH) as well as spent transformer oil (PCBs). The effectiveness of this treatment was confirmed by the results of both chromatographic and toxicological tests.

Supplementary Materials: The following are available online, Table S1: Comparison of PCB content in sterile soil samples A and C after inoculation with *Mycolicibacterium frederiksbergense* IN53, *Rhodococcococcus erythropolis* IN129,

Rhodococcus sp. IN306 and mixed culture M1 over 30 days. Measurement repetition number n = 8–10, <0.05., Table S2: Comparison of TPH, unidentified hydrocarons, alkanes (ΣnC10–nC22, ΣnC23–nC40 content in sterile soil samples B and C after by *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306 and mixed culture M1 after the 30 days biodegradation process. Measurement repetition number n = 8–10, <0.05, Table S3: Coefficients of first-order mathematical model describing PCB and TPH pollutants group biodegradation in non-sterile soil D (ex-situ prism method). Measurement repetition number n = 7–10, <0.05, Figure S1: Percentage share of identified pollutants in the sample (Soil D) comprising: (a) PCBs, (b) TPH, Figure S2: Scheme of the test stand for conducting the process of biodegradation of pollutants (PCBs and TPH) in soil D under semi-technical conditions (ex situ prism method), Certificate of Analysis Soil A, Ecotoxicological analyses–description of the methodology.

Author Contributions: Conceptualization, T.S., K.W.; formal analysis, T.S., K.W., P.K.; investigation, T.S., K.W., P.K.; methodology, T.S., K.W., P.K., J.B.; writing—original draft, T.S., K.W. All authors have read and agreed to the published version of the manuscript.

Funding: This research was financially supported by Polish Ministry of Science and Higher Education within statutory funding for Oil and Gas Institute-National Research Institute.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Field, J.A.; Sierra-Alvarez, R. Microbial transformation and degradation of polychlorinated biphenyls. *Environ. Pollut.* **2008**, *155*, 1–12. [CrossRef] [PubMed]
2. Vasilyeva, G.K.; Strijakova, E.R. Bioremediation of soils and sediments contaminated by polychlorinated biphenyls. *Microbiology* **2007**, *76*, 639–653. [CrossRef]
3. Pieper, D.H.; Seeger, M. Bacterial metabolism of polychlorinated biphenyls. *J. Mol. Microbiol. Biotechnol.* **2008**, *15*, 121–138. [CrossRef] [PubMed]
4. Furukawa, K.; Fujihara, H. Microbial degradation of polychlorinated biphenyls: Biochemical and molecular features. *J. Biosci. Bioeng.* **2008**, *105*, 433–449. [CrossRef]
5. Correa, P.A.; Lin, L.; Just, C.L.; Hu, D.; Hornbuckle, K.C.; Schnoor, J.L.; Van Aken, B. The effects of individual PCB congeners on the soil bacterial community structure and the abundance of biphenyl dioxygenase genes. *Environ. Int.* **2010**, *36*, 901–906. [CrossRef] [PubMed]
6. Dudášová, H.; Lukáčová, L.; Murínová, S.; Puškárová, A.; Pangallo, D.; Dercová, K. Bacterial strains isolated from PCB-contaminated sediments and their use for bioaugmentation strategy in microcosms. *J. Basic Microbiol.* **2014**, *54*, 253–260. [CrossRef]
7. Shaikh, N.S.; Parkin, S.; Luthe, G.; Lehmler, H.-J. The three-dimensional structure of 3,3′,4,4′-tetrachlorobiphenyl, a dioxin-like polychlorinated biphenyl (PCB). *Chemosphere* **2008**, *70*, 1694–1698. [CrossRef]
8. Erickson, M.D.; Kaley, R.G. Applications of polychlorinated biphenyls. *Environ. Sci Pollut. Res.* **2011**, *18*, 135–151. [CrossRef]
9. Borja, J.; Taleon, D.M.; Auresenia, J.; Gallardo, S. Polychlorinated biphenyls and their biodegradation. *Process Biochem.* **2005**, *40*, 1999–2013. [CrossRef]
10. Sharma, J.K.; Gautam, R.K.; Nanekar, S.V.; Weber, R.; Singh, B.K.; Singh, S.K.; Juwarkar, A.A. Advances and perspective in bioremediation of polychlorinated biphenyl-contaminated soils. *Environ. Sci Pollut. Res.* **2018**, *25*, 16355–16375. [CrossRef]
11. Blanco-Moreno, R.; Sáez, L.P.; Luque-Almagro, V.M.; Roldán, M.D.; Moreno-Vivián, C. Isolation of bacterial strains able to degrade biphenyl, diphenyl ether and the heat transfer fluid used in thermo-solar plants. *New Biotechnol.* **2017**, *35*, 35–41. [CrossRef] [PubMed]
12. Liang, Y.; Martinez, A.; Hornbuckle, K.C.; Mattes, T.E. Potential for polychlorinated biphenyl biodegradation in sediments from Indiana Harbor and Ship Canal. *Int. Biodeterior. Biodegrad.* **2014**, *89*, 50–57. [CrossRef] [PubMed]
13. Mukerjee-Dhar, G. *Bph* genes of the thermophilic PCB degrader, *Bacillus* sp. JF8: Characterization of the divergent ring-hydroxylating dioxygenase and hydrolase genes upstream of the Mn-dependent BphC. *Microbiology* **2005**, *151*, 4139–4151. [CrossRef] [PubMed]
14. Suyamud, B.; Inthorn, D.; Panyapinyopol, B.; Thiravetyan, P. Biodegradation of bisphenol A by a newly isolated *Bacillus megaterium* strain ISO-2 from a polycarbonate industrial wastewater. *Water Air Soil Pollut.* **2018**, *229*, 348. [CrossRef]
15. Ponce, B.L.; Latorre, V.K.; González, M.; Seeger, M. Antioxidant compounds improved PCB-degradation by *Burkholderia xenovorans* strain LB400. *Enzym. Microb. Technol.* **2011**, *49*, 509–516. [CrossRef] [PubMed]

16. Sierra, I.; Valera, J.L.; Marina, M.L.; Laborda, F. Study of the biodegradation process of polychlorinated biphenyls in liquid medium and soil by a new isolated aerobic bacterium (*Janibacter* sp.). *Chemosphere* **2003**, *53*, 609–618. [CrossRef]
17. Moody, J.; Doerge, D.; Freeman, J.; Cerniglia, C. Degradation of biphenyl by *Mycobacterium* sp. strain PYR-1. *Appl. Microbiol. Biotechnol.* **2002**, *58*, 364–369. [CrossRef]
18. Murínová, S.; Dercová, K. Potential Use of Newly Isolated Bacterial Strain *Ochrobactrum anthropi* in bioremediation of polychlorinated biphenyls. *Water Air Soil Pollut.* **2014**, *225*, 1980. [CrossRef]
19. Horváthová, H.; Lászlová, K.; Dercová, K. Bioremediation of PCB-contaminated shallow river sediments: The efficacy of biodegradation using individual bacterial strains and their consortia. *Chemosphere* **2018**, *193*, 270–277. [CrossRef]
20. Zorádová, S.; Dudášová, H.; Lukáčová, L.; Dercová, K.; Čertík, M. The effect of polychlorinated biphenyls (PCBs) on the membrane lipids of Pseudomonas stutzeri. *Int. Biodeterior. Biodegrad.* **2011**, *65*, 1019–1023. [CrossRef]
21. Tao, B.; Zhou, Y.; He, X.; Li, D. *Pseudomonas chengduensis* sp. nov., isolated from landfill leachate. *Int. J. Syst. Evol. Microbiol.* **2014**, *64*, 95–100. [CrossRef] [PubMed]
22. Ridl, J.; Suman, J.; Fraraccio, S.; Hradilova, M.; Strejcek, M.; Cajthaml, T.; Zubrova, A.; Macek, T.; Strnad, H.; Uhlik, O. Complete genome sequence of *Pseudomonas alcaliphila* JAB1 (=DSM 26533), a versatile degrader of organic pollutants. *Stand Genom. Sci.* **2018**, *13*, 3. [CrossRef]
23. Chakraborty, J.; Das, S. Characterization of the metabolic pathway and catabolic gene expression in biphenyl degrading marine bacterium *Pseudomonas aeruginosa* JP-11. *Chemosphere* **2016**, *144*, 1706–1714. [CrossRef] [PubMed]
24. Suenaga, H.; Fujihara, H.; Kimura, N.; Hirose, J.; Watanabe, T.; Futagami, T.; Goto, M.; Shimodaira, J.; Furukawa, K. Insights into the genomic plasticity of *Pseudomonas putida* KF715, a strain with unique biphenyl-utilizing activity and genome instability properties: Genomic plasticity of *Pseudomonas putida* KF715. *Environ. Microbiol. Rep.* **2017**, *9*, 589–598. [CrossRef] [PubMed]
25. Weiland-Bräuer, N.; Fischer, M.A.; Schramm, K.-W.; Schmitz, R.A. Polychlorinated biphenyl (PCB)-degrading potential of microbes present in a cryoconite of Jamtalferner Glacier. *Front. Microbiol.* **2017**, *8*, 1105. [CrossRef]
26. Chung, S.-Y.; Maeda, M.; Song, E.; Horikoshij, K.; Kudo, T. A Gram-positive polychlorinated biphenyl-degrading bacterium, *Rhodococcus erythropolis* strain TA421, isolated from a termite ecosystem. *Biosci. Biotechnol. Biochem.* **1994**, *58*, 2111–2113. [CrossRef]
27. Taguchi, K.; Motoyama, M.; Kudo, T. Multiplicity of 2,3-dihydroxybiphenyl dioxygenase genes in the gram-positive polychlorinated biphenyl degrading bacterium *Rhodococcus rhodochrous* K37. *Biosci. Biotechnol. Biochem.* **2004**, *68*, 787–795. [CrossRef] [PubMed]
28. Murínová, S.; Dercová, K.; Dudášová, H. Degradation of polychlorinated biphenyls (PCBs) by four bacterial isolates obtained from the PCB-contaminated soil and PCB-contaminated sediment. *Int. Biodeterior. Biodegrad.* **2014**, *91*, 52–59. [CrossRef]
29. Pham, T.T.M.; Pino Rodriguez, N.J.; Hijri, M.; Sylvestre, M. Optimizing polychlorinated biphenyl degradation by flavonoid-induced cells of the *Rhizobacterium Rhodococcus erythropolis* U23A. *PLoS ONE* **2015**, *10*, e0126033. [CrossRef]
30. Shumkova, E.S.; Olsson, B.E.; Kudryavtseva, A.V.; Plotnikova, E.G. Draft genome sequence of *Rhodococcus ruber* strain P25, an active polychlorinated biphenyl degrader. *Genome Announc.* **2015**, *3*. [CrossRef]
31. Atago, Y.; Shimodaira, J.; Araki, N.; Bin Othman, N.; Zakaria, Z.; Fukuda, M.; Futami, J.; Hara, H. Identification of novel extracellular protein for PCB/biphenyl metabolism in *Rhodococcus jostii* RHA1. *Biosci. Biotechnol. Biochem.* **2016**, *80*, 1012–1019. [CrossRef] [PubMed]
32. Xu, Y.; Yu, M.; Shen, A. Complete genome sequence of the polychlorinated biphenyl degrader *Rhodococcus* sp. WB1. *Genome Announc.* **2016**, *4*. [CrossRef] [PubMed]
33. Wang, H.; Hu, J.; Xu, K.; Tang, X.; Xu, X.; Shen, C. Biodegradation and chemotaxis of polychlorinated biphenyls, biphenyls, and their metabolites by *Rhodococcus* sp. *Biodegradation* **2018**, *29*, 1–10. [CrossRef] [PubMed]
34. Sato, T.; Todoroki, T.; Shimoda, K.; Terada, A.; Hosomi, M. Behavior of PCDDs/PCDFs in remediation of PCBs-contaminated sediments by thermal desorption. *Chemosphere* **2010**, *80*, 184–189. [CrossRef] [PubMed]
35. Wang, X.; Teng, Y.; Luo, Y.; Dick, R.P. Biodegradation of 3,3′,4,4′-tetrachlorobiphenyl by *Sinorhizobium meliloti* NM. *Bioresour. Technol.* **2016**, *201*, 261–268. [CrossRef]

36. Jing, R.; Fusi, S.; Kjellerup, B.V. Remediation of polychlorinated biphenyls (PCBs) in contaminated soils and sediment: State of knowledge and perspectives. *Front. Environ. Sci.* **2018**, *6*, 1–17. [CrossRef]
37. Pathiraja, G.; Egodawatta, P.; Goonetilleke, A.; Teo, V.S.J. Effective degradation of polychlorinated biphenyls by a facultative anaerobic bacterial consortium using alternating anaerobic aerobic treatments. *Sci. Total Environ.* **2019**, *659*, 507–514. [CrossRef]
38. Cervantes-González, E.; Cuevara-García, M.A.; García-Mena, J.; Ovando-Medina, V.M. Microbial diversity assessment of polychlorinated biphenyl–contaminated soils and the biostimulation and bioaugmentation processes. *Environ. Monit. Assess* **2019**, *191*, 118. [CrossRef]
39. Steliga, T.; Jakubowicz, P.; Kapusta, P. Changes in toxicity during in situ bioremediation of weathered drill wastes contaminated with petroleum hydrocarbons. *Bioresour. Technol.* **2012**, *125*, 1–10. [CrossRef]
40. Silva-Castro, G.A.; Rodriguez-Calvo, A.; Laguna, J.; González-López, J.; Calvo, C. Autochthonous microbial responses and hydrocarbons degradation in polluted soil during biostimulating treatments under different soil moisture. Assay in pilot plant. *Int. Biodeterior. Biodegrad.* **2016**, *108*, 91–98. [CrossRef]
41. Roy, A.; Dutta, A.; Pal, S.; Gupta, A.; Sarkar, J.; Chatterjee, A.; Saha, A.; Sarkar, P.; Sar, P.; Kazy, S.K. Biostimulation and bioaugmentation of native microbial community accelerated bioremediation of oil refinery sludge. *Bioresour. Technol.* **2018**, *253*, 22–32. [CrossRef] [PubMed]
42. Sarkar, P.; Roy, A.; Pal, S.; Mohapatra, B.; Kazy, S.K.; Maiti, M.K.; Sar, P. Enrichment and characterization of hydrocarbon-degrading bacteria from petroleum refinery waste as potent bioaugmentation agent for in situ bioremediation. *Bioresour. Technol.* **2017**, *242*, 15–27. [CrossRef] [PubMed]
43. Tyagi, M.; da Fonseca, M.M.R.; de Carvalho, C.C.C.R. Bioaugmentation and biostimulation strategies to improve the effectiveness of bioremediation processes. *Biodegradation* **2011**, *22*, 231–241. [CrossRef] [PubMed]
44. Havel, J.; Reineke, W. Degradation of Aroclor 1221 and survival of strains in soil microcosms. *Appl Microbiol. Biotechnol.* **1992**, *38*, 129–134. [CrossRef]
45. Egorova, D.O.; Demakov, V.A.; Plotnikova, E.G. Bioaugmentation of a polychlorobiphenyl contaminated soil with two aerobic bacterial strains. *J. Hazard. Mater.* **2013**, *261*, 378–386. [CrossRef]
46. Wu, M.; Li, W.; Dick, W.A.; Ye, X.; Chen, K.; Kost, D.; Chen, L. Bioremediation of hydrocarbon degradation in a petroleum-contaminated soil and microbial population and activity determination. *Chemosphere* **2017**, *169*, 124–130. [CrossRef]
47. Lászlová, K.; Dudášová, H.; Olejníková, P.; Horváthová, G.; Velická, Z.; Horváthová, H.; Dercová, K. The application of biosurfactants in bioremediation of the aged sediment contaminated with polychlorinated biphenyls. *Water Air Soil Pollut.* **2018**, *229*, 219. [CrossRef]
48. Passatore, L.; Rossetti, S.; Juwarkar, A.A.; Massacci, A. Phytoremediation and bioremediation of polychlorinated biphenyls (PCBs): State of knowledge and research perspectives. *J. Hazard. Mater.* **2014**, *278*, 189–202. [CrossRef]
49. Steliga, T.; Kapusta, P.; Jakubowicz, P. Effectiveness of bioremediation processes of hydrocarbon pollutants in weathered drill wastes. *Water Air Soil Pollut.* **2009**, *202*, 211–228. [CrossRef]
50. Steliga, T.; Jakubowicz, P.; Kapusta, P.; Kluk, D. Badania biodegradacji odpadów wiertniczych zanieczyszczonych substancjami ropopochodnymi (ang. Study on biodegradation of drilling wastes contaminated with petroleum hydrocarbons). *Przemysł Chem.* **2018**, *97*, 1666–1675. [CrossRef]
51. Brzeszcz, J.; Steliga, T.; Kapusta, P.; Turkiewicz, A.; Kaszycki, P. r-strategist versus K-strategist for the application in bioremediation of hydrocarbon-contaminated soils. *Int. Biodeterior. Biodegrad.* **2016**, *106*, 41–52. [CrossRef]
52. Pruesse, E.; Peplies, J.; Glöckner, F.O. SINA: Accurate high-throughput multiple sequence alignment of ribosomal RNA genes. *Bioinformatics* **2012**, *28*, 1823–1829. [CrossRef] [PubMed]
53. Ludwig, W.; Strunk, O.; Westram, R.; Richter, L.; Meier, H.; Yadhukumar; Buchner, A.; Lai, T.; Steppi, S.; Jobb, G.; et al. ARB: A software environment for sequence data. *Nucleic Acids Res.* **2004**, *32*, 1363–1371. [CrossRef]
54. Elangovan, S.; Pandian, S.B.S.; Geetha, S.J.; Joshi, S.J. Polychlorinated biphenyls (PCBs): Environmental fate, challenges and bioremediation. In *Microbial Metabolism of Xenobiotic Compounds*; Arora, P.K., Ed.; Springer: Singapore, 2019; pp. 165–188. [CrossRef]
55. Tu, C.; Teng, Y.; Luo, Y.; Li, X.; Sun, X.; Li, Z.; Liu, W.; Christie, P. Potential for biodegradation of polychlorinated biphenyls (PCBs) by *Sinorhizobium meliloti*. *J. Hazard. Mater.* **2011**, *186*, 1438–1444. [CrossRef]

56. Horváthová, H.; Lászlová, K.; Dercová, K. Bioremediation vs. nanoremediation: Degradation of polychlorinated biphenyls (PCBS) using integrated remediation approaches. *Water Air Soil Pollut.* **2019**, *230*, 204. [CrossRef]
57. Nabavi, B.; Nikaeen, M.; Amin, M.; Hatamzadeh, M. Isolation and identification of aerobic polychlorinated biphenyls degrading bacteria. *Int. J. Environ. Health Eng.* **2013**, *2*, 47. [CrossRef]
58. McLeod, M.P.; Warren, R.L.; Hsiao, W.W.L.; Araki, N.; Myhre, M.; Fernandes, C.; Miyazawa, D.; Wong, W.; Lillquist, A.L.; Wang, D.; et al. The complete genome of *Rhodococcus* sp. RHA1 provides insights into a catabolic powerhouse. *Proc. Natl. Acad. Sci. USA* **2006**, *103*, 15582–15587. [CrossRef]
59. Rodrigues, J.L.M.; Kachel, C.A.; Aiello, M.R.; Quensen, J.F.; Maltseva, O.V.; Tsoi, T.V.; Tiedje, J.M. Degradation of Aroclor 1242 dechlorination products in sediments by *Burkholderia xenovorans* LB400(*ohb*) and *Rhodococcus* sp. Strain RHA1(*fcb*). *Appl. Environ. Microbiol.* **2006**, *72*, 2476–2482. [CrossRef] [PubMed]
60. Ohmori, T.; Morita, H.; Tanaka, M.; Miyauchi, K.; Kasai, D.; Furukawa, K.; Miyashita, K.; Ogawa, N.; Masai, E.; Fukuda, M. Development of a strain for efficient degradation of polychlorinated biphenyls by patchwork assembly of degradation pathways. *J. Biosci. Bioeng.* **2011**, *111*, 437–442. [CrossRef] [PubMed]
61. McKay, D.B.; Prucha, M.; Reineke, W.; Timmis, K.N.; Pieper, D.H. Substrate specificity and expression of three 2,3-dihydroxybiphenyl 1,2-dioxygenases from *Rhodococcus globerulus* Strain P6. *J. Bacteriol.* **2003**, *185*, 2944–2951. [CrossRef] [PubMed]
62. Yang, X.; Sun, Y.; Qian, S. Biodegradation of seven polychlorinated biphenyls by a newly isolated aerobic bacterium (*Rhodococcus* sp. R04). *J. Ind. Microbiol. Biotechnol.* **2004**, *31*, 415–420. [CrossRef] [PubMed]
63. Hu, C.; Zhang, Y.; Tang, X.; Luo, W. PCB biodegradation and *bphA1* gene expression induced by salicylic acid and biphenyl with *Pseudomonas fluorescence* P2W and *Ralstonia eutropha* H850. *Pol. J. Environ. Stud.* **2014**, *23*, 1591–1598.
64. Zhang, H.; Jiang, X.; Lu, L.; Xiao, W. Biodegradation of polychlorinated biphenyls (PCBs) by the novel identified *cyanobacterium Anabaena* PD-1. *PLoS ONE* **2015**, *10*, e0131450. [CrossRef]
65. Furukawa, K. Molecular genetics and evolutionary relationship of PCB-degrading bacteria. *Biodegradation* **1994**, *5*, 289–300. [CrossRef]
66. Abbasian, F.; Lockington, R.; Megharaj, M.; Naidu, R. A review on the genetics of aliphatic and aromatic hydrocarbon degradation. *Appl. Biochem. Biotechnol.* **2016**, *178*, 224–250. [CrossRef] [PubMed]
67. Nwankwegu, A.S.; Orji, M.U.; Onwosi, C.O. Studies on organic and in-organic biostimulants in bioremediation of diesel-contaminated arable soil. *Chemosphere* **2016**, *162*, 148–156. [CrossRef] [PubMed]
68. Dercová, K.; Čičmanová, J.; Lovecká, P.; Demnerová, K.; Macková, M.; Hucko, P.; Kušnír, P. Isolation and identification of PCB-degrading microorganisms from contaminated sediments. *Int. Biodeterior. Biodegrad.* **2008**, *62*, 219–225. [CrossRef]
69. Guarino, C.; Zuzolo, D.; Marziano, M.; Conte, B.; Baiamonte, G.; Morra, L.; Benotti, D.; Gresia, D.; Stacul, E.R.; Cicchella, D.; et al. Investigation and assessment for an effective approach to the reclamation of polycyclic aromatic hydrocarbon (PAHs) contaminated site: SIN Bagnoli, Italy. *Sci. Rep.* **2019**, *9*, 11522. [CrossRef]
70. Adams, R.H.; Kanga-Leyva, K.; Guzmán-Osorio, F.J.; Escalante- Espinosa, E. Comparison of moisture management methods for the bioremediation of hydrocarbon contaminated soil. *Afr. J. Biotechnol.* **2011**, *19*, 394–404. [CrossRef]
71. Foucault, Y.; Durand, M.-J.; Tack, K.; Schreck, E.; Geret, F.; Leveque, T.; Pradere, P.; Goix, S.; Dumat, C. Use of ecotoxicity test and ecoscores to improve the management of polluted soils: Case of a secondary lead smelter plant. *J. Hazard. Mater.* **2013**, *246–247*, 291–299. [CrossRef]
72. Lima, T.M.S.; Procópio, L.C.; Brandão, F.D.; Leão, B.A.; Tótola, M.R.; Borges, A.C. Evaluation of bacterial surfactant toxicity towards petroleum degrading microorganisms. *Bioresour. Technol.* **2011**, *102*, 2957–2964. [CrossRef] [PubMed]
73. Oleszczuk, P.; Jośko, I.; Kuśmierz, M.; Futa, B.; Wielgosz, E.; Ligęza, S.; Pranagal, J. Microbiological, biochemical and ecotoxicological evaluation of soils in the area of biochar production in relation to polycyclic aromatic hydrocarbon content. *Geoderma* **2014**, *213*, 502–511. [CrossRef]
74. Niyommaneerat, W.; Nakajima, F.; Tobino, T.; Yamamoto, K. Development of a chronic sediment toxicity test using the benthic ostracod *Heterocypris incongruens* and their application to toxicity assessments of urban road dust. *Ecotoxicol. Environ. Saf.* **2017**, *143*, 266–274. [CrossRef] [PubMed]
75. Steliga, T. The use of biotests in estimation of bioremediation processes in weathered drilling wastes. *Arch. Environ. Prot.* **2011**, *2*, 61–70.

76. Baran, A.; Tarnawski, M. Phytotoxkit/Phytotestkit and Microtox® as tools for toxicity assessment of sediments. *Ecotoxicol. Environ. Saf.* **2013**, *98*, 19–27. [CrossRef]
77. Fai, P.B.; Grant, A. An assessment of the potential of the microbial assay for risk assessment (MARA) for ecotoxicological testing. *Ecotoxicology* **2010**, *19*, 1626–1633. [CrossRef]
78. Mamindy-Pajany, Y.; Hamer, B.; Roméo, M.; Géret, F.; Galgani, F.; Durmiši, E.; Hurel, C.; Marmier, N. The toxicity of composted sediments from Mediterranean ports evaluated by several bioassays. *Chemosphere* **2011**, *82*, 362–369. [CrossRef]
79. Gabrielson, J.; Kühn, I.; Colque-Navarro, P.; Hart, M.; Iversen, A.; McKenzie, D.; Möllby, R. Microplate-based microbial assay for risk assessment and (eco)toxic fingerprinting of chemicals. *Anal. Chim. Acta* **2003**, *485*, 121–130. [CrossRef]
80. Gabrielson, J. *Assessing the Toxic Impact of Chemicals Using Bacteria*; Microbiology and Tumöbiologist Centrum (MTC)/Microbiology and Tumor Biology Center (MTC); Karolinska University Press: Stockholm, Sweden, 2004; ISBN 978917140134.
81. Paskuliakova, A.; McGowan, T.; Tonry, S.; Touzet, N. Phycoremediation of landfill leachate with the chlorophyte *Chlamydomonas* sp. SW15aRL and evaluation of toxicity pre and post treatment. *Ecotoxicol. Environ. Saf.* **2018**, *147*, 622–630. [CrossRef]
82. Wadhia, K.; Dando, T.; Clive Thompson, K. Intra-laboratory evaluation of Microbial Assay for Risk Assessment (MARA) for potential application in the implementation of the Water Framework Directive (WFD). *J. Environ. Monit.* **2007**, *9*, 953. [CrossRef]
83. Kamber, M.; Fluckiger-Isler, S.; Engelhardt, G.; Jaeckh, R.; Zeiger, E. Comparison of the Ames II and traditional Ames test responses with respect to mutagenicity, strain specificities, need for metabolism and correlation with rodent carcinogenicity. *Mutagenesis* **2009**, *24*, 359–366. [CrossRef] [PubMed]
84. Vijay, U.; Gupta, S.; Mathur, P.; Suravajhala, P.; Bhatnagar, P. Microbial mutagenicity assay: Ames Test. *Biol. Protoc.* **2018**, *8*, 32763. [CrossRef]
85. Wang, D.; Lin, J.; Lin, J.; Wang, W.; Li, S. Biodegradation of petroleum hydrocarbons by *Bacillus subtilis* BL-27, a strain with weak hydrophobicity. *Molecules* **2019**, *24*, 3021. [CrossRef] [PubMed]
86. Wrenn, B.A.; Venosa, A.D. Selective enumeration of aromatic and aliphatic hydrocarbon degrading bacteria by a most-probable-number procedure. *Can. J. Microbiol.* **1996**, *42*, 252–258. [CrossRef]
87. Malińska, K. Application of a modified OxiTop® respirometer for laboratory composting studies. *Arch. Environ. Prot.* **2016**, *42*, 56–62. [CrossRef]
88. Steliga, T.; Wojtowicz, K. Wykorzystanie testów respirometrycznych do oceny efektywności biodegradacji osadów z instalacji kopalnianych. *Naft. Gaz* **2019**, *75*, 29–37. [CrossRef]
89. Steliga, T.; Uliasz, M. Spent drilling muds management and natural environment protection. *Gospod. Surowcami Min.* **2014**, *30*, 135–155. [CrossRef]
90. Tian, L.; Huang, D.; Shi, Y.; Han, F.; Wang, Y.; Ye, H.; Tang, Y.; Yu, H. Method for the analysis of 7 indictor polychlorinated biphenyls (PCBs) and 13 organochlorine pesticide residues in sediment by gas chromatography (GC). *IOP Conf. Ser. Earth Environ. Sci.* **2019**, *237*, 022053. [CrossRef]
91. Wojtowicz, K.; Jakubowicz, P. Opracowanie metodyki oznaczania polichlorowanych bifenyli w próbkach gleb. *Naft. Gaz* **2019**, *75*, 420–429. [CrossRef]
92. Chaîneau, C.H.; Yepremian, C.; Vidalie, J.F.; Ducreux, J.; Ballerini, D. Bioremediation of a crude oil-polluted soil: Biodegradation, leaching and toxicity assessments. *Water Air Soil Pollut.* **2003**, *144*, 419–440. [CrossRef]

Sample Availability: Samples of the compounds are not available from the authors.

© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (http://creativecommons.org/licenses/by/4.0/).

molecules

MDPI

Suplementary materials

Assessment of biodegradation efficiency of polychlorinated biphenyls (PCBs) and petroleum hydrocarbons (TPH) in soil using three individual bacterial strains and their mixed culture

Teresa Steliga[1], Katarzyna Wojtowicz[1], Piotr Kapusta[2] and Joanna Brzeszcz[2]

[1] Department of Production Technology of Reservoir Fluids, Oil and Gas Institute – National Research Institute, 31-503 Krakow, ul. Lubicz 25 A, Poland

[2] Department of Microbiology, Oil and Gas Institute-National Research Institute, 31-503 Krakow, ul. Lubicz 25A, Poland

* Correspondence: teresa.steliga@inig.pl

Table S1. Comparison of PCB content in sterile soil samples A and C after inoculation with *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306 and mixed culture M1 over 30 days. Measurement repetition number n=8–10, <0.05.

	Soil A (PCB µg/kg s.m. contaminated sterile soil)				
	Soil control	IN53	IN129	IN306	M1
PCBs	13,100 ± 915	8,391 ± 421	7,391 ± 402	5,971 ± 354	7,050 ± 371
TriCB	494 ± 28.5	202 ±15.9	162 ±11.5	123 ±9.1	134 ±10.3
TetraCBs	2,026 ± 94.5	962 ± 44.4	820 ± 41.5	678 ± 39.4	740 ± 41.2
PentaCs	4,838 ± 185.1	2,862 ± 112.2	2,814 ± 127.9	2,172 ± 99.6	2,618 ± 104.8
HexaCBs	3,873 ± 121.2	2,967 ± 105.1	2,392 ± 92.1	1,962 ± 71.6	2,349 ±88.2
HeptCBs	1,680 ± 82.1	1,398 ± 71.2	1,202 ± 65.9	1,034 ± 51.6	1,109 ± 64.9
	Soil C (PCB µg/kg s.m. contaminated sterile soil)				
	Soil control	IN53	IN129	IN306	M1
PCBs	13,100 ± 915	9,243 ± 358	8,001 ± 454	6,853 ± 387	6,309 ± 365
TriCB	494 ±28.5	241 ± 19.5	192 ±15.8	151 ±11.9	146 ±10.9
TetraCBs	2,026 ± 94.5	1,168 ± 55.3	934 ± 45.4	764 ± 38.9	761 ± 38.5
PentaCs	4,838 ± 185.1	3,250 ± 132.8	3,055 ± 145.6	2,525 ± 93.3	2,277 ±99.2
HexaCBs	3,873 ± 121.2	3,137 ± 111.4	2,562 ± 89.5	2,233 ± 84.1	2,060 ± 82.6
HeptCBs	1,680 ± 82.1	1,449 ± 88.2	1,256 ± 51.3	1,184 ± 53.5	1,059 ± 51.9

molecules

MDPI

Table S2. Comparison of TPH, unidentified hydrocarons, alkanes (ΣnC_{10}–nC_{22}, ΣnC_{23} – nC_{40} content in sterile soil samples B and C after by *Mycolicibacterium frederiksbergense* IN53, *Rhodococcus erythropolis* IN129, *Rhodococcus* sp. IN306 and mixed culture M1 after the 30 day biodegradation process. Measurement repetition number n=8–10, <0.05.

	Soil B (TPH mg/kg s.m. contaminated sterile soil)				
	Soil control	IN53	IN129	IN306	M1
TPH	12,515 ± 787	7,863 ± 398	8,951 ± 451	9,396 ± 487	8.309 ± 402
Unident. hydrocarb.	5,687±258	3,825± 198	4,115 ± 214	4,223 ± 247	3,990 ± 214
ratio nC_{17}/pristane	0.818 ± 0.04	0.473 ± 0.02	0.536 ± 0.02	0.594 ± 0.02	0.522 ± 0.02
ratio nC_{18}/phytane	1.657 ± 0.09	0.985 ± 0.07	1.154 ± 0.08	1.232 ±0.08	1.029 ± 0.08
Σ nC_{10} – nC_{22}	4,058 ± 187	2,154 ± 135	2.589 ±128	2.788 ±112	2,292 ± 108
Σ nC_{23} – nC_{40}	1,956 ± 178	1,183 ± 99	1,546 ± 139	1,658 ± 147	1,334± 129
	Soil C (TPH mg/kg s.m. contaminated sterile soil)				
	Soil control	IN53	IN129	IN306	M1
TPH	12,365 ±769	8,282 ± 411	9,426 ± 499	9,838 ± 529	8,085 ±396
Unident. hydrocarb.	5,536 ± 247	3,909 ± 196	4,345 ± 247	4,541 ± 274	3,874 ± 228
ratio nC_{17}/pristane	0.819 ± 0.04	0.541± 0.02	0.489 ± 0.02	0.601 ± 0.02	0.495 ± 0.02
Ratio nC_{18}/phytane	1.657 ±0.09	1.167 ± 0.08	0.996 ± 0.07	1.266 ± 0.08	1.009 ± 0.07
Σ nC_{10}-nC_{22}	4,076 ± 288	2,336 ± 145	2,732 ± 165	2,876 ± 128	2,235 ±114
Σ nC_{23}-nC_{40}	1,956 ± 198	1,321 ± 117	1,614 ± 142	1,686 ± 145	1,264 ± 109

Table S3. Coefficients of first-order mathematical model describing PCB and TPH pollutants group biodegradation in non-sterile soil D (ex-situ prism method). Measurement repetition number n=7–10, <0.05.

PCB pollutants group	k [day^{-1}]	$(C/C_{PCB209})_0$	Determination coefficient (r^2)
PCBs	0.0109 ± 0.0009	16,102 ± 80.5	0.9928
TriCB	0.0170 ± 0.0012	240.6 ± 12.0	0.9883
TetraCBs	0.0110 ± 0.0008	545.8 ± 27.3	0.9897
PentaCBs	0.0095 ± 0.0007	593.1 ± 29.7	0.9908
HexaCBs	0.0082 ± 0.0006	216.1 ± 10.8	0.9853
HeptCB	0.0050 ± 0.0003	21.0 ± 1.1	0.9741
Petroleum pollutants group	k [day^{-1}]	$(C/C_H)_0$	Determination coefficient (r^2)
TPH	0.0070 ± 0.0004	4,087.1 ± 204.4	0.9841
Σ nC_{11} – nC_{22}	0.0130 ± 0.0009	1,892.2 ± 94.6	09941
Σ nC_{23} – nC_{40}	0.0049 ± 0.0003	867.5 ± 43.4	0.9976

molecules




Figure S1. Percentage share of identified pollutants in the sample (Soil D) comprising: (a) PCBs, (b) TPH




Figure S2. Scheme of the test stand for conducting the process of biodegradation of pollutants (PCBs and TPH) in soil D under semi-technical conditions (*ex situ* prism method)

molecules

MDPI

Certificate of Analysis

Certified Reference Material

PCB Congeners in Soil

Product ID SQC068-50G
Lot LRAB537Z
Expiration Date December 31, 2021
Storage Conditions Store at room temperature

Analyte	Units	Certified [1,4] Value	k [5]	Standard [2] Deviation	Acceptance [3] Interval
PCBs, total	ug/Kg	13,100 ± 459	2.00	1,965	7,205 to 18,995
2,4,4'-Trichlorobiphenyl (PCB 28) Trichlorobiphenyl, 2,4,4'-	ug/Kg	494 ± 17.3	2.00	74.1	272 to 716
2,2',5,5'-Tetrachlorobiphenyl (PCB 52) Tetrachlorobiphenyl, 2,2',5,5'-	ug/Kg	607 ± 21.2	2.00	91.1	334 to 880
3,3',4,4'-Tetrachlorobiphenyl (PCB 77) Tetrachlorobiphenyl, 3,3',4,4'-	ug/Kg	635 ± 22.2	2.00	95.3	349 to 921
3,4,4',5-Tetrachlorobiphenyl (PCB 81) Tetrachlorobiphenyl, 3,4,4',5-	ug/Kg	784 ± 27.4	2.00	118	431 to 1,137
2,2',4,5,5'-Pentachlorobiphenyl (PCB 101) Pentachlorobiphenyl, 2,2',4,5,5'-	ug/Kg	803 ± 28.1	2.00	120	442 to 1,164
2,3,3',4,4'-Pentachlorobiphenyl (PCB 105) Pentachlorobiphenyl, 2,3,3',4,4'-	ug/Kg	733 ± 25.7	2.00	110	403 to 1,063
2,3',4,4',5-Pentachlorobiphenyl (PCB 118) Pentachlorobiphenyl, 2,3',4,4',5-	ug/Kg	548 ± 19.2	2.00	82.2	301 to 795
2,3',4,4',5'-Pentachlorobiphenyl (PCB 123) Pentachlorobiphenyl, 2,3',4,4',5'-	ug/Kg	917 ± 32.1	2.00	138	504 to 1,330
2,3,4,4',5-Pentachlorobiphenyl (PCB 114) Pentachlorobiphenyl, 2,3,4,4',5-	ug/Kg	577 ± 20.2	2.00	86.6	317 to 837
3,3',4,4',5-Pentachlorobiphenyl (PCB 126) Pentachlorobiphenyl, 3,3',4,4',5-	ug/Kg	1,260 ± 44.1	2.00	189	693 to 1,827
2,2',3,4,4',5'-Hexachlorobiphenyl (PCB 138) Hexachlorobiphenyl, 2,2',3,4,4',5'-	ug/Kg	608 ± 21.3	2.00	91.2	334 to 882
2,2',4,4',5,5'-Hexachlorobiphenyl (PCB 153) Hexachlorobiphenyl, 2,2',4,4',5,5'-	ug/Kg	571 ± 20.0	2.00	85.7	314 to 828
2,3,3',4,4',5'-Hexachlorobiphenyl (PCB 157) Hexachlorobiphenyl, 2,3,3',4,4',5'-	ug/Kg	535 ± 18.7	2.00	80.3	294 to 776
2,3,3',4,4',5-Hexachlorobiphenyl (PCB 156) Hexachlorobiphenyl, 2,3,3',4,4',5-	ug/Kg	757 ± 26.5	2.00	114	416 to 1,098
2,3',4,4',5,5'-Hexachlorobiphenyl (PCB 167) Hexachlorobiphenyl, 2,3',4,4',5,5'-	ug/Kg	635 ± 22.2	2.00	95.3	349 to 921
3,3',4,4',5,5'-Hexachlorobiphenyl (PCB 169) Hexachlorobiphenyl, 3,3',4,4',5,5'-	ug/Kg	767 ± 26.8	2.00	115	422 to 1,112
2,2',3,4,4',5,5'-Heptachlorobiphenyl (PCB 180) Heptachlorobiphenyl, 2,2',3,4,4',5,5'-	ug/Kg	1,160 ± 40.6	2.00	174	638 to 1,682
2,3,3',4,4',5,5'-Heptachlorobiphenyl (PCB 189) Heptachlorobiphenyl, 2,3,3',4,4',5,5'-	ug/Kg	520 ± 18.2	2.00	78.0	286 to 754
PCB (156)+(157)	ug/Kg	1,290 ± 45.2	2.00	194	710 to 1,871
PCB (20)+(28)	ug/Kg	494 ± 17.3	2.00	74.1	272 to 716
PCB (90)+(101)+(113)	ug/Kg	803 ± 28.1	2.00	120	442 to 1,164

ANAB ACCREDITED ANAB ACCREDITED

SIGMA-ALDRICH
2631 Soldier Springs Rd. Laramie, Wyoming 82070 USA
307-742-5452
rtstechgroup@sial.com www.sigma-aldrich.com

molecules

MDPI

PCB (129)+(138)+(163)	ug/Kg	608 ± 21.3	2.00	91.2	334 to 882
PCB (153)+(168)	ug/Kg	571 ± 20.0	2.00	85.7	314 to 828
PCB (180)+(193)	ug/Kg	1,160 ± 40.6	2.00	174	638 to 1,682

Sample Information

DESCRIPTION

The sample size provided is 50 g of soil.
The soil has been sterilized to minimize degradation of the sample.
The sample has been sized to 100 mesh.
The sample has been intentionally prepared with an apparent headspace.

PREPARATION INSTRUCTIONS

Mix Prior to use.
Recommended minimum sampling size is 1 gram.

STORAGE

The sample should be stored at Room Temperature. It has been determined to be stable for the duration of the expiration date.
The shelf life of the product was determined by historic stability of similar CRM's. The expiration date may be extended based on stock and popularity upon successful stability testing by a 17025 accredited laboratory.
Stability and shelf life after opening must be determined by the user, taking into account sampling frequency/volume and all local conditions.

1 Certified value - based on the robust mean of round robin interlaboratory study and analytically verified by RTC with associated uncertainties from the preparation and analytical procedures.
2 The standard deviation is the robust statistics standard deviation from the round robin interlaboratory study.
3 Acceptance limits are based on Interlaboratory Study Results. These ranges are recommendations only.
4 Ucrm - Uncertainty values in this document are expressed as Expanded Uncertainty (Ucrm) corresponding to the 95% confidence interval. Ucrm is derived from the combined standard uncertainty multiplied by the coverage factor k, which is obtained from a t-distribution and degrees of freedom. The components of combined standard uncertainty include the uncertainties due to characterization, homogeneity, long term stability, and short term stability (transport). The components due to stability are generally considered to be negligible unless otherwise indicated by stability studies. The mathematical representation of the Ucrm calculation is as follows:

$$u_{CRM} = \sqrt{u^2_{char} + u^2_{bb\ (between)} + u^2_{stab}}$$

5 k: Coverage factor derived from a t distribution table, based on the degrees of freedom of the data set. Confidence interval = 95%

Traceability. The standard was manufactured under an ISO/IEC 17025:2005 certified quality system. The balance used to weigh raw materials is accurate to +/- 0.0001 g and calibrated regularly using mass standards traceable to NIST. All dilutions were performed gravimetrically. Additionally, individual analytes are traceable to NIST SRMs where available and specified above.

Homogeneity. Homogeneity was assessed in accordance with ISO Guide 35. Completed units were sampled using a random stratified sampling protocol. The results of chemical analysis were then compared using a one-way analysis of variance approach as described by TNI EL-V3-2009 Appendix A.2. See instructions for minimum sub-sample size.

THIS PRODUCT WAS DESIGNED, PRODUCED AND VERIFIED FOR ACCURACY AND STABILITY IN ACCORDANCE WITH ISO/IEC 17025:2005 (ANAB Cert AT-1467) and ISO GUIDE 34:2009 (ANAB Cert AR-1478).

Robert O'Brien - QC Supervisor

Mark Pooler - QA Supervisor

Certification Date July 24, 2017
Version 6ZZ-7242017

molecules

MDPI

Ecotoxicological analyses –description of the methodology

Ostracodtoxkit(F)™ (MicroBioTest Inc. Belgium

Ostracodtoxkit(F) test belongs to direct contact tests of chronic toxicity estimation with the use of crustaceans *Heterocypris incongruens*. The direct contact test is done with the application of young bottom crustaceans (*Heterocypris incongruens*), which hatched from cysts during 52 hours (according to the producer's procedure). The test is carried out on 6-hole polystyrene microplates. A standardised nutrient is an algae suspension, which is applied to the suspension in portions of 2 ml. Portions of 10 ostracods are added to each cell of the microplate. 0.5 g of a control soil is applied to the row A, whereas 0.5 g of the soil sample is added to the consecutive rows. Incubation lasts for 6 days in a temperature of 25°C. Results reading consists of estimation of life microforms amount in each hole of the microplate and measurement of their length. Growth inhibition of H. incongruens was calculated as

$$GI = 100 - (\frac{A}{B} * 100)$$

where: A – ostracods length in tested soil, B – ostracods length in reference soil

Microtox® Solid Phase Test (SDI, USA)

Microtox Test created in the USA in 1979 as the first bioindication test, combines typical bioindication and analytic precision. Luminescence bacteria *Vibro fischeri*, which uses about 10% of its metabolism in order to emit light, was used as a bioindicator. In the presence of toxical substances, there is decrease in luminescence and increase in general toxicity of a sample. In an electron transportation system of the bacteria, lucipherasa enzyme (alcane oxygenaze) catalyses oxidation of a reduced substrate (reduced flavine mononucleotide, riboflavin phosphate or flavine adenin dinucleotide) and during this process luminescence, which can be measured with a photometer, takes place. The obtained substrates of this reaction are oxygen and long-chain aldehyde. In the presence of substances that have negative influence on cell metabolism, the decrease in luminescence of bacteria is immediate[1].

Test Microtox®, produced by SDI Company (USA), enables a direct contact of luminescence bacteria *Vibro fischeri* with a wastewater sample, which leads not only to determination of substances dissolved in water, but also to recognition of lipophilicity systems and poorly dissolved systems in water. Lyophilised bacteria *Vibro fischeri* can be stored in a period of a year in temperature of -20°C and then used for testing immediately after suspending in deionised water. A tests with serial dilutions of the wastewater were done according to a standard procedure and results were obtain in Delta Tox analyser. In this analysis, the concentration of wastewater, which reduces the luminescence by 50%, was determined (half maximal effective concentration, EC_{50}). In this presentation a lower of EC_{50} denotes a higher toxicity. To simplify this interpretation, EC_{50} values were converted to Toxicity Units (TU) in which a higher value denotes a higher toxicity according to the following formula.

molecules

$$TU = \frac{1}{EC_{50}} * 100$$

Phytotoxkit™ test (MicroBioTests Inc., Belgium)

Chronic toxicity assessment test *Phytotoxkit™* is based on the evaluation of germination and early growth of plants (root elongation inhibition measurement. Three types of plants selected according to germination rate and root growth rate are used in the test, which allows making a complete determination in 3 days of incubation: monocotyledonous - sorghum *(Sorghum saccharatum)*, and dicotyledonous - cress *(Lepidium sativum)* and white mustard *(Sinapis alba)*. The determination process was carried out using three repetitions for each test plant. The tests were performed on transparent polystyrene test plates. *Incubation conditions:* temperature T = 25°C in darkness, incubation time t = 72 h. *Test reaction:* inhibition of germination and early root growth.

MARA environmental risk test

Evaluation of the data produced with the MARA is assessed by examining the growth inhibitory effect on the pellet formation at each sample dilution. The assessment can be made for each individual species or by looking at the mean software computed value of all the constituent species. Using the latter the toxicity of a sample can be expressed with reference to a threshold value or interval assigned for toxicity classification. For example, an undiluted sample exhibiting an overall inhibitory effect (across 11 species) of o20% is considered to be non-toxic. This is an effective means of obtaining a rapid indication (or assessment) of the results.

In order to provide a comprehensive and optimal assessment utilising the significant feature of the MARA as a multi species test, a determination referred to as the Microbial Toxic Concentration (MTC) is computed. The MTC value is determined as follows

$$MTC = c_{min} \times d^{(P_{tot}/P_0)-1}$$

where c_{min}=lowest concentration in the gradient, P_0=pellet size in the control, d= dilution factor and P_{tot}= six pellet sizes down the concentration gradient.

The MTC is a means of computing a value equivalent to the EC_{50} determination. The computation uses all available points above and below the growth curve in the 0–100% inhibition range[3].

As the MARA plate generates 11 independent values from 11 microbial strains, a test result can be presented either as the mean inhibition value or as the minimum inhibition value if a single numerical value is required. The mean value is probably a more "realistic" value, a common criticism against existing bacterial assays is that the result from one strain only is not representative. On the other hand,

molecules MDPI

the minimum value is a valuable result as it shows at which concentration level the most susceptible species may be affected. However, most of the information gained in a multi-species test is lost if a single value is used for evaluation instead of the whole toxic fingerprint[4].




Figure S3. Scheme of MARA test conduct

AMES (Muta-ChromoPlate™ Kit (EBPI, Canada).

Numerous petroleum contaminants and metabolites created in biodegradation of pollutants include carcinogenic compounds. In order to estimate potentially mutagenic properties of the wastewater, Ames Test was applied. It enables determination of reverse mutations from histidine auxotrophy to prototrophy in mutants, which are test strains of *Salmonella typhimurium*. The test strains of the bacteria can be characterised by various types of mutation in genes, which are responsible for histidine synthesis. Therefore, the presence of histidine in a substratum is required in order to provide the growth of the strains. Generally, TA98 and TA100 strains are used[5]. They include proper mutations consisting of so-called reading frame displacement, pair base substitution and two mutagenes (2-nitro-fluoren and sodium azide). The test strains are mutants, which are unable to histidine synthesis. A mutagenous substance causes reversion in mutation and histidine synthesis can take place, which results in appearance of revertants colonies in the histidine-free basis.

In order to lead Ames Test, a high-tech microplate AMES MPF test was applied. It can be used for detection of genotoxic activity of water contaminated with petroleum substances and metabolites, which are the effects of biodegradation. Delivered microforms of T-98 and

molecules

T-100 *Salmonella typhimurium* undergo strict quality control (genotype and phenotype). The test results in colour reaction, where yellow means reverse mutation, whereas scarlet – the lack of mutation.




Figure S4. Scheme of Ames Test conduct

Petroleum contaminants from the samples were extracted with the use of dichloromethane, which was then vaporised and the obtained deposit was dissolved in dimethylosulfoxide (DMSO)[6]. As an activator, S9 microsomal fraction from a rat liver was used. The role of the activator is to transform the tested substance, which leads to disclosure of its potentially mutagenous properties.